



# Forecasting the Mediterranean Sea marine heatwave of summer 2022

Ronan McAdam, Giulia Bonino, Emanuela Clementi, and Simona Masina

CMCC Foundation – Euro-Mediterranean Center on Climate Change, Bologna, Italy

**Correspondence:** Ronan McAdam (ronan.mcadam@cmcc.it)

Received: 24 July 2023 – Discussion started: 28 August 2023

Revised: 11 January 2024 – Accepted: 6 February 2024 – Published: 30 September 2024

**Abstract.** Early warning of marine heatwaves requires short-term forecasts to provide precise information on timings, local-scale coverage, and intensities of coming events. Here, we describe our successful efforts to track the onset, peak, and decay of the Mediterranean Sea marine heatwave of summer 2022 with the Copernicus MedFS short-term (10 d) forecast system. First, we show that the 2022 event eclipses the economically and ecologically damaging event of 2003 in terms of marine heatwave (MHW) activity (a measure of intensity and duration). Forecasts of MHW area and activity provide a means of basin-wide validation, highlighting the capability of MedFS to capture regional behaviour. On local scales, we found that the MHW occurrence in the Ligurian Sea and Gulf of Taranto, two regions of economic and ecological importance, was also reliably forecast. Encouragingly, we note that the forecast has demonstrated skill in capturing not just the season-long MHW cycle but also breaks in MHW persistence and abrupt changes in local activity. Subseasonal forecasts do not yet demonstrate the capacity to predict MHW response to short-lived weather patterns, but this study confirms that short-term forecasts, at least in the Mediterranean Sea, can fill this gap.

## 1 Introduction

Disease outbreaks, mass mortality events, and the redistribution of species induced by marine heatwaves (MHWs) lead to economic losses to fisheries and aquaculture farms, as well as hampering conservation efforts (Barbeaux et al., 2020; Smith et al., 2021; Garrabou et al., 2022; Smith et al., 2023). The need to prepare for and mitigate these MHW-induced impacts has driven developments in understanding the drivers and predictability of MHWs (Holbrook et al., 2019; Rodrigues et al., 2019; Amaya et al., 2020; Sen Gupta et al., 2020; Li et al., 2020; Schlegel et al., 2021) and in quantifying the skill of forecasts of MHWs (Benthuisen et al., 2021; Jacox et al., 2022; McAdam et al., 2023). The Mediterranean Sea is a particular “hot spot” for MHWs, with much literature documenting the increases in intensity, duration, frequency, and impacts (Darmaraki et al., 2019; Ibrahim et al., 2021; Juza et al., 2022; Dayan et al., 2023). Despite this, there is currently less information on forecasting capability and event predictability in the Mediterranean region than in others (e.g. the North Pacific: Jacox et al., 2019; de Boises-

son et al., 2022). With marine services projected to play an ever-increasing role in global sustainability and economic security (Rayner et al., 2019), early-warning systems of heat extremes can aid their planning and day-to-day management (Hartog et al., 2023).

While interannual variability of MHW occurrence and characteristics is derived from ocean warming and preconditioning (de Boissezon et al., 2022), ENSO (Jacox et al., 2022), and atmospheric teleconnections (e.g. Rossby wave trains; Rodrigues et al., 2019), short-lived atmospheric processes and weather systems can disrupt MHW persistence or halt their continuation completely (Benthuisen et al., 2021). The definition of MHWs assumes persistent conditions are harmful to marine life if the duration is 5 d or longer (Hobday et al., 2016), although this number is quite arbitrary and in principle should be species-dependent. The average duration of MHWs across most of the global ocean, as well as in the Mediterranean Sea, falls within the definition of short-term forecasting (< 2 weeks) (Oliver et al., 2021; Dayan et

**Table 1.** Product table.

| Product ref. no | Product ID and type  | Data access                                  | Documentation  |
|-----------------|--|--|--|
| 1               | SST_MED_SST_L4_REP_OBSERVATIONS_010_021;<br>Satellite observations | EU Copernicus Marine Service Product (2023a) | PUM: Pisano et al. (2023a)<br>QUID: Pisano et al. (2023b)  |
| 2               | MEDSEA_ANALYSISFORECAST_PHY_006_013;<br>Numerical models           | EU Copernicus Marine Service Product (2023b) | PUM: Lecci et al. (2023)<br>QUID: Goglio et al. (2023)   |
| 3               | MED_MULTIYEAR_PHYS_006_004;<br>Numerical models                    | EU Copernicus Marine Service Product (2022)  | PUM: Lecci et al. (2022)<br>QUID: Escudier et al. (2022)   |
| 4               | ECMWF Integrated Forecast System (IFS) Forecast & Analysis         | Copernicus Climate Changes Service           | <a href="https://www.ecmwf.int/en/forecasts/datasets/set-i#I-i-a_fc">https://www.ecmwf.int/en/forecasts/datasets/set-i#I-i-a_fc</a><br>(last access: 13 February 2022) |
| 5               | ERA5   | Copernicus Climate Changes Service           | Hersbach et al. (2023)   |

al., 2023). A short-term view of MHWs is therefore crucial to understanding their predictability and their impacts.

Short-term forecasting of MHWs has a range of potential roles in marine activities. While some contingency plans for extreme heat events in the aquaculture and fishing industries require several months notice (e.g. relocating or switching species), others should be performed at the latest possible moment in order to avoid or minimize losses (e.g. early harvesting or cooling of farm water) (Holsman et al., 2019; Galappaththi et al., 2020). In these cases, accurate information on daily timescales is crucial. Short-term forecasts are also useful for marine protected areas (MPAs), allowing them to prepare to monitor ecosystem damage (e.g. coral bleaching) and recovery, which in turn helps assess the effectiveness of their conservation efforts (McLeod et al., 2009). Forecasts of SST can also, in theory, be coupled to distribution models to forecast changes in species habitat for highly mobile species (Abrahms et al., 2019). “Early warnings” are a key means of climate resilience for marine services (Galappaththi et al., 2020); an assessment of their ability to track MHWs will contribute to further uptake by these services and unlock potential socio-economic benefits.

During the summer of 2022, the Copernicus Marine Service Mediterranean Physical Forecasting system (MedFS) was employed to monitor and forecast sea surface temperature (SST) increases which eventually evolved into a record-breaking MHW for the region. MedFS has already demonstrated skill in detecting past extreme events in the Mediterranean Sea: the “aqua alta” flooding in Venice in 2019 (Giesen et al., 2021), Medicane Ianos (Clementi et al., 2022), and Storm Gloria (Álvarez-Fanjul et al., 2022) in 2020 provides evidence that the forecast system has the ability to capture a wide range of concurrent conditions (e.g. high surface air temperatures, moisture, atmospheric instability for medicanes; Cavicchia et al., 2014). Here, we provide a basin-wide description of the event and demonstrate the ability of

MedFS to accurately predict many facets of the event (e.g. the onset, spread, persistence, and decay). First, we introduce the high-resolution regional forecast system and the satellite-derived SST data used to identify MHWs. Then, the record-breaking characteristics (intensity, geographic extent) of the 2022 event are described. We demonstrate the system’s ability to predict the MHW spread across the basin and daily temperature variability in regions of key economic and ecological importance. Finally, we explore the potential role of short-term forecasting in the early warning of MHWs compared to other forecasting timescales.

## 2 Dataset and methods

Here, MHWs are detected with a  $0.05^\circ$  resolution reprocessing of a blend of satellite-derived products provided by the ESA Climate Change Initiative (CCI) and the Copernicus Climate Change Service (C3S) initiatives, including AVHRR Pathfinder dataset version 5.3 to increase the input observation coverage (Table 1, product ref. no. 1). The dataset provides daily SST of the Mediterranean Sea from 1 January 1982 to the present (currently up to 6 months before real time).

The Mediterranean Near Real Time Analysis and Forecast is a 3D coupled hydrodynamic–wave modelling system implemented at  $1/24^\circ$  ( $\sim 4$  km) horizontal spatial resolution, which produces analysis and 10 d forecasts of the main ocean essential variables (Table 1, product ref. no. 2). The analysis system assimilates satellite sea level anomalies as well as in situ temperature and salinity observations and nudges SST towards an ultrahigh-resolution satellite product. The same model framework is used to provide a multi-decadal reanalysis of the ocean, extending from 1987 to the present (Table 1, product ref. no. 3). Forecasts are made daily; once a week (on Tuesdays) an analysis is used to initialize forecasts, while on other days a hindcast is used. A schematic of the provision of

forecast and analysis data is found in the Quality Information Document (QUID) and Product User Manual (PUM) of the product.


Forecasts of 2 m temperature (T2M) and wind speed are obtained from the European Centre for Medium-Range Weather Forecasts (ECMWF) operational forecast and analysis (Table 1, product ref. no. 4) distributed by the Italian National Meteorological Service (USAM/CNMA). Variables are available at 1-hourly resolution for the first 3 d of forecast, 3-hourly for the following 3 d, and 6-hourly for the following 4 d. The horizontal resolution is  $0.1^\circ$ . The T2M anomaly is calculated using the same variables in the ERA5 reanalysis over the period 1987–2021.

MHWs are defined as SSTs which persist above the 90th percentile for 5 d or longer (Hobday et al., 2016). Here, the 90th percentile threshold corresponds to the 33-year baseline period of 1987–2021 calculated individually for satellite-derived and model-derived data and smoothed with an 11 d moving window. MHWs in MedFS are defined relative to the climatology of the physical reanalysis. Although there are benefits of detrending SST prior to detecting MHWs (Amaya et al., 2023), we chose not to detrend in order to present the true values of temperature because they are of more relevance to species impacts (e.g. Galli et al., 2017).

We use the MHW activity as a means to describe the event on a basin scale and to validate the forecast ability to capture the spatial scale of the event. Previously, activity has been defined as the product of event intensity, duration, and area over a target period (Simon et al., 2022). Here, in order to study basin-wide spread at daily resolution, we define activity as the sum of the intensity over the area undergoing an MHW in the Mediterranean Basin. We assume that all MHW activity in the basin corresponds to the same event, unlike more novel methods of MHW tracking which employ spatial clustering (Bonino et al., 2023). Nonetheless, the activity metric used here identifies very similar phases of MHW activity during the 2003 event as the more advanced clustering method (Fig. 1a; Bonino et al., 2023).

### 3 Results


First, we describe the event on a basin scale using satellite observations. We remind the reader that references to specific dates are for indication only, as the precise timings of peaks and onsets may differ when using different datasets and climatologies. In 2022, the onset of summer MHW conditions began in mid-May; by 23 May, 35 % of the area of the Mediterranean Sea was already experiencing MHW conditions (Fig. 1c). Maps of temperature anomaly confirm that the onset occurred mostly in the western regions and the Adriatic Sea (Fig. 2a). The geographic extent of MHW extended into the central and eastern parts of the basin (e.g. Fig. 2b), and MHW area remained above a third of the total basin area until the decay at the end of September. Peak area



**Figure 1.** MHW activity across the Mediterranean Sea. (a) MHW activity defined by reprocessed satellite observations for 2022 and the 3 previous record years according to Simon et al. (2022). (b) Comparison between satellite observations and forecasts of 2022 MHW activity. (c) Area of the Mediterranean Sea experiencing an MHW (as a percentage of total basin area). Activity is defined as the sum of the intensity over the area undergoing an MHW. Shown here are the first 8 d of forecasts initiated on Tuesdays. Forecast start dates are shown by the blue dots.

(70 %) was reached on 6 June, while notable peaks of activity occurred later on 29 June, 6 July, and 27 July (Fig. 1a). The peak temperature anomaly of  $6.45^\circ\text{C}$  (above the 1987–2019 average) was reached in the Gulf of Lion on 18 July.

Prior to 2022, the MHWs of the summers of 2003, 2015, and 2018 had been found to have the highest activities on record (using a slightly different definition of activity, but which is still based on intensity and duration; Simon et al., 2022). Here, we find that the activity in 2022 clearly eclipses that of 2015 and 2018 in terms of both maxima and persistence of activity (Fig. 1a). Though the summer of 2003 reached similarly high peaks of activity (twice, in mid-June and at the end of August), the total activity during the summer (defined as the area under the curve) is lower ( $82 \times 10^6 \text{ }^\circ\text{C km}^2$ ) than for 2022 ( $139 \times 10^6 \text{ }^\circ\text{C km}^2$ ). While



**Figure 2.** Snapshots of SST anomalies and MHW occurrence during the different stages of the 2022 MHW. (a, c, e) Reprocessed satellite observations. (b, d, f) Forecasts with a lead time of 4 d. Areas in which SST is above the 90th percentile threshold are indicated by the dark red contour. The day of 13 May highlights the MHW onset, 9 July highlights the peak activity, and 26 August highlights the (first) decay. Regions used in Figs. 3 and 4 are highlighted: Ligurian Sea (magenta), Gulf of Taranto (yellow), and the western Mediterranean (blue).

in 2003 the MHW activity returned to zero in late May and mid-July, in 2022 it persisted throughout the summer above at least  $0.5^{\circ}\text{C km}^2$  each day. Using the total activity as a measure, the summer of 2022 now holds the record for MHW activity.

Using the MHW activity provides an efficient, if not complete, means of validating MedFS on the basin scale. It is important to remember that activity time series cannot identify where and when MHWs are occurring (we study forecast ability in specific regions later). Here, we show both the activity (Fig. 1b) and the area (Fig. 1c) predicted to infer whether forecast inaccuracies are caused by an inability to capture the geographic extent or the temperature intensity. Overall, we find that MedFS was able to forecast the evolution of basin-wide MHW activity (Fig. 1b). In particular, we highlight the accurate predictions of the timings of the May onset, the various peaks throughout the summer, the two stages of the decay, and the September rebound. On several occasions, MHW activity rapidly increases, often doubling or tripling over the period of less than a week; such increases in activity are predicted by the forecasts in mid-May, early June, mid-June (twice), and mid-July. Timings of declines in activity are also correctly forecast on all occasions, in some cases even with lead times of 5 d or more (e.g. early June). Similar capabilities are found for the forecasts of the area of MHW conditions. Ability to capture both the activity and the area implies accuracy in capturing the intensity as well, although this analysis does not yet determine the geographic distribution of MHW intensity.

There are indeed forecast inaccuracies to highlight. Firstly, there are instances of peaks of activity being overestimated

(e.g. by up to a third in early June) and of false alarms about growth being raised (e.g. early August; Fig. 1b). Overestimations of both activity and area occur throughout the summer, except for the onset in May. Then, there are instances in which MHW area tendencies follow the activity but are overestimated (late July to early August); given that activity continues to increase while area decreases (e.g. early September), there is an implied overestimation of the sea surface temperatures. Elsewhere, we see fluctuations in activity on daily timescales which dominate the longer-term growth tendencies (e.g. during the growth period beginning at the end of June). Throughout the summer of 2022, we see various examples of the activity forecasts being unable to detect this higher-frequency variability. Variability in area, on the other hand, is lower and the forecasts generally follow the observed tendencies. In summary, the forecasts sometimes persist or increase temperature anomalies for too long, suggesting that they fail to capture sporadic cooling. Candidate drivers for short-term cooling mechanisms not captured by the forecasts include cloud cover changes or winds.

It is important to also consider the ability to capture the spread of MHW occurrence. The geographical distribution of intensity in key phases of the MHW life cycle in forecasts agrees well with observations (Fig. 2). During the onset, forecasts capture the basin-wide patterns, with MHW occurrence at this stage correctly forecast in the Tyrrhenian Sea, Gulf of Lion, and parts of the Adriatic Sea. The spread of the MHW conditions during the peak was correctly predicted to cover the southern part of the Alboran Sea, the Ionian, and southern regions of the Levantine Basin. Meanwhile, the Aegean Sea was predicted to be shielded from MHWs and instead expe-



rience cold anomalies, most likely caused by cooling related to the Etesian winds (Poupkou et al., 2011). Lastly, the first decay phase at the end of August produced very inhomogeneous MHW conditions across the basin. This “patchiness”, indicative of local-scale processes acting to cool the ocean such as increased cloud cover or weak winds, was indeed predicted, but how well the forecast matches observations depends greatly on the local regions of interest and the exact day and lead time considered. Although it is not possible to draw rigorous conclusions from snapshots, the accuracy of basin-wide occurrence (Fig. 1b) suggests that forecast ability to capture MHW patterns and spread was generally high across the entire summer.

While basin-scale analysis allows an overview of forecasting skill, local-scale testing is imperative as forecasting tools are expected to be used for local-scale analysis (Dayan et al., 2023). Here, we also provide MHW forecasts for two key areas of maritime activity in the Mediterranean Sea: the Ligurian Sea and the Gulf of Taranto (Fig. 3). Each region experienced MHW conditions at different times during the summer, and in each case the forecasts accurately predicted the onset, persistence, intensity, and decays. The Ligurian Sea, bordered by Italy and France, is a crucial location for marine conservation; it doubles as a marine protected area (the Pelagos Sanctuary for Mediterranean Marine Mammals) which is home to unique species of fin whales and striped dolphins, amongst other species (Notarbartolo-de-Sciara et al., 2008). The Ligurian Sea experienced 115 d of MHWs throughout the summer, and temperature anomalies reached a maximum of 4.46 °C above the 1987–2019 average at the end of July, coinciding with the peak temperature of the summer (28.74 °C). The forecasts of SST were highly accurate; the root mean squared difference (RMSD) of the forecasts shown in Fig. 3a (across lead times) was 0.12 °C, lower than the reported RMSDs averaged over the entire Mediterranean Sea between the analysis and independent observations (Goglio et al., 2023). For an indication of forecast reliability, we highlight the false alarms (MHW days forecast but not observed) and misses (MHW days observed but not forecast). First we note that in the 89 % of days the correct conditions are forecast, with a few sporadic exceptions. For example, the forecast made on 10 May captured the sharp rise in SST but not the MHW conditions at the end of the week. However, reducing the lead time (i.e. checking forecasts made on 12 or 13 May) correctly forecast the MHW state.

The Gulf of Taranto, situated in the northern Ionian Sea, is one of the most productive areas of shellfish (mussels) farming in Italy (Prioli, 2004), but there are no data yet on MHW-induced mass mortality or economic loss in this region (Garrabou et al., 2022). Unlike the Ligurian Sea, the Gulf of Taranto experienced three short but intense periods of MHW occurrence in June and July, adding up to 61 d of MHWs in total. The peak temperature anomaly was 4.76 °C on 6 June, though peak temperatures occurred later in the season. As in the Ligurian Sea, the forecasts were highly ac-


**Table 2.** Root mean square differences of forecasts of summer 2022 MHW activity and atmospheric conditions (Fig. 4). Values in parenthesis are RMSD values normalized by standard deviation over the summer. Differences in MHW activity and area are relative to reprocessed satellite observations, while differences in T2M anomaly and wind speed are relative to ECMWF analysis. Each column corresponds to a different lead time.

| RMSD<br>(normalized) | Lead: 1 d   | Lead: 4 d    | Lead: 7 d    |
|----------------------|-------------|--------------|--------------|
| MHW activity         | 0.16 (0.48) | 0.20 (0.59)  | 0.28 (0.82)  |
| MHW area             | 8.88 (0.33) | 11.65 (0.43) | 16.50 (0.61) |
| T2M anomaly          | 0.18 (0.21) | 0.31 (0.38)  | 0.52 (0.62)  |
| Wind speed           | 0.22 (0.18) | 0.52 (0.42)  | 0.94 (0.76)  |

curate (demonstrating an RMSD of 0.08 °C). Regarding the reliability of MHW forecasts, the continuation of the start of the heatwave in early May was missed by the forecast of 17 May, while the forecast of 2 August missed several days of MHW occurrence. However, in both cases, SST increases were predicted. None of the forecasts shown in Fig. 3 raised false alarms.

So far we have studied the accuracy of the entire forecast period, but, in some applications, it might be necessary or of more interest to have a specific warning time (e.g. 4 d). As MedFS produces forecasts every day, we now study forecast accuracy for the summer of 2022 at different lead times; we focus on the western Mediterranean Sea, as opposed to the entire basin, in order to investigate local forcings as reasons for poor skill (Fig. 4). Table 2 quantifies the error of forecasts over the western Mediterranean Sea. The overestimation of MHW activity in July and August occurred in forecasts with lead time of 1 d. In many instances, lead time 1 and lead time 4 are similarly far from the observed values, while lead time 7 further overestimates the peaks in activity (Table 2). MHW area, on the other hand, while predicted more accurately than activity on all lead times, is typically underestimated by the forecasts. This implies an overestimation of SST (i.e. MHW intensity) during the activity peaks. The RMSD normalized by the standard deviation indicates where errors fall within the range of natural variability (normalized RMSD < 1); in all lead times, this is true for both area and activity, suggesting their skill is similar to atmospheric variables (Table 2).

The decreases in skill with lead time can partly be explained by the decrease in skill of the ECMWF atmospheric forecasts used to force MedFS. T2M and wind speed correlate strongly and significantly with the MHW activity (correlation values of 0.89 and 0.50 with the ECMWF analysis respectively), evidencing their role in MHW formation. Errors of forecasts of T2M and wind speed grow with time but do not exceed natural variability at lead time 7 (Table 2). In the first half of the summer, forecasts at lead time 7 of both T2M and wind speed are frequently out of phase with the observed changes. In fact, the underestimations of MHW area



**Figure 3.** Time series of SST and MHW occurrence in summer 2022. Orange shading highlights MHW occurrence in reprocessed satellite observations. Forecast start dates are shown by the blue dots. Definitions of the Ligurian Sea (a) and Gulf of Taranto (b). Note that the climatology lines (red) correspond to the satellite data, not to the model output (analysis and forecasts). Crosses correspond to misses (red) and false alarms (blue) in the forecast output.


in this period occur simultaneously with underestimations of T2M. For example, the underestimation of MHW area at the end of May, by an area of roughly 30% of the western Mediterranean, corresponds to overestimations of wind speed by up to  $1 \text{ m s}^{-1}$  and temperature anomalies roughly  $1 \text{ }^{\circ}\text{C}$  weaker than observed. However, the overestimation of activity in July and August, found to be linked to overestimations of SST, does not correspond to overestimations of T2M, implying that other phenomena are not well represented. It should be noted that the use of area-averaged atmospheric variables may hide sub-regional-scale processes which impact the MHW location and intensity.

#### 4 Discussion and summary

The MHW of summer 2022 in the Mediterranean Sea was record-breaking, eclipsing 2003 in terms of basin-wide activity (defined as the integral of intensity, duration, and area). Other contributions to the Ocean State Report 8 also define the MHW of 2022 as a record-breaking event using other definitions (e.g. local SST records). Here, we provide a basin-wide view of the MHW conditions. The Copernicus Mediterranean Physical forecasting system was used to track this event, serving as the first validation of MHW prediction for this system. Forecasts captured the full life cycle of the MHWs several days in advance: onset (mid-May) in the western part of the basin, spread into the Adriatic and

Ionian Sea, sporadic local-scale occurrences in the Levantine Basin, persistence of peak conditions throughout July and August, breaks in MHW persistence and abrupt changes in local occurrence, and the gradual decay (September). The forecasts also identified regions shielded from MHWs, e.g. during cooling in the Aegean Seas. Subseasonal forecasts do not yet demonstrate the capacity to predict MHW response to weather patterns (Benthuisen et al., 2021), but this study confirms that short-term forecasts, at least in the Mediterranean Sea, can fill this gap.

A full analysis of potential drivers and attribution of forecast skill to certain processes was outside the scope of this study, but the dependence on accurate atmospheric forcings (here provided by ECMWF forecasts) has been shown to be crucial for accurate forecasts of the 2022 event. Unlike the other extreme events, the common drivers of MHWs in the Mediterranean are yet to be identified. The MHW of 2022, as well as the concurrent and record-breaking atmospheric heatwave which occurred over western Europe, appears to be linked to the northward extension of the subtropical ridge (Barriopedro et al., 2023), while model studies have suggested that mid-latitude MHWs in summer typically arise from reduced ocean heat loss to the atmosphere and reduced vertical diffusion (Vogt et al., 2022). Here, the decay in skill of MHW forecasts matches the decay in skill of key atmospheric conditions (T2M and wind speed). Erroneous forecasts of early summer in the western Mediterranean Sea are



**Figure 4.** Effect of lead time and atmospheric forcings on forecasts of MHW activity and area. Comparison between reprocessed satellite observations and forecasts of 2022 MHW activity (a) and area (b). Each forecast time series corresponds to a different lead time (i.e. how many days in advance the forecast was made). Forecasts of MHW activity were calculated for forecasts initiated every day; the lead time from each forecast was extracted to construct the time series. Area-averaged 2 m temperature anomaly (c) and wind speed (d) from the ECMWF analysis and forecasts used to force the MedFS system. All time series correspond to the western Mediterranean Sea (Fig. 2).

explained by inaccurate forecasts of these atmospheric conditions. Peak summer conditions, such as the overestimation of MHW activity, are not yet understood, meaning further studies of short-term forecasting of MHWs are necessary.

The timescale of forecasting determines the information that can be provided and the type of response to that information. Here, we make the case for using short-term forecasting in MHW tracking tools and studies. Seasonal forecasting informs management decisions and contingency plans, while subseasonal forecasting can update these plans (White et al., 2017). Short-term forecasting, on the other hand, can then be used to determine the precise timings of events and in-

struct users on when to implement urgent response actions. Longer-term forecasts are typically global in scale and have a relatively low model resolution, while short-term forecasting centres, benefitting from the reduced timescale, can put more computational power towards regional-scale forecasting at a finer scale more relevant to stakeholders. In principle, for MHWs, this means the following: seasonal forecasts provide forewarning of extreme summer temperatures (e.g. seasonal averages above the 90th percentile, identification of ocean basins affected). Sub-seasonal forecasts then update this to provide forewarning of MHW occurrence (e.g. daily temperatures persisting above 90th percentile, greater detail

on geographic spread). Finally, short-term forecasts can provide key details such as the start date, onset rate, and breaks in occurrence on a local to regional scale. Currently, more effort is being devoted to seasonal forecasting of MHWs (de Burgh-Day et al., 2022; Liu et al., 2018; Jacox et al., 2022). With the level of accuracy for local-scale MHW indicators shown here, such tools should be complemented with daily short-term updates.

In particular, we found that the MHW occurrence in the Ligurian Sea and Gulf of Taranto, two regions of economic and ecological importance, was also reliably forecast. There is, though, a need to include subsurface temperatures or heat content to report MHWs occurring at depth (Dayan et al., 2023; McAdam et al., 2023). For example, caged fish have been observed to avoid the top of cages when surface temperatures increase (Gamperl et al., 2021), meaning truly stakeholder-relevant tracking tools need a 3D view. The near-real-time analysis, as well as the forecast system, provides 3D temperatures and can track subsurface propagation of MHWs (unlike satellite observations). The MHW record in the analysis aligns exceptionally well with satellite observations for the two target regions shown (Fig. 3), suggesting a high level of accuracy (the same is found for the basin-wide MHW activity; not shown). However, a subsurface validation with in situ data should be performed in the near future, before using the analysis and forecast to track subsurface MHWs.

**Code availability.** Codes used to analyse data and produce figures in this study are available at <https://github.com/RJMcAdam> (last access: 8 April 2024; <https://doi.org/10.5281/zenodo.10944086>, McAdam et al., 2024).

**Data availability.** The underlying data can be found in the product table (Table 1).

**Author contributions.** RM, GB, SM, and EC conceived the study. RM and GB performed the analysis and prepared the figures. RM wrote the paper. GB, SM, and EC contributed to the interpretation of the results and to the paper writing. RM, GB, SM, and EC reviewed the paper.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

**Acknowledgements.** We acknowledge the CMCC Foundation for providing computational resources.

**Review statement.** This paper was edited by Pierre Brasseur and reviewed by Francisco Pastor and one anonymous referee.

## References

- Abrahms, B., Welch, H., Brodie, S., Jacox, M. G., Becker, E. A., Bograd, S. J., Irvine, L. M., Palacios, D. M., Mate, B. R., and Hazen, E. L.: Dynamic ensemble models to predict distributions and anthropogenic risk exposure for highly mobile species, *Divers Distrib.*, 25, 1182–1193, <https://doi.org/10.1111/ddi.12940>, 2019.
- Álvarez Fanjul, E., Pérez Gómez, B., Alfonso Alonso-Muñoyerro, M. D., Lorente Jiménez, P., García Sotillo, M., Lin-Ye, J., Aznar Lecocq, R., Ruiz Gil de la Serna, M., Pérez-Rubio, S., Clementi, E., and Coppini, G.: Western Mediterranean record-breaking storm Gloria: An integrated assessment based on models and observations, in: Copernicus Ocean State Report, Issue 6, *J. Oper. Oceanogr.*, 15, s151–s159, <https://doi.org/10.1080/1755876X.2022.2095169>, 2022.
- Amaya, D. J., Miller, A. J., Xie, S. P., and Kosaka, Y.: Physical drivers of the summer 2019 North Pacific marine heatwave, *Nat. Commun.*, 11, 1903, <https://doi.org/10.1038/s41467-020-15820-w>, 2020.
- Amaya, D. J., Jacox, M. G., Fewings, M. R., Saba, V. S., Stuecker, M. F., Rykaczewski, R. R., Ross, A. C., Stock, C. A., Capotondi, A., Petrik, C. M., and Bograd, S. J.: Marine heatwaves need clear definitions so coastal communities can adapt, *Nature*, 616, 29–32, 2023.
- Barbeaux, S. J., Holsman, K., and Zador, S.: Marine heatwave stress test of ecosystem-based fisheries management in the Gulf of Alaska Pacific cod fishery, *Front. Mar. Sci.*, 7, 703, <https://doi.org/10.3389/fmars.2020.00703>, 2020.
- Barriopedro, D., García-Herrera, R., Ordóñez, C., Miralles, D. G., and Salcedo-Sanz, S.: Heat waves: Physical understanding and scientific challenges, *Rev. Geophys.*, 61, e2022RG000780, <https://doi.org/10.1029/2022RG000780>, 2023.
- Benthuisen, J. A., Smith, G. A., Spillman, C. M., and Steinberg, C. R.: Subseasonal prediction of the 2020 Great Barrier Reef and Coral Sea marine heatwave, *Environ. Res. Lett.*, 16, 124050, <https://doi.org/10.1088/1748-9326/ac3aa1>, 2021.
- Bonino, G., Masina, S., Galimberti, G., and Moretti, M.: Southern Europe and western Asian marine heatwaves (SEWA-MHWs): a dataset based on macroevents, *Earth Syst. Sci. Data*, 15, 1269–1285, <https://doi.org/10.5194/essd-15-1269-2023>, 2023.
- Cavicchia, L., von Storch, H., and Gualdi, S.: A long-term climatology of medicanes, *Clim. Dynam.*, 43, 1183–1195, 2014.
- Clementi, E., Korres, G., Cossarini, G., Ravdas M., Federico, I., Goglio, A. C., Salon, S., Zacharioudaki, A., Pattanaro, M., and Coppini, G.: The September 2020 Medicanes Ianos predicted by the Mediterranean Forecasting systems, in: Copernicus Ocean State Report, Issue 6, *J. Oper. Oceanogr.*, 15, s168–s175, <https://doi.org/10.1080/1755876X.2022.2095169>, 2022.



- Darmaraki, S., Somot, S., Sevault, F., and Nabat, P.: Past variability of Mediterranean Sea marine heatwaves, *Geophys. Res. Lett.*, 46, 9813–9823, 2019.
- Dayan, H., McAdam, R., Juza, M., Masina, S., and Speich, S.: Marine heat waves in the Mediterranean Sea: An assessment from the surface to the subsurface to meet national needs, *Front. Mar. Sci.*, 10, 142, <https://doi.org/10.3389/fmars.2023.1045138>, 2023.
- de Boissesson, E., Balmaseda, M., Mayer, M., and Zuo, H.: Monitoring and predictions of the series of marine heatwave events impacting the Northeast Pacific in 2020, in: Copernicus Ocean State Report, Issue 6, *J. Oper. Oceanogr.*, 15, s168–s175, <https://doi.org/10.1080/1755876X.2022.2095169>, 2022.
- de Burgh-Day, C. O., Spillman, C. M., Smith, G., and Stevens, C. L.: Forecasting extreme marine heat events in key aquaculture regions around New Zealand, *Journal of Southern Hemisphere Earth Systems Science*, 72, 58–72, <https://doi.org/10.1071/ES21012>, 2022.
- Escudier, R., Clementi, E., Nigam, T., Aydogdu, A., Fini, E., Pistoia, J., Grandi, A., and Miraglio, P.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea Physics Reanalysis, MED-SEA\_MULTIYEAR\_PHY\_006\_004, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-004.pdf> (last access: 23 June 2023), 2022.
- EU Copernicus Marine Service Product: Mediterranean Sea Physics Reanalysis, Mercator Ocean International [data set], [https://doi.org/10.25423/CMCC/MEDSEA\\_MULTIYEAR\\_PHY\\_006\\_004\\_E3R1](https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_PHY_006_004_E3R1), 2022.
- EU Copernicus Marine Service Product: Mediterranean Sea – High Resolution L4 Sea Surface Temperature Reprocessed, Mercator Ocean International [data set], <https://doi.org/10.48670/moi-00173>, 2023a.
- EU Copernicus Marine Service Product: Mediterranean Sea Physics Analysis and Forecast, Mercator Ocean International [data set], [https://doi.org/10.25423/CMCC/MEDSEA\\_ANALYSISFORECAST\\_PHY\\_006\\_013\\_EAS7](https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS7), 2023b.
- Galappaththi, E. K., Ichien, S. T., Hyman, A. A., Aubrac, C. J., and Ford, J. D.: Climate change adaptation in aquaculture, *Rev. Aquacult.*, 12, 2160–2176, <https://doi.org/10.1111/raq.12427>, 2020.
- Galli, G., Solidoro, C., and Lovato, T.: Marine heat waves hazard 3D maps and the risk for low motility organisms in a warming Mediterranean Sea, *Front. Mar. Sci.*, 4, 136, <https://doi.org/10.3389/fmars.2017.00136>, 2017.
- Gamperl, A. K., Zrini, Z. A., and Sandrelli, R. M.: Atlantic salmon (*Salmo salar*) cage-site distribution, behavior, and physiology during a Newfoundland heat wave, *Front. Physiol.*, 12, 719594, <https://doi.org/10.3389/fphys.2021.719594>, 2021.
- Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Ponti, M., Schlegel, R., Bensoussan, N., Turicchia, E., Sini, M., Gerovasileiou, V., Teixido, N., Mirasole, A., Tamburello, L., Cebrian, E., Rilov, G., Ledoux, J.-B., Souissi, J. B., Khamassi, F., Ghanem, R., Benabdi, M., Grimes, S., Ocaña, O., Bazairi, H., Hereu, B., Linares, C., Kersting, D. K., la Rovira, G., Ortega, J., Casals, D., Pagès-Escalà, M., Margarit, N., Capdevila, P., Verdura, J., Ramos, A., Izquierdo, A., Barbera, C., Rubio-Portillo, E., Anton, I., López-Sendino, P., Díaz, D., Vázquez-Luis, M., Duarte, C., Marbà, N., Aspillaga, E., Espinosa, F., Grech, D., Guala, I., Azzurro, E., Farina, S., Gambi, M. C., Chimentì, G., Montefalcone, M., Azzola, A., Pulido Mantas, T., Frascchetti, S., Ceccherelli, G., Kipson, S., Bakran-Petricioli, T., Petricioli, D., Jimenez, C., Katsanevakis, S., Kizilkaya, I. T., Kizilkaya, Z., Sartoretto, S., Elodie, R., Ruitton, S., Comeau, S., Gattuso, J. P., and Harmelin, J.-G.: Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea, *Glob. Change Biol.*, 28, 5708–5725, <https://doi.org/10.1111/gcb.16301>, 2022.
- Giesen, R., Clementi, E., Bajo, M., Federico, I., Stoffelen, A., and Santoleri, R.: The November 2019 record high water levels in Venice, Italy, in: Copernicus Ocean State Report, Issue 5, *J. Oper. Oceanogr.*, 14, s156–s162, <https://doi.org/10.1080/1755876X.2021.1946240>, 2021.
- Goglio, A. C., Clementi, E., Grandi, A., Moulin, A., Giurato, M., Aydogdu, A., Pistoia, J., Miraglio, P., Mariani, A., and Drudi, M.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea Physics Analysis and Forecast, MED-SEA\_ANALYSISFORECAST\_PHY\_006\_013, Issue 2.4, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-013.pdf> (last access: 14 January 2024), 2023.
- Hartog, J. R., Spillman, C. M., Smith, G., and Hobday, A. J.: Forecasts of marine heatwaves for marine industries: Reducing risk, building resilience and enhancing management responses, *Deep-Sea Res. Pt. II*, 209, 105276, <https://doi.org/10.1016/j.dsr2.2023.105276>, 2023.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., Benthuisen, J. A., Burrows, M. T., Donat, M. G., Feng, M., and Holbrook, N. J.: A hierarchical approach to defining marine heatwaves, *Prog. Oceanogr.*, 141, 227–238, 2016.
- Holbrook, N. J., Scannell, H. A., Sen Gupta, A., Benthuisen, J. A., Feng, M., Oliver, E. C., Alexander, L. V., Burrows, M. T., Donat, M. G., Hobday, A. J., and Moore, P. J.: A global assessment of marine heatwaves and their drivers, *Nat. Commun.*, 10, 2624, <https://doi.org/10.1038/s41467-019-10206-z>, 2019.
- Holsman, K. K., Hazen, E. L., Haynie, A., Gourguet, S., Hollowed, A., Bograd, S. J., Samhuri, J. F., and Aydin, K.: Towards climate resiliency in fisheries management, *ICES J. Mar. Sci.*, 76, 1368–1378, <https://doi.org/10.1093/icesjms/fsz031>, 2019.
- Ibrahim, O., Mohamed, B., and Nagy, H.: Spatial variability and trends of marine heat waves in the eastern mediterranean sea over 39 years, *Journal of Marine Science and Engineering*, 9, 643, <https://doi.org/10.3390/jmse9060643>, 2021.
- Jacox, M. G., Tommasi, D., Alexander, M. A., Hervieux, G., and Stock, C. A.: Predicting the evolution of the 2014–2016 California Current System marine heatwave from an ensemble of coupled global climate forecasts, *Front. Mar. Sci.*, 6, 497, <https://doi.org/10.3389/fmars.2019.00497>, 2019.
- Jacox, M. G., Alexander, M. A., Amaya, D., Becker, E., Bograd, S. J., Brodie, S., Hazen, E. L., Pozo Buil, M., and Tommasi, D.: Global seasonal forecasts of marine heatwaves, *Nature*, 604, 486–490, <https://doi.org/10.1038/s41586-022-04573-9>, 2022.

- Juza, M., Fernández-Mora, À., and Tintoré, J.: Sub-Regional marine heat waves in the Mediterranean Sea from observations: Long-term surface changes, Sub-surface and coastal responses, *Front. Mar. Sci.*, 9, 785771, <https://doi.org/10.3389/fmars.2022.785771>, 2022.
- Lecci, R., Drudi, M., Grandi, A., Cretì, S., and Clementi, E.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea Physics Reanalysis, MED-SEA\_MULTIIYEAR\_PHY\_006\_004, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-004.pdf> (last access: 23 June 2023), 2022.
- Lecci, R., Drudi, M., Grandi, A., Cretì, S., and Clementi, E.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea Physics Analysis and Forecast, MED-SEA\_ANALYSISFORECAST\_PHY\_006\_013, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-013.pdf> (last access: 14 February 2024), 2023.
- Li, Z., Holbrook, N. J., Zhang, X., Oliver, E. C., and Coughon, E. A.: Remote forcing of Tasman Sea marine heatwaves, *J. Climate*, 33, 5337–5354, <https://doi.org/10.1175/JCLI-D-19-0641.1>, 2020.
- Liu, G., Eakin, C. M., Chen, M., Kumar, A., De La Cour, J. L., Heron, S. F., Geiger, E. F., Skirving, W. J., Tirak, K. V., and Strong, A. E.: Predicting heat stress to inform reef management: NOAA Coral Reef Watch's 4-month coral bleaching outlook, *Front. Mar. Sci.*, 5, 57, <https://doi.org/10.3389/fmars.2018.00057>, 2018.
- McAdam, R., Masina, S., and Gualdi, S.: Seasonal forecasting of subsurface marine heatwaves, *Commun. Earth Environ.*, 4, 225, <https://doi.org/10.1038/s43247-023-00892-5>, 2023.
- McAdam, R., Bonino, G., Clementi, E., and Masina, S.: Forecasting the Mediterranean Sea marine heatwave of summer 2022, Zenodo [data set], <https://doi.org/10.5281/zenodo.10944086>, 2024.
- McLeod, E., Salm, R., Green, A., and Almany, J.: Designing marine protected area networks to address the impacts of climate change, *Front. Ecol. Environ.*, 7, 362–370, <https://doi.org/10.1890/070211>, 2009.
- Notarbartolo-di-Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T., and Van Klaveren, P.: The Pelagos sanctuary for Mediterranean marine mammals, *Aquat. Conserv.*, 18, 367–391, <https://doi.org/10.1002/aqc.855>, 2008.
- Oliver, E. C., Benthuisen, J. A., Darmaraki, S., Donat, M. G., Hobday, A. J., Holbrook, N. J., Schlegel, R. W., and Sen Gupta, A.: Marine heatwaves, *Annu. Rev. Mar. Sci.*, 13, 313–342, <https://doi.org/10.1146/annurev-marine-032720-095144>, 2021.
- Pisano, A., Fanelli, C., Tronconi, C., Cesarini, C., La Padula, F., and Buongiorno Nardelli, B.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea – High Resolution L4 Sea Surface Temperature Reprocessed Product, SST\_MED\_SST\_L4\_REP\_OBSERVATIONS\_010\_021, Issue 3.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-021-022-041-042.pdf> (last access: 14 February 2024), 2023a.
- Pisano, A., Fanelli, C., Tronconi, C., Cesarini, C., La Padula, F., and Buongiorno Nardelli, B.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea – High Resolution L4 Sea Surface Temperature Reprocessed Product, SST\_MED\_SST\_L4\_REP\_OBSERVATIONS\_010\_021, Issue 3.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-021-022-041-042.pdf> (last access: 14 February 2024), 2023b.
- Poupkou, A., Zanis, P., Nastos, P., Papanastasiou, D., Melas, D., Tourpali, K., and Zerefos, C.: Present climate trend analysis of the Etesian winds in the Aegean Sea, *Theor. Appl. Climatol.*, 106, 459–472, 2011.
- Prioli, G.: Shellfish farming: technologies and production, *Vet. Res. Commun.*, 28, 51–56, <https://doi.org/10.1023/b:verc.0000045378.40910.7e>, 2004.
- Rayner, R., Jolly, C., and Gouldman, C.: Ocean observing and the blue economy, *Front. Mar. Sci.*, 6, 330, <https://doi.org/10.3389/fmars.2019.00330>, 2019.
- Rodrigues, R. R., Taschetto, A. S., Sen Gupta, A., and Foltz, G. R.: Common cause for severe droughts in South America and marine heatwaves in the South Atlantic, *Nat. Geosci.*, 12, 620–626, <https://doi.org/10.1038/s41561-019-0393-8>, 2019.
- Schlegel, R. W., Oliver, E. C., and Chen, K.: Drivers of marine heatwaves in the Northwest Atlantic: The role of air–sea interaction during onset and decline, *Front. Mar. Sci.*, 8, 627970, <https://doi.org/10.3389/fmars.2021.627970>, 2021.
- Sen Gupta, A., Thomsen, M., Benthuisen, J. A., Hobday, A. J., Oliver, E., Alexander, L. V., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., and Perkins-Kirkpatrick, S.: Drivers and impacts of the most extreme marine heatwave events, *Sci. Rep.*, 10, 19359, <https://doi.org/10.1038/s41598-020-75445-3>, 2020.
- Simon, A., Plecha, S. M., Russo, A., Teles-Machado, A., Donat, M. G., Auger, P. A., and Trigo, R. M.: Hot and cold marine extreme events in the Mediterranean over the period 1982–2021, *Front. Mar. Sci.*, 9, 892201, <https://doi.org/10.3389/fmars.2022.892201>, 2022.
- Smith, K. E., Burrows, M. T., Hobday, A. J., Sen Gupta, A., Moore, P. J., Thomsen, M., Wernberg, T., and Smale, D. A.: Socioeconomic impacts of marine heatwaves: Global issues and opportunities, *Science*, 374, eabj3593, <https://doi.org/10.1126/science.abj3593>, 2021.
- Smith, K. E., Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., Thomsen, M. S., Wernberg, T., and Smale, D. A.: Biological impacts of marine heatwaves, *Annu. Rev. Mar. Sci.*, 15, 119–145, <https://doi.org/10.1146/annurev-marine-032122-121437>, 2023.
- Vogt, L., Burger, F. A., Griffies, S. M., and Frölicher, T. L.: Local drivers of marine heatwaves: a global analysis with an earth system model, *Front. Clim.*, 4, 847995, <https://doi.org/10.3389/fclim.2022.847995>, 2022.
- White, C. J., Carlsen, H., Robertson, A. W., Klein, R. J. T., Lazo, J. K., Kumar, A., Vitart, F., Coughlan de Perez, E., Ray, A. J., Murray, V., Bharwani, S., MacLeod, D., James, R., Fleming, L., Morse, A. P., Eggen, B., Graham, R., Kjellström, E., Becker, E., Pegion, K. V., Holbrook, N. J., McEvoy, D., Depledge, M., Perkins-Kirkpatrick, S., Brown, T. J., Street, R., Jones, L., Remenyi, T. A., Hodgson-Johnston, I., Buontempo, C., Lamb, R., Meinke, H., Arheimer, B., and Zebiak, S. E.: Potential applications of subseasonal-to-seasonal (S2S) predictions, *Met. Apps*, 24, 315–325, <https://doi.org/10.1002/met.1654>, 2017.