



Mediterranean marine heatwave 2023: ecosystem and fisheries impacts in Italian waters

Riccardo Martellucci¹, Francesco Tiralongo², Sofia F. Darmaraki³, Michela D'Alessandro¹,
Giorgio Mancinelli⁴, Emanuele Mancini^{4,5}, Roberto Simonini⁶, Milena Menna¹, Annunziata Pirro¹,
Diego Borme¹, Rocco Auriemma¹, Marco Graziano¹, and Elena Mauri¹

¹Oceanography Section, National Institute of Oceanography and Applied Geophysics, Trieste, Italy

²Department of Biological, Geological and Environmental Sciences, University of Catania, Catania, Italy

³Foundation for Research and Technology Hellas, Heraklion, Greece

⁴Department of Biological and Environmental Sciences and Technologies, University of Salento, Lecce, Italy

⁵National Biodiversity Future Center (NBFC), Palermo, Italy

⁶Department of Life Sciences, University of Modena and Reggio Emilia, Modena, Italy

Correspondence: Riccardo Martellucci (rmartellucci@ogs.it) and Sofia F. Darmaraki
(sofia.darmaraki@dal.ca)

Received: 30 August 2024 – Discussion started: 20 September 2024

Revised: 7 March 2025 – Accepted: 13 May 2025 – Published: 30 September 2025

Abstract. In 2023, the Mediterranean Sea experienced the longest recorded marine heatwave (MHW) in four decades, affecting marine biodiversity, fisheries and coastal livelihoods. In this study, we assess the effects of this extreme event on the spread of the invasive species *Callinectes sapidus* (Atlantic blue crab) and *Hermodice carunculata* (bearded fireworm) along the Italian coasts. We focus on the coastal area of the Po River Delta in the northern Adriatic and on the northern and southern coasts of Sicily and investigate to what extent the increased seawater temperatures contribute to the increase in the monthly biomass of these species. Considering that the spread of the Atlantic blue crab is responsible for significant economic losses in the shellfish fishery and that the spread of the bearded fireworm poses a health risk to the artisanal fishing industry, we also assess the socio-economic impact of this MHW by analysing fish market data and online surveys. Finally, we discuss possible strategies to mitigate the spread and ecological impact of these invasive species. We take into account the aggressive feeding behaviour of the fireworm and the thermophilic nature of the fireworm, whose toxic antennae also pose a health risk to humans. Overall, the sustainability of marine ecosystems and coastal communities in the Mediterranean requires robust interdisciplinary collaboration to address the challenges posed by biological invasions and climate change in this region.

1 Introduction

The Mediterranean Sea, one of the most biodiverse marine ecosystems in the world (Coll et al., 2010), is currently facing unprecedented challenges due to extreme temperature events caused by climate change, known as marine heatwaves (MHWs) (Darmaraki et al., 2019). In recent decades, the frequency, intensity and duration of these record-breaking episodes have increased in the region, mainly due to the mean sea surface temperature (SST) warming trend in the Mediterranean Sea, which ranges between 0.035–0.041 °C yr⁻¹ (EU

Copernicus Marine Service Information, 2022a), almost twice as high as the corresponding global SST trend of 0.015 ± 0.001 °C yr⁻¹ (EU Copernicus Marine Service Information, 2022b). This has significant implications for the region's biodiversity and economy, as the warming trend and MHWs may facilitate the proliferation of invasive species (Joyce et al., 2024).

Species that pose a significant threat include the Atlantic blue crab, *Callinectes sapidus*, and the bearded fireworm, *Hermodice carunculata*, which have attracted attention due

Table 1. Products used in the present work. Complete references for the articles in Prod. 1, Prod. 2, Prod. 3 and Prod. 6 are reported in the References.

Ref. no.	Product name & type	Documentation
Copernicus products		
1	Copernicus Marine SST_MED_SST_L4_REP_OBSERVATIONS_010_021 Mediterranean Sea – High Resolution L4 Sea Surface Temperature Reprocessed	Quality Information Document (QUID): Pisano et al. (2016, 2024a); Product User Manual (PUM): Pisano et al. (2024b)
2	Copernicus Marine MEDSEA_MULTIYEAR_PHY_006_004 Mediterranean Sea Physics reanalysis	Quality Information Document (QUID): Escudier al. (2022); Product User Manual (PUM): Lecci et al. (2022)
Non-Copernicus products		
3	Crab and clam fishery data	CONSORZIO COOPERATIVE PESCATORI DEL POLESINE Organizzazione di Produttori Soc. Coop. A r.l., Via della Sacca, 11 45018 Scardovari (RO) – ITALIA. P.IVA 00224140293
4	Worms Out questionnaire	Link: bit.ly/3L3TWUc https://www.facebook.com/MonitoraggioVermocane (last access: 7 February 2025)
5	Righi et al. (2020) questionnaire	https://doi.org/10.12681/mms.23117
6	iNaturalist	https://www.inaturalist.org/ (last access: 7 February 2025)

to their rapid spread and negative impact on Italian fisheries (e.g. Heilskov et al., 2006; Riera et al., 2014; Simonini et al., 2019; Righi et al., 2020; Bardelli et al., 2023; Tiralongo et al., 2023). In particular, *C. sapidus*, which is native to the Atlantic, has rapidly colonised the Italian coasts (Mancinelli et al., 2021). *C. sapidus* is characterised by its voracious predatory behaviour and opportunistic feeding habits (Mancinelli et al., 2021) and has led to considerable economic losses in shellfish fisheries and enormous challenges for native species (Clavero et al., 2022). Similarly, *H. carunculata*, a thermophilic polychaete, has spread in Italian waters, affecting artisanal fisheries by both ruining the catch and posing health risks to human health (Heilskov et al., 2006; Riera et al., 2014; Simonini et al., 2019; Righi et al., 2020; Tiralongo et al., 2023). In fact, *H. carunculata* with its toxic setae can cause painful stings in humans, leading to burns and redness upon physical contact and posing a health risk for tourists in coastal areas and for the fishing community, especially when cleaning nets (Tiralongo et al., 2023). In addition, *H. carunculata* is an ecological disruptor as well as a direct threat to the well-being of coastal communities. The resilience of both species to environmental stressors and the rapid spread of the population emphasise the urgency of addressing the intensifying risks of climate change and bioinvasers with comprehensive management strategies.

The year 2023 marked a turning point when average global air temperatures reached an unprecedented high (Copernicus, 2024). The European continent experienced its second warmest year on record, with the Mediterranean basin experiencing a series of extreme temperature events (Marullo et al., 2023). Of particular concern was the occurrence of the longest recorded and one of the strongest surface MHWs of the last four decades, which persisted in the north-western Mediterranean from May 2022 to boreal spring 2023 (Marullo et al., 2023; Pirro et al., 2024). At its peak in July 2022, this MHW covered almost the entire western Mediterranean basin, with maximum daily SST anomalies reaching about 2.6 and 4.3 °C and anomalously warm conditions comparable to the 2003 summer MHW (Guinaldo et al., 2023). The long duration of the event was attributed to a combination of anomalously low wind speeds, high solar radiation and weak vertical mixing in the ocean (Marullo et al., 2023). These warming-related events had far-reaching impacts on marine life and coastal communities (He and Silliman, 2019).

The aim of this study is to investigate the spread and increase in the abundance of *C. sapidus* and *H. carunculata* associated with moderate and extreme warming in two coastal areas of Italy, particularly during the MHWs of 2022/2023. Relevant socio-economic impacts were also assessed by analysing fish market data and responses to questionnaires

handed out to the local fishing community and completed online, which addressed issues related to the bioinvasion of these species. In addition, we discuss possible solutions to mitigate the invasion of these species.

2 Methods

2.1 Study areas

The study was conducted in two different regions within Italian waters: two adjacent lagoons in the northern Adriatic Sea (Canarin and Scardovari) and two coastal areas of Sicily (Fig. 1).

2.1.1 The Po River Delta

The two lagoons studied in the northern Adriatic are transitional and shallow-water environments located in the Po River Delta and connected to the sea and various river branches (Fig. 1b and c). As they are directly influenced by the outflows of the Po River, these regions have highly dynamic hydro-morphological characteristics and are subject to rapid changes due to biotic and abiotic forces (Maicu et al., 2018; Franzoi et al., 2023). Although the lagoons are subject to various forms of anthropogenic pressure that have progressively altered their natural ecological characteristics (Franzoi et al., 2023), the lagoons support several mussel and oyster farms, which represent the main productive activities and vital economic resources at both local and regional level (Turolla, 2008; Bordignon et al., 2020; Chiesa et al., 2025). The Scardovari lagoon covers an area of 32 km², has an average depth of 1.5 m (Mistri et al., 2018), and is connected to the sea via two bays in the north-east and south-west of the basin. The Canarin lagoon covers an area of about 6.4 km² in the southern part of the study area and has an average depth of 0.9 m. It is connected to the Adriatic Sea by a shallow, approximately 200 m wide estuary with a maximum depth of 2.5 m.

2.1.2 Sicily

The island of Sicily (Fig. 1d) lies at the convergence of the eastern and western basins of the Mediterranean, which are influenced by both the relatively cooler Atlantic waters and the warmer Levantine waters. In particular, the eastern coast of Sicily (Ionian Sea) is significantly affected by quasidecadal reversals of the Northern Ionian Gyre driven by the mechanisms of the bimodal oscillation system in the Ionian Sea (Gačić et al., 2021; Menna et al., 2022). The distribution of water masses is altered by the bringing-in of warm and salty Levantine water during the cyclonic phase (anti-clockwise) and transportation of cooler Atlantic water during the anticyclonic phase (clockwise). This dynamic influences marine ecosystems and favours the occurrence of Levantine species during the cyclonic phase and vice versa

(Civitarese et al., 2023). In comparison, the northern coast of Sicily (Tyrrhenian Sea) experiences less saline and relatively cooler Atlantic water flowing in through the Sardinian Channel (Vetrano et al., 2010), while the southern coast of Sicily is characterised by cold water due to semi-permanent upwelling (Raffa et al., 2017).

2.2 Biological characteristics of the two species

2.2.1 *Callinectes sapidus*

The Atlantic blue crab *C. sapidus* Rathbun, 1896 is a species native to the western coasts of the Atlantic Ocean and is naturally distributed from Nova Scotia to northern Argentina (Millikin and Williams, 1984). It was first recorded in Europe in 1901 and in the Mediterranean in 1947 (Mancinelli et al., 2021). The first record in Italian waters dates back to 1949 from the lagoon of Venice, and ballast water is considered the most likely cause (Nehring, 2011). In Mediterranean waters, *C. sapidus* is considered one of the 100 most invasive alien species (Zenetos et al., 2005; Katsanevakis et al., 2018; Tsirintanis et al., 2022) and is present in at least seven of the nine southern European marine ecoregions (Mancinelli et al., 2017a, b). In the last 10 years, it has rapidly expanded its range to new ecosystems throughout the Mediterranean, such as the European Atlantic waters of Portugal, France, Belgium and Germany but also in Italian waters (Tiralongo et al., 2021; Bardelli et al., 2023). This eurythermal and euryhaline species is a voracious predator characterised by aggressive behaviour, high fecundity, excellent swimming abilities and high fertility (Tsirintanis et al., 2022), and it inhabits lagoons, estuaries and other coastal environments. *C. sapidus* has a complex, biphasic life cycle consisting of marine planktonic larvae (zoea) and benthic postlarvae (megalopa), with juveniles and adults living in estuaries, lagoons and other coastal habitats (Lipcius et al., 2007). In marine waters, this species lives mainly on soft substrates at depths between 1 and 90 m. Their life cycle is very complex and includes different habitats depending on sex and ontogenetic stage: adults can reach a relatively large size, with a carapace up to 25 cm wide in males and 18 cm in females (Millikin and Williams, 1984), and reside in lagoons and estuaries where males settle and moult. After mating, the egg-laying females migrate to the sea where they lay their eggs. The young return to the transitional areas and, after rapid growth, reach sexual maturity in their second year of life (Millikin and Williams, 1984; Taylor and Fehon, 2021). *C. sapidus* exhibits opportunistic feeding behaviour and feeds mainly on fish and invertebrates, especially bivalves and polychaetes, and may consume detritus and macrophytes when other food sources are scarce (Mancinelli et al., 2017a; Tiralongo et al., 2024b). Recent studies in the Po River Delta have shown that *C. sapidus* significantly affects the aquaculture of *Ruditapes philippinarum* (Manila clam). Predation by *C. sapidus* resulted in mussel losses of up to 100 % in certain areas, with up to 56 % of




Figure 1. Map of Italy (a) with the two study areas highlighted in red boxes. The map shows the main water basins in the region, with the blue line indicating 10 m rivers and lakes (retrieved from <https://www.naturalearthdata.com>, last access: 7 February 2025). The Po River Delta (b) and a detailed enlargement of the Scardovari (Sc) and Canarin (Ca) are highlighted in panel (c) (maps were downloaded from the EMODnet Digital Bathymetry available at <https://portal.emodnet-bathymetry.eu/>, last access: 7 February 2025). The examined study (coastal) areas around Sicily (d). For images of *C. sapidus* and *H. carunculata*, refer to Figs. S4 and S5.

mussel shells showing signs of predation and a complete absence of seeds in natural recruitment zones (Azzurro et al., 2024; Chiesa et al., 2025). Although several studies have been conducted on this species, the impacts and interactions of *C. sapidus* on native species and Mediterranean aquatic ecosystems are still poorly understood and require further investigation (Mancinelli et al., 2017b; Clavero et al., 2022).

2.2.2 *Hermodice carunculata*

The thermophilic amphinomid *H. carunculata* (Pallas, 1766), commonly known as the bearded fireworm, is a large predator/scavenger polychaeta found in warm and temperate areas of the Caribbean, Atlantic, Red Sea (Fishelson, 1971; Ahrens et al., 2013; Ramos and Schizas, 2023) and Mediterranean Sea (Simonini et al., 2018; Toso et al., 2022, 2024). Although it is native to the Mediterranean region, it is also considered a highly invasive species due to its increasing spread. Previous studies indicate that its abundance across the Mediterranean has increased in recent years, likely due to warmer temperatures favouring its northward spread (Righi et al., 2020; Toso et al., 2022). This may have a detrimental effect on the region's ecosystems and associated species, as well as human health, as it is resilient to natural and anthropogenic stressors (Schulze et al., 2017) and may also become a carrier of new pathogens (Sussman et al., 2003; Schulze et al., 2017). Similarly, coastal and anthropogenic activities such as fishing (Fig. S5 in the Supplement) and

bathing can also be affected (Celona and Comparetto, 2010; Cosentino and Giacobbe, 2011; Schulze et al., 2017; Simonini et al., 2018; Righi et al., 2020; Toso et al., 2020; Tiralongo et al., 2023). *H. carunculata* can grow to over 70 cm in length and reach a lifespan of 9 years (Simonini and Ferri, 2022). Its metamers are equipped with dorsal calcareous chaetae filled with a venom that is very effective against predators (Kicklighter and Hay, 2006; Schulze et al., 2017; Simonini et al., 2018, 2021; Righi et al., 2021, 2022). The presence of these defence mechanisms makes the polychaete highly resistant to predators, as none of the species identified in the Mediterranean region are able to effectively prey on it (Ladd and Shantz, 2016; Righi et al., 2021; Simonini et al., 2021). In contrast, *H. carunculata* is a voracious predator of sessile and benthic invertebrates (Wolf and Nugues, 2013; Wolf et al., 2014; Jumars et al., 2015; Barroso et al., 2016; Schulze et al., 2017; Simonini et al., 2018; Righi et al., 2020), and its ability to regenerate favours its dispersal (Toso et al., 2024). In addition, this species is characterised by a remarkable dispersal ability, which is attributed to the production of planktotrophic and particularly long-lived larvae (Ahrens et al., 2013; Schulze et al., 2017; Toso et al., 2020). In Italian waters, *H. carunculata* is widespread on rocky substrates between 1 and 20 m (Righi et al., 2020; Simonini et al., 2021), but in some areas of the Mediterranean it reaches greater depths and has also been observed in association with coralligenous and precoralligenous bioformations (Fishelson, 1971; Righi et al., 2020).

2.3 Temperature datasets

To identify surface MHWs on the study areas we obtained daily SST data from the Mediterranean Sea SST Analysis L4 product of the Copernicus Marine Service, covering the period 1982–2023 (Table 1, product ref. 1). This dataset provides gap-free, optimally interpolated, satellite-based estimates of SST with a resolution of $0.05^\circ \times 0.05^\circ$. For the analysis of subsurface temperatures in the areas of interest, daily vertical temperature profiles were obtained from the Mediterranean Sea Physics Reanalysis dataset for the period from 1993 to 2023 (Table 1, product ref. 2) with a spatial resolution of $0.042^\circ \times 0.042^\circ$. MHWs are detected whenever the SST exceeds a daily, 40-year (1982–2023) climatological threshold for at least 5 consecutive days, based on the identification framework proposed by Hobday et al. (2016).

2.4 The crab and clam fishery data

To assess the impact of the spread of *C. sapidus* on the local fishing industry, we use data on the production of mussels (*Ruditapes philippinarum*) provided by the Scardovari and Canarin Cooperative, which has been farming this species in the Po River Delta for years. The dataset contains monthly values for waste and sales of *C. sapidus*, representing the sum of daily harvests by fishers before reaching the market for fish sales (Table 1, ref. 3). Recent studies in the area have shown that *C. sapidus* eats mussels, as evidenced by claw marks on the mussels. Compared to previous years, mussel production in 2023 has decreased by 75 % in the Scardovari lagoon and 100 % in the Canarin lagoon (Azzurro et al., 2024; Chiesa et al., 2025), which is why we use discard data as an indicator of *C. sapidus* biomass and damage to fisheries.

2.5 Questionnaire for *Hermodice carunculata*

In recent decades, citizen participation in the collection of data useful for science has increased thanks to numerous awareness-raising initiatives (Turrini et al., 2018) and has already been recognised as a valuable resource for research, biodiversity monitoring and conservation (Lopez et al., 2019; Toivonen et al., 2019). Although in some cases this information lacks a solid scientific basis and needs to be validated by experts in the field, it offers the advantage that it can be collected over wide geographical areas at low cost (Ballard et al., 2017; Tirelli et al., 2021; Sun et al., 2021). For this reason, citizen science projects are currently on the rise in various areas, particularly as a tool for solving environmental and conservation issues (Kullenberg and Kasperowski, 2016; Turrini et al., 2018). For example, citizen participation is widely used in projects and initiatives for the sighting of non-indigenous, invasive and rare species such as the *Alien-Fish* project (<https://www.facebook.com/alienfish>, last access: 10 February 2025), *avvistAPP* (<https://www.avvistapp.it/>,

last access: 10 February 2025), *Monitoraggio Vermocane* (<https://www.facebook.com/MonitoraggioVermocane>) and *iNaturalist* (<https://www.inaturalist.org/>). To assess the impact of *H. carunculata* on human activities such as fishing and tourism, a questionnaire was developed and distributed to the fishing community as an online survey. The questionnaire, which builds on the observations of Righi et al. (2020), was distributed to a large number of people in 2023, who provided a total of 151 responses. It was primarily distributed via websites such as “Monitoraggio del Vermocane” and “Fauna Marina Mediterranea” (<https://www.facebook.com/groups/230601830399549>, last access: 10 February 2025) as well as via social media pages of various authors and also Italian diving centres. The questionnaire included 19 questions, of which 4 single-choice questions focusing on the frequency of sightings, the abundance of specimens and the perception of the species as a potential problem were analysed in this study. Most respondents reported having observed the fireworm while diving (74 %), spearfishing (10 %) and snorkelling (15 %), while no questions were asked about respondents’ age or employment status. Due to its distinctive morphology and bright colouration, *H. carunculata* is unlikely to be confused with other species, as there are no comparable organisms in the Mediterranean. This study was carried out as part of the “Worms Out” project funded by the National Institute of Oceanography and Applied Geophysics and the ECCSEL NatLab Italy project (Table 1, ref. 5). In addition to the real-time data collected by marine users, the use of this questionnaire is important to identify and assess the occurrence of this species. This is to address the current lack of scientific reports on this species, which is probably due to the limited number of sampling efforts (e.g. Frascchetti et al., 2002; Giangrande et al., 2003; Corriero et al., 2004; Mastrototaro et al., 2010). To address the scarcity of scientific reports, we integrated observations from various sources such as the *iNaturalist* observations, the results of Righi et al. (2020) and the most recent observations from the 2023 online survey. Integrating this information with the scientific literature and observations is crucial for monitoring biological invasions and studying native invasive species (Azzurro et al., 2019; Giovos et al., 2019; Toivonen et al., 2019).

3 Results

3.1 Northern Adriatic

In the first half of 2023, mussel production in the area was around 400 t per month and peaked in August (800 t) before drastically declining from September to December (Fig. 2a). In contrast, the sale of crabs began in the summer months, with the highest discard observed in August 2023 with a total of 300 t. Towards the end of 2023, crab discards were comparable to sales (Fig. 2b). Throughout 2023, the study area in the northern Adriatic was also characterised by particularly high SSTs with six MHWs observed (Fig. 2c and d). The

first event occurred in March, lasted 5 d and had a moderate intensity, while the remaining events occurred during the summer and autumn seasons. The most intense MHW was observed at the end of August 2023 and was characterised by temperatures that were more than 3 °C above normal and lasted 11 d (Table S1 in the Supplement). The longest event was observed in October and lasted 36 d with a strong intensity (> 2.6 °C). The high temperatures of 2023 also affected the entire water column, with the strongest subsurface temperature anomalies (> 4 °C) observed between 4 m and 15 m depth during the summer season (Fig. 2d). This indicates a significant increase in subsurface temperatures in this area, as typical deviations from the 1993–2016 mean are between 0.8 and 1.2 °C (Fig. S1). Regarding the potential cumulative effects of temperature on the life cycle of *C. sapidus*, although the SST in 2023 was not below the threshold for hibernation (5 °C), it exceeded the temperatures for reproductive activity (10.8 °C) and larval development (19 °C) by 300 and 170 d, respectively (dashed lines in Fig. 2c).

3.2 Sicily

In the two coastal regions of northern and eastern Sicily, the SST remained above the climatological values throughout 2023 (Fig. 3b, c): eastern Sicily experienced three MHWs that lasted about 60 d in total (Table S1). The most intense event occurred in July and lasted 21 d, and the longest was in October with a duration of 30 d. On the north coast of Sicily, there were slightly longer MHWs on average: the first event occurred in March, lasted 5 d and had an intensity of 1.4 °C. The most intense MHW (> 2.5 °C) was observed between July and August with a duration of 25 d, while the longest event (49 d) occurred in autumn with an intensity of 1.6 °C. Compared to the period 1993–2016, temperatures in the entire water column in both regions were around 1.2–4 °C warmer than normal and even warmer on the north coast of Sicily. The upper 80 m of the water column shows the highest temperature anomalies (> 2 °C) throughout the year, especially in the autumn months (> 2.5 °C) in both areas. However, during some days in summer, the subsurface layers between 10 and 50 m depth show negative temperature anomalies (down to –2 °C) (Fig. 3d, e), with temperatures dropping below 14° at greater depths. Typically, subsurface temperatures deviate by about 0.8–2 °C from the 1993–2016 mean, with the highest values observed in the upper 20–80 m depth between June–November (Fig. S1). The progressive temperature increase of around 0.03 °C yr^{–1} observed on the north coast of Sicily seems to be consistent with an increasing trend in the records of *Hermodice carunculata* over the last 20 years, especially in 2007–2008, 2014–2015 and 2023, when the highest number of observations was recorded (Fig. 3a). This increasing trend is also supported by the results of our proposed questionnaire, which shows a significant increase in the frequency of sightings in recent years compared to 2018, particularly during recreational ac-

tivities. In particular, the results of the questionnaire show that the presence of this species is increasingly recognised as a problem, particularly as *H. carunculata* is observed in areas where it was not previously seen (Fig. 3h and j). The data from the *Hermodice* questionnaire were compared with those from the iNaturalist platform (Table 1, ref. 6), and the two datasets showed similar observations, particularly in relation to the observed trend (Fig. 2a).

4 Discussion

The two Mediterranean regions analysed in the present study experienced multiple, prolonged and strong surface MHWs throughout 2023, with temperature anomalies ranging between 1.6–2.6 °C on the eastern and northern coasts of Sicily and between 2–3 °C on the northern Adriatic coast during the events. These events were associated with a general warming trend that was evident in increased monthly temperature anomalies occurring throughout the year, especially during summer in the northern Adriatic. Additionally, a trend of 0.03 °C per year was particularly evident in the sea surface temperature along the coast of northern Sicily during the period 2000–2023. Compared to the study areas in Sicily, the northern Adriatic showed a slightly stronger warming of the entire water column, which is probably due to its shallower depth. Also, lower-than-normal subsurface temperatures were observed in the upper 40 m of the two study areas in Sicily in summer and autumn. This is likely related to a stronger stratification of the upper ocean during these periods and a shift in the thermocline (see Figs. S2, S3), resulting in cooler temperatures near the surface, similar to what Pirro et al. (2024) observed. Overall, a significant warming was observed in all study areas in 2023, which in turn may have led to an increase in the abundance of the two invasive species in both areas.

In the case of *Callinectes sapidus*, the higher temperatures may have triggered a positive feedback loop in the phenology of the larval and adult stages, ultimately leading to higher survival and reproduction rates and driving the population increase (Costlow, 1967; Gencer, 2024). For marine invertebrates with complex life cycles, the effects of climate warming are particularly pronounced during critical stages (Libralato et al., 2015; Alter et al., 2024) such as larval development, reproductive activity or winter dormancy (dashed line in Fig. 2c). A similar mechanism has been proposed for other invasive crayfish such as *Hemigrapsus takanoi* as well as for other native brachyurans (Valdes et al., 1991; Anger, 2001; van den Brink et al., 2012; Oh and Lee, 2020). In this context, rising temperatures may have increased the invasiveness of the species in regions where established populations remained at low levels, as has been observed in the northern Adriatic Sea, where *C. sapidus* has been detected episodically since 1949 (Manfrin et al., 2016). Considering that in this species winter dormancy, adult reproduction and




Figure 2. Northern Adriatic study area: **(a)** time series of daily, spatially averaged SST anomalies of 2023 relative to the period 1993–2016 (red line) and monthly evolution of sold clams (black cross). **(b)** Monthly evolution of sold (green triangles) and discharged (purple diamonds) *C. sapidus* during 2023. **(c)** Time series of daily, spatially averaged SST during 2023 (black), smoothed SST climatology (blue) and 90th percentile threshold of SST (green) based on the 1982–2023 period. MHWs are indicated in red and identified using the Hobday et al. (2016) definition. The three dashed lines represent the temperature thresholds for winter dormancy (5 °C), reproductive activity (10.8 °C) and larval development (19 °C) of *C. sapidus*. **(d)** Vertical profile of temperature anomalies during 2023, relative to the period 1993–2016, spatially averaged at each depth. Temperature data were obtained from Copernicus Marine Service (Table 1, product ref. 1, 2). Clams and crab data were obtained from the Consorzio Cooperative Pescatori del Polesine (Table 1, product ref. 3).

early life stages (egg maturation, zoea and megalopa development) are strongly regulated by temperature minima (Brylawsky and Miller 2006; Rogers et al., 2022; Schneider et al., 2024), an increase in water temperatures may have additionally accelerated metamorphosis from zoea to megalopa, reducing predation risk and promoting survival and ultimately population abundance. On the other hand, a further increase in temperature above 26 °C may have an opposite effect by reducing the number of moults per larva and causing a reduction in blue crab growth (Gencer, 2024).

An increase in water temperatures also accelerates development and growth rates in *Hermodice carunculata* (Libralato et al., 2015; Alter et al., 2024), and several studies identify water temperature as a decisive factor in the spread of the species (Righi et al., 2020; Tiralongo et al., 2023; Toso et al., 2024) by influencing the species range shifts and facilitating its establishment and spread (Stachowicz et al., 2002; Samperio-Ramos et al., 2015). Our questionnaires and iNatu-

ralist observations confirm this spread and show a significant increase in sightings over the last 5 years. Our results are thus consistent with the documented increase in the presence of the species and its bathymetric extent over the last two decades, which coincides with a general increase in water temperatures along the coasts of Sicily (Pisano et al., 2020; Righi et al., 2020; Tiralongo et al., 2023; Kubin et al., 2023), which has been particularly evident since 2023 (Fig. 3a). Given the ongoing warming trend in the Mediterranean, it is likely that this thermophilic species will continue to spread along the north-western Mediterranean coast, where it was frequently observed in shallow-water conditions, especially in the summer months (Schulze et al., 2017; Encarnação et al., 2019; Righi et al., 2020). Noticeably, both the remarkable increase in the monthly biomass of *C. sapidus* in the coastal areas of the northern Adriatic and the annual records of *H. carunculata* on the coasts of Sicily share a number of common implications. In particular, we show that the prolif-




Figure 3. (a) Yearly-averaged time series and linear trend of SST (red), spatially averaged over the northern coast of Sicily, and yearly records of *H. carunculata* (black cross) with their linear trend, based on Righi et al. (2020) and our questionnaire for the period 2000–2023 (Table 1, product ref. 4 and 5); the green crosses represent the biannual mean of *H. carunculata* from iNaturalist. Daily and spatially averaged SST time series during 2023 (black), smoothed SST climatology (blue), and 90th percentile threshold of SST (green) based on the 1982–2023 period for the northern **(b)** and eastern **(c)** coasts of Sicily. MHWs are indicated in red and identified using the Hobday et al. (2016) definition. Vertical profile of spatially averaged temperature anomalies during 2023, relative to the climatological period of 1993–2016 for the northern **(d)** and eastern **(e)** coasts of Sicily. The climatological depth of the 14 °C isotherm is displayed by the dashed blue line, whereas the depth of the 14 °C isotherm during 2023 by the solid black line. **(f)–(j)** Main results of the Worms Out questionnaire (Table 1, product ref. 4). Temperature data were obtained from Copernicus Marine Service (Table 1, product ref. 1, 2).

eration of *C. sapidus* in the Po River Delta significantly disrupted mussel production in 2023 (Fig. S4) and that, under current conditions, no resurgence of mussel populations is expected in 2024 (Chiesa et al., 2025). The dramatic impact on the mussel fishery is exacerbated by the cost of removing the crabs discarded by the fishery (Fig. S4). This expansion is also expected to disrupt benthic habitats, altering community composition and leading to remarkable changes at the level of the entire food web, as observed in the Ebro River Delta (Clavero et al., 2022) and recently in the Po River Delta in the Goro lagoon (Gavioli et al., 2025).

Similarly, *H. carunculata* poses a threat to both marine biodiversity and the economic stability of local fisheries, as has been observed for other annelids (Berke et al., 2010; Pires et al., 2015). Although comprehensive information on the impact of the species has only recently been presented (Tiralongo et al., 2023), the fireworm is known to cause both direct and indirect damage to fisheries. Direct damage includes the severing of secondary lines attached to hooks, either by the worm's teeth or by hiding among the rocks after consuming the bait (Tiralongo et al., 2023). To minimise this impact on the target species, the soaking time of the fishing gear has been reduced, which leads to a decrease in catch rates (Simonini et al., 2021; Tiralongo et al., 2020). The scavenging activities of *H. carunculata* not only affect the efficiency of fisheries, but also impact the fishing industry by damaging fish catches and reducing the market value of fish. The economic impact is estimated at around EUR 7.32 kg⁻¹ of damaged fish, resulting in significant annual losses given the total weight of commercially valuable catches (Tiralongo et al., 2023). The continued temperature-induced spread of *H. carunculata*, as observed in our results and documented by Righi et al. (2020) and Tiralongo et al. (2023), emphasises the urgent need for effective mitigation strategies to address the impacts caused by climate change on fisheries, tourism and the coastal economy.

Implication for human life and solutions for stakeholder

The ongoing invasion of *C. sapidus* provides an opportunity to evaluate strategies and measures to contain this spread and mitigate its ecological impact, even in freshwater ecosystems (Scalici et al., 2022; Tiralongo et al., 2024b, a; Bedmar et al., 2024). As the economic value of the species is already internationally recognised, this species can be used for both food and other purposes. Several studies have shown that over-harvesting plays an important role in the control of invasive species (Mancinelli et al., 2017b; Giakoumi et al., 2019), as exemplified by the commercial harvest of *C. sapidus*, which supports a significant fishery on the coasts of the USA (Hines, 2007; Kennedy et al., 2007; Bunnell et al., 2010); accordingly, control measures should target similar commercialisation strategies. Despite its introduction to the Italian market, the consumption of *C. sapidus* is not yet widespread, complicating efforts to eradicate it (Azzurro et al., 2024). Ef-

fective management strategies should therefore include the cultural integration of this species. For example, the Italian government has promoted *C. sapidus* by showcasing it at the 2024 G7 summit and distributing online promotional content. Expanding outreach efforts through targeted events and educational campaigns could further boost consumption of this species, similar to what has been done successfully with other species. Furthermore, the extraction of chitosan and astaxanthin from crab shells could support the ongoing shellfish market while providing valuable compounds with multiple applications in pharmaceutical, biomedical, cosmetic, agricultural and biotechnological fields (Ambati et al., 2014; see also Demir et al., 2016, and Baron et al., 2017, for recent examples on *C. sapidus*).

Similarly, the increasing frequency of *H. carunculata* sightings emphasises the need for effective management strategies to contain its spread. As an efficient scavenger, predatory generalist and opportunistic consumer that can also feed on carrion, *H. carunculata* has also been found in large numbers under aquaculture net cages and in places with high anthropogenic pressure and organic enrichment, such as artisanal fishing harbours (Heilskov et al., 2006; Riera et al., 2014; Righi et al., 2020). Due to its ability to tolerate captivity, *H. carunculata* offers potential applications for the disposal of waste from the production and processing of marine products. Recent biorefinery research is investigating the use of *H. carunculata* in the processing of molluscan waste, particularly expired mussels from retail outlets, to recover and valorise the shells. Preliminary results show that *H. carunculata* consumes mussel meat at high rates and leaves the shells almost completely clean (Simonini et al., 2024). Given the ability to maintain them at high densities without substrate, the species could prove useful for the valorisation of waste shells, with clean shells serving as a source of “green” calcium carbonate (Seesanong et al., 2023). In addition, experiments could be conducted to determine the effectiveness of this species when used in IMTA (integrated multitrophic aquaculture) systems to determine its performance as a bioremediator organism (Giangrande et al., 2020). The development of practical applications for this invasive species could also support the elimination of areas where *H. carunculata* becomes a pest (Simonini et al., 2024).

Thus, the costs of managing and controlling invaded habitats may ultimately yield gains for the local population while greatly reducing the impact of the invader and even enhancing the ecosystem goods and services provided by coastal habitats. Collaborative efforts are essential to formulate adaptation measures that protect both marine ecosystems and the livelihoods of Mediterranean coastal communities. Through interdisciplinary collaboration and proactive management strategies, it is possible to mitigate the negative impacts of climate change and the spread of invasive species, thus ensuring the long-term sustainability of the Mediterranean marine environment and the well-being of coastal communities.

5 Conclusions

Overall, global warming and biological invasions in marine ecosystems are so far recognised to be closely linked, although the extent of their interactions and the role of climate change as a driving force remain controversial, as they vary along the invasion process and are influenced by species-specific responses to warming (Blackburn et al., 2011; Katsanevakis et al., 2014; Joyce et al., 2024). These responses affect the distribution, demography and life histories of invasive species. Currently, there is an urgent need for additional studies investigating the relationship between climate warming and the phenology of bioinvasive species to elucidate the indirect effects that may occur on both the distribution and abundance of the latter. The present study, focusing on *Callinectes sapidus* and *Hermodice carunculata*, is one of the first attempts to address this important issue in the Mediterranean. However, temperature alone may not be the only determinant of increases in invasive species abundance: for example, for *H. carunculata*, Simonini et al. (2021) suggest that fishing practices may facilitate the spread, establishment and survival of the species, with fish waste and other organic debris resulting from the cleaning of nets potentially providing increased trophic resource availability. In addition, releasing pregnant *C. sapidus* females after capture can significantly increase the number of larvae and increase populations (Hines et al., 2008). In the Po River Delta, this practise has become common among fishing companies given the low value of egg-laying females in local fish markets (Tiralongo, personal observation; see also Fig. S4). These examples illustrate how inadequate management practises can inadvertently favour the spread of invasive species and ultimately highlight the need for integrated control and mitigation strategies that take into account the diverse range of factors that can actually contribute to the success of a bioinvasion.

Code and data availability. The codes and data are available upon request to Annunziata Pirro (rmartellucci@ogs.it). Further information can also be found in Table 1.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/sp-6-osr9-9-2025-supplement>.

Author contributions. RM, FT, SFD, MD'A, GM and EmM contributed to the original draft writing, conceptualisation, investigation, data curation and formal analysis. RM and SFD were also responsible for visualisation, while FT and SFD coordinated project management. MD'A provided additional resources, and GM contributed supervision. RS, MM, AP, DB, RA and MG were involved in review and editing, supporting investigation, data maintenance, resourcing, and validation. EIM also oversaw project administration and acquired funding.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. Sofia F. Darmaraki acknowledge the financial support from the Hellenic Foundation for Research and Innovation (HFRI) under the third call of the "Research projects to support Post-Doctoral Researchers" scheme, project number 7077, acronym TexMed. This work was partly funded by the ECCSEL-NatLab Italy project, made by the extraordinary MIUR contribution (FOE) for the participation of Italy in the activities related to the international ECCSEL infrastructure (European Carbon Dioxide Capture and Storage Laboratory Infrastructure), with the aim of developing and maintaining the natural laboratories of Panarea (Aeolian Islands) and Latera (Lazio).

The publication has been funded by EU – Next Generation EU Mission 4, Component 2 – CUP B53C22002150006 – Project IR0000032 – ITINERIS – Italian Integrated Environmental Research Infrastructures System.

Additional financial support was provided to Emanuele Mancini by the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 – Call for tender No. 3138 of 16 December 2021, rectified by Decree no. 3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – Next Generation EU Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP D33C22000960007, Project title "National Biodiversity Future Center – NBFC".

Financial support. This research has been supported by the EU – Next Generation (grant nos. CUP B53C22002150006 and CUP D33C22000960007), the Hellenic Foundation for Research and Innovation (HFRI, project number 7077, TexMed), and the ECCSEL-NatLab Italy project.

Review statement. This paper was edited by Marta Marcos and reviewed by Rodrigo Riera and one anonymous referee.

References

- Ahrens, J. B., Borda, E., Barroso, R., Paiva, P. C., Campbell, A. M., Wolf, A., Nuges, M. M., Rouse, G. W., and Schulze, A.: The curious case of *Hermodice carunculata* (Annelida: Amphionomidae): evidence for genetic homogeneity throughout the Atlantic Ocean and adjacent basins, *Mol. Ecol.*, 22, 2280–2291, <https://doi.org/10.1111/mec.12263>, 2013.
- Alter, K., Jacquemont, J., Claudet, J., Lattuca, M. E., Barrantes, M. E., Royer, C., Cormier, M., Ferrari, R., Cominassi, L., Crochelet,

- E., Boudouresque, C. F., and Lejeune, C.: Hidden impacts of ocean warming and acidification on biological response of marine animals revealed through meta-analysis, *Nat. Commun.*, 15, 2885, <https://doi.org/10.1038/s41467-024-47064-3>, 2024.
- Ambati, R. R., Phang, S. M., Ravi, S., and Aswathanarayana, R. G.: Astaxanthin: sources, extraction, stability, biological activities and its commercial applications – A review, *Mar. Drugs*, 12, 128–152, <https://doi.org/10.3390/md12010128>, 2014.
- Anger, K.: The biology of decapod crustacean larvae, AA Balkema Publishers, Lisse, Vol. 14, 1–420, ISBN 9026518285, 2001.
- Azzurro, E., Sbragaglia, V., Cerri, J., Bariche, M., Bolognini, L., Ben Souissi, J., and Moschella, P.: Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge, *Glob. Change Biol.*, 25, 2779–2792, <https://doi.org/10.1111/gcb.14724>, 2019.
- Azzurro, E., Bonanomi, S., Chiappi, M., De Marco, R., Luna, G. M., Cella, M., Guicciardi, S., Tiralongo, F., Bonifazi, A., and Strafella, P.: Uncovering unmet demand and key insights for the invasive blue crab (*Callinectes sapidus*) market before and after the Italian outbreak: Implications for policy-makers and industry stakeholders, *Mar. Policy*, 167, 106295, <https://doi.org/10.1016/j.marpol.2024.106295>, 2024.
- Ballard, H. L., Dixon, C. G., and Harris, E. M.: Youth-focused citizen science: Examining the role of environmental science learning and agency for conservation, *Biol. Conserv.*, 208, 65–75, <https://doi.org/10.1016/j.biocon.2017.01.021>, 2017.
- Bardelli, R., Mancinelli, G., Mazzola, A., and Vizzini, S.: The Atlantic blue crab *Callinectes sapidus* spreading in the Tyrrhenian sea: Evidence of an established population in the Stagnone di Marsala (Sicily, southern Italy), *NAŠE MORE: znanstveni časopis za more i pomorstvo*, 70, 177–183, <https://doi.org/10.17818/NM/2023/SI6>, 2023.
- Baron, R. D., Pérez, L. L., Salcedo, J. M., Córdoba, L. P., and Sobral, P. J. d. A.: Production and characterization of films based on blends of chitosan from blue crab (*Callinectes sapidus*) waste and pectin from Orange (*Citrus sinensis* Osbeck) peel, *Int. J. Biol. Macromol.*, 98, 676–683, <https://doi.org/10.1016/j.ijbiomac.2017.01.061>, 2017.
- Barroso, R., Almeida, D., Contins, M., Filgueiras, D., and Dias, R.: *Hermodice carunculata* (Pallas, 1766) (Polychaeta: Amphinomidae) preying on starfishes, *Mar. Biodivers.*, 46, 333–334, <https://doi.org/10.1007/s12526-015-0394-9>, 2016.
- Bedmar, S., Oficialdegui, F. J., and Clavero, M.: Far-reaching blues: Long-distance migration of the invasive Atlantic blue crab, *Aquat. Conserv.*, 34, e4136, <https://doi.org/10.1002/aqc.4136>, 2024.
- Berke, S. K., Mahon, A. R., Lima, F. P., Halanych, K. M., Wetthey, D. S., and Woodin, S. A.: Range shifts and species diversity in marine ecosystem engineers: Patterns and predictions for European sedimentary habitats, *Global Ecol. Biogeogr.*, 19, 223–232, <https://doi.org/10.1111/j.1466-8238.2009.00509.x>, 2010.
- Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., and Richardson, D. M.: A proposed unified framework for biological invasions, *Trends Ecol. Evol.*, 26, 333–339, <https://doi.org/10.1016/j.tree.2011.03.023>, 2011.
- Bordignon, F., Zomeño, C., Xiccato, G., Birolo, M., Pascual, A., and Trocino, A.: Effect of emersion time on growth, mortality and quality of Pacific oysters (*Crassostrea gigas*, Thunberg 1973) reared in a suspended system in a lagoon in Northern Italy, *Aquaculture*, 528, 735481, <https://doi.org/10.1016/j.aquaculture.2020.735481>, 2020.
- Bunnell, D. B., Lipton, D. W., and Miller, T. J.: The bioeconomic impact of different management regulations on the Chesapeake Bay blue crab fishery, *N. Am. J. Fish. Manage.*, 30, 1505–1521, <https://doi.org/10.1577/M09-182.1>, 2010.
- Celona, A. and Comparetto, G.: Prime osservazioni sulla predazione opportunistica del “vermocene” *Hermodice carunculata* (Pallas, 1766), ai danni della piccola pesca artigianale nelle acque di Lampedusa (Is. Pelagie), *Annali della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 20, 15–20, <http://www.dlib.si/details/URN:NBN:SI:DOC-YNXLVSY> (last access: 7 February 2025), 2010.
- Chiesa, S., Petochi, T., Brusà, R. B., Raicevich, S., Cacciatore, F., Franceschini, G., Antonini, C., Vallini, C., Bernarello, V., Oselladore, F., Ciani, M., Di Blasio, L., Campolunghi, M. P., Baldessin, F., Boldrin, L., and Marino, G.: Impacts of the blue crab invasion on Manila clam aquaculture in Po Delta coastal lagoons (Northern Adriatic Sea, Italy), *Estuar. Coast. Shelf Sc.*, 312, 109037, <https://doi.org/10.1016/j.ecss.2024.109037>, 2025.
- Civitaresi, G., Gačić, M., Batistić, M., Bensi, M., Cardin, V., Dulčić, J., and Menna, M.: The BiOS mechanism: history, theory, implications, *Prog. Oceanogr.*, 216, 103056, <https://doi.org/10.1016/j.pocean.2023.103056>, 2023.
- Clavero, M., Franch, N., Bernardo, R., López, V., Abelló, P., and Mancinelli, G.: Severe, rapid, and widespread impacts of an Atlantic blue crab invasion, *Mar. Pollut. Bull.*, 176, 113479, <https://doi.org/10.1016/j.marpolbul.2022.113479>, 2022.
- Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Ben Rais Lasram, F., Aguzzi, J., and Voultsiadou, E.: The biodiversity of the Mediterranean Sea: estimates, patterns, and threats, *PLOS ONE*, 5, e11842, <https://doi.org/10.1371/journal.pone.0011842>, 2010.
- Copernicus: 2023 is the hottest year on record, with global temperatures close to the 1.5 °C limit, Copernicus, <https://climate.copernicus.eu/copernicus-2023-hottest-year-record> (last access: 7 February 2025), 2024.
- Corriero, G., Gherardi, M., Giangrande, A., Longo, C., Mercurio, M., and others: Inventory and distribution of hard bottom fauna from the marine protected area of Porto Cesareo (Ionian Sea): Porifera and Polychaeta, *Ital. J. Zool.*, 71, 237–245, <https://doi.org/10.1080/11250000409356578>, 2004.
- Cosentino, A. and Giacobbe, S.: The new potential invader *Linopherus canariensis* (Polychaeta: Amphinomidae) in a Mediterranean coastal lake: Colonization dynamics and morphological remarks, *Mar. Pollut. Bull.*, 62, 236–245, <https://doi.org/10.1016/j.marpolbul.2010.10.013>, 2011.
- Costlow, J. D.: The effect of salinity and temperature on survival and metamorphosis of megalops of the blue crab *Callinectes sapidus*, *Helgoländ. Wiss. Meer.*, 15, 84–97, <https://doi.org/10.1007/BF01618611>, 1967.
- Darmaraki, S., Somot, S., Sevault, F., Nabat, P., Cabos Narvaez, W. D., Cavicchia, L., and Sein, D. V.: Future evolution of marine heatwaves in the Mediterranean Sea, *Clim. Dynam.*, 53, 1371–1392, <https://doi.org/10.1007/s00382-019-04661-z>, 2019.
- Demir, D., Öfkeli, F., Ceylan, S., and Bölgen, N.: Extraction and characterization of chitin and chitosan from blue crab and synthesis of chitosan cryogel scaffolds, *Journal of the*

- Turkish Chemical Society Section A: Chemistry, 3, 131–144, <https://doi.org/10.28978/nesciences.970546>, 2016.
- Encarnação, J., Morais, P., Baptista, V., Cruz, J., and Teodósio, M. A.: New evidence of marine fauna tropicalization off the Southwestern Iberian Peninsula, *Diversity*, 11, 48, <https://doi.org/10.3390/d11040048>, 2019.
- Escudier, R., Clementi, E., Nigam, T., Aydogdu, A., Fini, E., Pistoia, J., Grandi, A., and Miraglio, P.: EU Copernicus Marine Service Quality Information Document for Mediterranean Sea Physics Reanalysis, MED SEA_MULTIYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-004.pdf> (last access: 10 February 2025), 2022.
- EU Copernicus Marine Service Information (CMEMS): Mediterranean Sea Surface Temperature time series and trend from Observations Reprocessing, Mercator Ocean International, Marine Data Store (MDS) [data set], <https://doi.org/10.48670/moi-00268>, 2022a.
- EU Copernicus Marine Service Information (CMEMS): Global Ocean Sea Surface Temperature time series and trend from Observations Reprocessing, Marine Data Store (MDS) [data set], <https://doi.org/10.48670/moi-00242>, 2022b.
- Fishelson, L.: Ecology and distribution of the benthic fauna in the shallow waters of the Red Sea, *Mar. Biol.*, 10, 113–133, <https://doi.org/10.1007/BF00354828>, 1971.
- Franzoi, P., Facca, C., Redolfi Bristol, S., Boschiero, M., Matteo, Z., and Scapin, L.: Application of the Habitat Fish Biological Index (HFBI) for the assessment of the ecological status of Po Delta lagoons (Italy), *Italian Journal of Freshwater Ichthyology*, 9, 91–106, <https://iris.unive.it/handle/10278/5046422> (last access: 7 February 2025), 2023.
- Fraschetti, S., Giangrande, A., Terlizzi, A., Miglietta, M. P., Della Tommasa, L., and Boero, F.: Spatio-temporal variation of hydroids and polychaetes associated with *Cystoseira amentacea* (Fucales: Phaeophyceae), *Mar. Biol.*, 140, 949–958, <https://doi.org/10.1007/s00227-001-0770-9>, 2002.
- Gačić, M., Ursella, L., Kovačević, V., Menna, M., Malačič, V., Bensi, M., Negretti, M.-E., Cardin, V., Orlić, M., Sommeria, J., Viana Barreto, R., Viboud, S., Valran, T., Petelin, B., Siena, G., and Rubino, A.: Impact of dense-water flow over a sloping bottom on open-sea circulation: laboratory experiments and an Ionian Sea (Mediterranean) example, *Ocean Sci.*, 17, 975–996, <https://doi.org/10.5194/os-17-975-2021>, 2021.
- Gavioli, A., Mancinelli, G., Turolla, E., Lanzoni, M., Paesanti, V., Soana, E., and Castaldelli, G.: Impacts of the invasive blue crab *Callinectes sapidus* on small-scale fisheries in a Mediterranean lagoon using fishery landing data, *Sci. Total Environ.*, 974, 179236, <https://doi.org/10.1016/j.scitotenv.2025.179236>, 2025.
- Gencer, Ö.: The impact of an abiotic variable, temperature, on larvae of the blue crab, *Callinectes sapidus* Rathbun, 1896 (Brachyura, Portunidae), *Crustaceana*, 97, 137–150, <https://doi.org/10.1163/15685403-bja10239>, 2024.
- Giakoumi, S., Katsanevakis, S., Albano, P. G., Azzurro, E., Cardoso, A. C., Cebrian, E., and Sghaier, Y. R.: Management priorities for marine invasive species, *Sci. Total Environ.*, 688, 976–982, <https://doi.org/10.1016/j.scitotenv.2019.06.282>, 2019.
- Giangrande, A., Delos, A. L., Fraschetti, S., Musco, L., and Licciano, M.: Polychaete assemblages along a rocky shore on the South Adriatic coast (Mediterranean Sea): patterns of spatial distribution, *Mar. Biol.*, 143, 1109–1116, <https://doi.org/10.1007/s00227-003-1162-0>, 2003.
- Giangrande, A., Pierri, C., Arduini, D., Borghese, J., Licciano, M., Trani, R., Corriero, G., Basile, G., Cecere, E., Petrocelli, A., Stabili, L., and Longo, C.: An innovative IMTA system: Polychaetes, sponges and macroalgae co-cultured in a Southern Italian in-shore mariculture plant (Ionian Sea), *Journal of Marine Science and Engineering*, 8, 733, <https://doi.org/10.3390/jmse8100733>, 2020.
- Giovos, I., Kleitou, P., Poursanidis, D., Batjakas, I., Bernardi, G., Crocetta, F., Doumpas, N., Kalogirou, S., Kampouris, T. E., Keramidas, I., Langeneck, J., and Azzurro, E.: Citizen-science for monitoring marine invasions and stimulating public engagement: a case project from the eastern Mediterranean, *Biol. Invasions*, 21, 3707–3721, <https://doi.org/10.1007/s10530-019-02083-w>, 2019.
- Guinaldo, T., Voldoire, A., Waldman, R., Saux Picart, S., and Roquet, H.: Response of the sea surface temperature to heatwaves during the France 2022 meteorological summer, *Ocean Sci.*, 19, 629–647, <https://doi.org/10.5194/os-19-629-2023>, 2023.
- He, Q. and Silliman, B. R.: Climate change, human impacts, and coastal ecosystems in the Anthropocene, *Curr. Biol.*, 29, R1021–R1035, <https://doi.org/10.1016/j.cub.2019.08.042>, 2019.
- Heilskov, A. C., Alperin, M., and Holmer, M.: Benthic fauna bio-irrigation effects on nutrient regeneration in fish farm sediments, *J. Exp. Mar. Biol. Ecol.*, 339, 204–225, <https://doi.org/10.1016/j.jembe.2006.08.002>, 2006.
- Hines, A. H.: Ecology of Juvenile and Adult Blue Crabs, in: *The Blue Crab: Callinectes sapidus*, edited by: Kennedy, V. S. and Cronin, L. E., Maryland Sea Grant College, College Park, MD, 565–654, ISBN 978-0-943676-67-8, 2007.
- Hines, A. H., Johnson, E. G., Young, A. C., Aguilar, R., Kramer, M. A., Goodison, M., and Zohar, Y.: Release strategies for estuarine species with complex migratory life cycles: stock enhancement of Chesapeake blue crabs (*Callinectes sapidus*), *Rev. Fish. Sci.*, 16, 175–185, <https://doi.org/10.1080/10641260701678090>, 2008.
- Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C. J., Benthuyssen, J. A., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Moore, P. J., Scannell, H. A., Sen Gupta, A., and Wernberg, T.: A hierarchical approach to defining marine heatwaves, *Prog. Oceanogr.*, 141, 227–238, <https://doi.org/10.1016/j.pocean.2015.12.014>, 2016.
- Joyce, P. W. S., Tong, C. B., Yip, Y. L., and Falkenberg, L. J.: Marine heatwaves as drivers of biological and ecological change: implication of current research patterns and future opportunities, *Mar. Biol.*, 171, 20, <https://doi.org/10.1007/s00227-023-04340-y>, 2024.
- Jumars, P. A., Dorgan, K. M., and Lindsay, S. M.: Diet of worms emended: an update of polychaete feeding guilds, *Annu. Rev. Mar. Sci.*, 7, 497–520, <https://doi.org/10.1146/annurev-marine-010814-020007>, 2015.
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M. E., Öztürk, B., Grabowski, M., Golani, D., and Cardoso, A. C.: Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review, *Aquat. Invasions*, 9, 391–423, <https://doi.org/10.3391/ai.2014.9.4.01>, 2014.

- Katsanevakis, S., Rilov, G., and Edelist, D.: Impacts of marine invasive alien species on European fisheries and aquaculture—plague or boon?, *CIESM Monograph*, 50, 125–132, 2018.
- Kennedy, V. S., Oesterling, M., and Van Engel, W. A.: History of Blue Crab Fisheries on the US Atlantic and Gulf Coasts, in: *The Blue Crab: Callinectes sapidus*, edited by: Kennedy, V. S. and Cronin, L. E., Maryland Sea Grant College, 655–710, ISBN 978-0-943676-67-8, 2007.
- Kicklighter, C. E. and Hay, M. E.: Integrating prey defensive traits: contrasts of marine worms from temperate and tropical habitats, *Ecol. Monogr.*, 76, 195–215, [https://doi.org/10.1890/0012-9615\(2006\)076\[0195:IPDTCO\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2006)076[0195:IPDTCO]2.0.CO;2), 2006.
- Kubin, E., Menna, M., Mauri, E., Notarstefano, G., Mieruch, S., and Poulain, P. M.: Heat content and temperature trends in the Mediterranean Sea as derived from Argo float data, *Front. Mar. Sci.*, 10, 1271638, <https://doi.org/10.3389/fmars.2023.1271638>, 2023.
- Kullenberg, C. and Kasperowski, D.: What is citizen science? – A scientometric meta-analysis, *PLOS ONE*, 11, e0147152, <https://doi.org/10.1371/journal.pone.0147152>, 2016.
- Ladd, M. C. and Shantz, A. A.: Novel enemies—previously unknown predators of the bearded fireworm, *Front. Ecol. Environ.*, 14, 342–343, <https://doi.org/10.1002/fee.1310>, 2016.
- Lecci, R., Drudi, M., Grandi, A., Creti, S., and Clementi, E.: EU Copernicus Marine Service Product User Manual for For Mediterranean Sea Physics Reanalysis, MED SEA_MULTIYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-004.pdf> (last access: 10 February 2025), 2022.
- Libralato, S., Caccin, A., and Pranovi, F.: Modeling species invasions using thermal and trophic niche dynamics under climate change, *Front. Mar. Sci.*, 2, 29, <https://doi.org/10.3389/fmars.2015.00029>, 2015.
- Lipcius, R. N., Eggleston, D. B., Heck, K. L., Seitz, R. D., and Van Montfrans, J.: The Blue Crab: *Callinectes sapidus*, edited by: Kennedy, V. S. and Cronin, L. E., Maryland Sea Grant College, 535–564, ISBN 978-0-943676-67-8, 2007.
- Lopez, B. E., Magliocca, N. R., and Crooks, A. T.: Challenges and Opportunities of Social Media Data for Socio-Environmental Systems Research, *Land*, 8, 107, <https://doi.org/10.3390/land8070107>, 2019.
- Maicu, F., De Pascalis, F., Ferrarin, C., and Umgiesser, G.: Hydrodynamics of the Po River-Delta-Sea System, *J. Geophys. Res.-Oceans*, 123, 6349–6372, <https://doi.org/10.1029/2017JC013601>, 2018.
- Mancinelli, G., Alujević, K., Guerra, M. T., Raho, D., Zotti, M., and Vizzini, S.: Spatial and seasonal trophic flexibility of the Atlantic blue crab *Callinectes sapidus* in invaded coastal systems of the Apulia region (SE Italy): a stable isotope analysis, *Estuar. Coast. Shelf Sc.*, 198, 421–431, <https://doi.org/10.1016/j.ecss.2017.10.013>, 2017a.
- Mancinelli, G., Chainho, P., Cilenti, L., Falco, S., Kapiris, K., Katselis, G., and Ribeiro, F.: The Atlantic blue crab *Callinectes sapidus* in southern European coastal waters: Distribution, impact and prospective invasion management strategies, *Mar. Pollut. Bull.*, 119, 5–11, <https://doi.org/10.1016/j.marpolbul.2017.03.024>, 2017b.
- Mancinelli, G., Bardelli, R., and Zenetos, A.: A global occurrence database of the Atlantic blue crab *Callinectes sapidus*, *Scientific Data*, 8, 111, <https://doi.org/10.1038/s41597-021-00879-x>, 2021.
- Manfrin, C., Comisso, G., Dall'Asta, A., Bettoso, N., and Sook Chung, J.: The return of the Blue Crab, *Callinectes sapidus* Rathbun, 1896, after 70 years from its first appearance in the Gulf of Trieste, northern Adriatic Sea, Italy (Decapoda: Portunidae), *Check List*, 12, 1–7, <https://doi.org/10.15560/12.6.2006>, 2016.
- Marullo, S., Serva, F., Iacono, R., Napolitano, E., di Sarra, A., Meloni, D., and Santoleri, R.: Record-breaking persistence of the 2022/23 marine heatwave in the Mediterranean Sea, *Environ. Res. Lett.*, 18, 114041, <https://doi.org/10.1088/1748-9326/acfdcb>, 2023.
- Mastrototaro, F., d'Onghia, G., Corriero, G., Matarrese, A., and Maiorano, P.: Biodiversity of the white coral bank off Cape Santa Maria di Leuca (Mediterranean Sea): An update, *Deep-Sea Res. Pt. II*, 57, 412–430, <https://doi.org/10.1016/j.dsr2.2009.10.010>, 2010.
- Menna, M., Gačić, M., Martellucci, R., Notarstefano, G., Fedele, G., Mauri, E., and Poulain, P. M.: Climatic, Decadal, and Interannual Variability in the Upper Layer of the Mediterranean Sea Using Remotely Sensed and In-Situ Data, *Remote Sens.*, 14, 1322, <https://doi.org/10.3390/rs14061322>, 2022.
- Millikin, M. R. and Williams, A. B.: Synopsis of biological data on blue crab, *Callinectes sapidus* Rathbun, *FAO Fisheries Synopsis*, 38, <https://repository.library.noaa.gov/view/noaa/5574> (last access: 10 February 2025), 1984.
- Mistri, M., Borja, A., Aleffi, I. F., Lardicci, C., Tagliapietra, D., and Munari, C.: Assessing the ecological status of Italian lagoons using a biomass-based index, *Mar. Pollut. Bull.*, 126, 600–605, <https://doi.org/10.1016/j.marpolbul.2017.09.048>, 2018.
- Nehring, S.: Invasion History and Success of the American Blue Crab *Callinectes sapidus* in European and Adjacent Waters, in: *In the Wrong Place – Alien Marine Crustaceans: Distribution, Biology and Impacts*, edited by: Galil, B. S., Clark, P. F., and Carlton, J. T., *Invading Nature – Springer Series in Invasion Ecology*, Springer Netherlands, 607–624, <https://doi.org/10.1007/978-94-007-0591-3>, 2011.
- Oh, I. K. and Lee, S. W.: Effects of temperature on the survival and larval development of *Deiratonotus japonicus* (Brachyura, Camptandriidae) as a biological indicator, *Journal of Marine Science and Engineering*, 8, 213, <https://doi.org/10.3390/jmse8030213>, 2020.
- Pallas, P. S.: *Miscellanea zoologica: quibus novae imprimis atque obscurae animalium species describuntur et observationibus iconibusque illustrantur*, apud Sam et Joan Luchtmans, <https://www.biodiversitylibrary.org/bibliography/44341> (last access: 10 February 2025), 1766.
- Pires, A., Martins, R., Magalhães, L., Soares, A. M. V. M., Figueira, Quintino, E. V., Rodrigues A. M., and Freitas, R.: Expansion of lugworms towards southern European habitats and their identification using combined ecological, morphological and genetic approaches, *Mar. Ecol. Prog. Ser.*, 533, 177–190, <https://doi.org/10.3354/meps11315>, 2015.
- Pirro, A., Martellucci, R., Gallo, A., Kubin, E., Mauri, E., Juza, M., Notarstefano, G., Pacciaroni, M., Bussani, A., and Menna, M.: Subsurface warming derived from Argo floats during the 2022 Mediterranean marine heat wave, in: *8th edition of the Copernicus Ocean State Report (OSR8)*, edited by: von Schuck-

- mann, K., Moreira, L., Grégoire, M., Marcos, M., Staneva, J., Brasseur, P., Garric, G., Lionello, P., Karstensen, J., and Neukermans, G., Copernicus Publications, State Planet, 4-osr8, 18, <https://doi.org/10.5194/sp-4-osr8-18-2024>, 2024.
- Pisano, A., Nardelli, B. B., Tronconi, C., and Santoleri, R.: The new Mediterranean optimally interpolated pathfinder AVHRR SST Dataset (1982–2012), *Remote Sens. Environ.*, 176, 107–116, <https://doi.org/10.1016/j.rse.2016.01.019>, 2016.
- Pisano, A., Marullo, S., Artale, V., Falcini, F., Yang, C., Leonelli, F. E., and Buongiorno Nardelli, B.: New evidence of Mediterranean climate change and variability from sea surface temperature observations, *Remote Sens.*, 12, 132, <https://doi.org/10.3390/rs12010132>, 2020.
- Pisano, A., Fanelli, C., Massi, A., Tronconi, C., Cesarini, C., La Padula, F., Buongiorno Nardelli, B., and Ciani D.: Sea Surface Temperature Production Centre Mediterranean Sea and Black Sea Sea Surface Temperature Reprocessing, Sea Surface Temperature Production Centre Mediterranean Sea and Black Sea Sea Surface Temperature Reprocessing, Issue 4.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-021-022-041-042.pdf> (last access: 10 February 2025), 2024a.
- Pisano, A., Fanelli, C., Massi, A., Tronconi, C., Cesarini, C., La Padula, F., Buongiorno Nardelli, B., and Ciani D.: EU Copernicus Marine Service Product User Manual for Mediterranean Sea and Black Sea L3S and L4 SST Reprocessed Products, SST_MED_SST_L4_REP_OBSERVATIONS_010_021, Issue 4.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-021-022-041-042.pdf> (last access: 10 February 2025) 2024b.
- Raffa, F., Ludeno, G., Patti, B., Soldovieri, F., Mazzola, S., and Serafino, F.: X-band wave radar for coastal upwelling detection off the southern coast of Sicily, *J. Atmos. Sol.-Terr. Phys.*, 34, 21–31, <https://doi.org/10.1175/JTECH-D-16-0049.1>, 2017.
- Ramos, M. A. C. and Schizas, N. V.: Population structure of the fireworm *Hermodice carunculata* in the wider Caribbean, Atlantic and Mediterranean Sea, *J. Mar. Biol. Assoc. UK*, 103, e14, <https://doi.org/10.1017/S0025315422001114>, 2023.
- Riera, R., Pérez, O., Rodríguez, M., Ramos, E., and Monterroso, Ó.: Are assemblages of the fireworm *Hermodice carunculata* enhanced in sediments beneath offshore fish cages?, *Acta Oceanol. Sin.*, 33, 108–111, <https://doi.org/10.1007/s13131-014-0449-y>, 2014.
- Righi, S., Prevedelli, D., and Simonini, R.: Ecology, distribution and expansion of a Mediterranean native invader, the fireworm *Hermodice carunculata* (Annelida), *Mediterr. Mar. Sci.*, 21, 558–574, <https://doi.org/10.12681/mms.23117>, 2020.
- Righi, S., Savioli, M., Prevedelli, D., Simonini, R., and Malferari, D.: Unraveling the ultrastructure and mineralogical composition of fireworm stinging bristles, *Zoology*, 144, 125851, <https://doi.org/10.1016/j.zool.2020.125851>, 2021.
- Righi, S., Forti, L., Simonini, R., Ferrari, V., Prevedelli, D., and Mucci, A.: Novel Natural Compounds and Their Anatomical Distribution in the Stinging Fireworm *Hermodice carunculata* (Annelida), *Mar. Drugs*, 20, 585, <https://doi.org/10.3390/md20090585>, 2022.
- Rogers, T. L., Gouhier, T. C., and Kimbro, D. L.: Distinct temperature stressors acting on multiple ontogenetic stages influence the biogeography of Atlantic blue crabs, *Mar. Ecol. Prog. Ser.*, 690, 97–111, <https://doi.org/10.3354/meps14039>, 2022.
- Samperio-Ramos, G., Olsen, Y. S., Tomas, F., and Marbà, N.: Ecophysiological responses of three Mediterranean invasive seaweeds (*Acrothamnion preissii*, *Lophocladia lallemandii* and *Caulerpa cylindracea*) to experimental warming, *Mar. Pollut. Bull.*, 96, 418–423, <https://doi.org/10.1016/j.marpolbul.2015.05.024>, 2015.
- Scalici, M., Chiesa, S., Mancinelli, G., Rontani, P. M., Voccia, A., and Nonnis Marzano, F.: Euryhaline aliens invading Italian inland waters: The case of the Atlantic blue crab *Callinectes sapidus* Rathbun, 1896, *Appl. Sci.*, 12, 4666, <https://doi.org/10.3390/app12094666>, 2022.
- Schneider, A. K., Fabrizio, M. C., and Lipcius, R. N.: Reproductive phenology of the Chesapeake Bay blue crab population in a changing climate, *Front. Ecol. Evol.*, 11, 1304021, <https://doi.org/10.3389/fevo.2023.1304021>, 2024.
- Schulze, A., Grimes, C. J., and Rudek, T. E.: Tough, armed and omnivorous: *Hermodice carunculata* (Annelida: Amphinomidae) is prepared for ecological challenges, *J. Mar. Biol. Assoc. UK*, 97, 1075–1080, <https://doi.org/10.1017/S0025315417000091>, 2017.
- Seesanong, S., Seangarun, C., Boonchom, B., Laohavisuti, N., Thompho, S., Boonmee, W., and Rungrojchaipon, P.: Bio-green synthesis of calcium acetate from oyster shell waste at low cost and reducing the emission of greenhouse gases, *Sustainable Environment Research*, 33, 26, <https://doi.org/10.1186/s42834-023-00187-6>, 2023.
- Simonini, R. and Ferri, A.: Prime stime della longevità del verme di fuoco *Hermodice carunculata* (Annelida) dedotte grazie al contributo dei fotografi subacquei alla scienza partecipata, *Atti della Società dei Naturalisti e Matematici di Modena*, 153, 207–222, 2022.
- Simonini, R., Maletti, I., Righi, S., Fai, S., and Prevedelli, D.: Laboratory observations on predator–prey interactions between the bearded fireworm (*Hermodice carunculata*) and Mediterranean benthic invertebrates, *Freshwater Behaviour and Physiology*, 51, 145–158, <https://doi.org/10.1080/10236244.2018.1502031>, 2018.
- Simonini, R., Prevedelli, D., and Righi, S.: Esemplari mediterranei del verme di fuoco *Hermodice carunculata* (Annelida) catalogati nelle raccolte zoologiche di musei europei, *Atti della Società dei Naturalisti e Matematici di Modena*, 150, 145–159, <https://iris.unimore.it/handle/11380/1184301> (last access: 10 February 2025), 2019.
- Simonini, R., Righi, S., Zanetti, F., Fai, S., and Prevedelli, D.: Development and catch efficiency of an attracting device to collect and monitor the invasive fireworm *Hermodice carunculata* in the Mediterranean Sea, *Mediterr. Mar. Sci.*, 22, 706–714, <https://doi.org/10.12681/mms.26916>, 2021.
- Simonini, R., Ferri, A., Righi, S., Cenni, E., Ferrari, V., Sabia, C., and Prevedelli, D.: Potenziali applicazioni biotecnologiche dei policheti *Halla parthenopeia* (Oeononidae) e *Hermodice carunculata* (Amphinomidae), *Biologia Marina Mediterranea*, 28, 27–30, 2024.
- Stachowicz, J. J., Terwin, J. R., Whitlatch, R. B., and Osman, R. W.: Linking climate change and biological invasions: ocean warming facilitates nonindigenous species

- invasions, P. Natl. Acad. Sci. USA, 99, 15497–15500, <https://doi.org/10.1073/pnas.242437499>, 2002.
- Sun, C. C., Hurst, J. E., and Fuller, A. K.: Citizen science data collection for integrated wildlife population analyses, *Front. Ecol. Evol.*, 9, 682124, <https://doi.org/10.3389/fevo.2021.682124>, 2021.
- Sussman, M., Loya, Y., Fine, M., and Rosenberg, E.: The marine fireworm *Hermodice carunculata* is a winter reservoir and spring-summer vector for the coral-bleaching pathogen *Vibrio shiloi*, *Environ. Microbiol.*, 5, 250–255, <https://doi.org/10.1046/j.1462-2920.2003.00392.x>, 2003.
- Taylor, D. L. and Fehon, M. M.: Blue Crab (*Callinectes sapidus*) population structure in Southern New England tidal rivers: Patterns of shallow-water, unvegetated habitat use and quality, *Estuar. Coast.*, 44, 1320–1343, <https://doi.org/10.1007/s12237-020-00867-1>, 2021.
- Tiralongo, F., Crocetta, F., Riginella, E., Lillo, A. O., Tondo, E., Macali, A., and Azzurro, E.: Snapshot of rare, exotic and overlooked fish species in the Italian seas: A citizen science survey, *J. Sea Res.*, 164, 101930, <https://doi.org/10.1016/j.seares.2020.101930>, 2020.
- Tiralongo, F., Villani, G., Arciprete, R., and Mancini, E.: Filling the gap on Italian records of an invasive species: first records of the Blue Crab, *Callinectes sapidus* Rathbun, 1896 (Decapoda: Brachyura: Portunidae), in Latium and Campania (Tyrrhenian Sea), *Acta Adriat.*, 61, 99–104, <https://doi.org/10.32582/aa.62.1.8>, 2021.
- Tiralongo, F., Marino, S., Ignoto, S., Martellucci, R., Lombardo, B. M., Mancini, E., and Scacco, U.: Impact of *Hermodice carunculata* (Pallas, 1766) (Polychaeta: Amphinomidae) on artisanal fishery: A case study from the Mediterranean Sea, *Mar. Environ. Res.*, 192, 106227, <https://doi.org/10.1016/j.marenvres.2023.106227>, 2023.
- Tiralongo, F., Nota, A., Di Pasquale, C., Muccio, E., and Felici, A.: Trophic interactions of *Callinectes sapidus* (Blue Crab) in Vendicari Nature Reserve (Central Mediterranean, Ionian Sea) and first record of *Penaeus aztecus* (Brown Shrimp), *Diversity*, 16, 724, <https://doi.org/10.3390/d16120724>, 2024a.
- Tiralongo, F., Marcelli, M., Anselmi, G., Gattelli, R., and Felici, A.: Invasion of freshwater systems by the Atlantic blue crab *Callinectes sapidus* Rathbun, 1896 – new insights from Italian regions, *Acta Adriat.*, 65, 193–204, <https://doi.org/10.32582/aa.65.2.7>, 2024b.
- Tirelli, V., Goruppi, A., Riccamboni, R., and Tempesta, M.: Citizens' Eyes on Mnemiopsis: How to Multiply Sightings with a Click!, *Diversity*, 13, 224, <https://doi.org/10.3390/d13060224>, 2021.
- Toivonen, T., Heikinheimo, V., Fink, C., Hausmann, A., Hiippala, T., Karhu, M., Orellana, J., and Tenkanen, H.: Social media data for conservation science: A methodological overview, *Biol. Conserv.*, 233, 298–315, <https://doi.org/10.1016/j.biocon.2019.02.013>, 2019.
- Toso, A., Boulamail, S., Lago, N., Pierri, C., Piraino, S., and Giangrande, A.: First description of early developmental stages of the native invasive fireworm *Hermodice carunculata* (Annelida, Amphinomidae): a cue to the warming of the Mediterranean Sea, *Mediterr. Mar. Sci.*, 21, 442–447, <https://doi.org/10.12681/mms.22043>, 2020.
- Toso, A., Furfaro, G., Fai, S., Giangrande, A., and Piraino, S.: A sea of fireworms? New insights on ecology and seasonal density of *Hermodice carunculata* (Pallas, 1766) (Annelida) in the Ionian Sea (SE Italy), *European Zoological Journal*, 89, 1104–1114, <https://doi.org/10.1080/24750263.2022.2113156>, 2022.
- Toso, A., Mammone, M., Rossi, S., Piraino, S., and Giangrande, A.: Effect of temperature and body size on anterior and posterior regeneration in *Hermodice carunculata* (Polychaeta, Amphinomidae), *Mar. Biol.*, 171, 152, <https://doi.org/10.1007/s00227-024-04468-5>, 2024.
- Tsirintanis, K., Azzurro, E., Crocetta, F., Dimiza, M., Frogia, C., Gerovasileiou, V., Langeneck, J., Mancinelli, G., Rosso, A., Stern, N., Triantaphyllou, M., Tsiamis, K., Turon, X., Verlaque, M., Zenetos, A., and Katsanevakis, S.: Bioinvasion impacts on biodiversity, ecosystem services, and human health in the Mediterranean Sea, *Aquat. Invasions*, 17, 308–352, <https://doi.org/10.3391/ai.2022.17.3.01>, 2022.
- Turolla, E.: La venericoltura in Italia. Estado actual de cultivo y manejo de moluscos bivalvos y su proyección futura: factores que afectan su sustentabilidad en América Latina, edited by: Lovatelli, A., Fariás, A., and Uriarte, I., Taller Técnico Regional de la FAO, 20–24 August 2007, Puerto Montt, Chile, FAO Actas de Pesca y Acuicultura, No. 12. Roma, FAO, 177–188, ISBN 978-92-5-306115-0, 2008.
- Turrini, T., Dörler, D., Richter, A., Heigl, F., and Bonn, A.: The threefold potential of environmental citizen science – Generating knowledge, creating learning opportunities and enabling civic participation, *Biol. Conserv.*, 225, 176–186, <https://doi.org/10.1016/j.biocon.2018.06.003>, 2018.
- Valdes, L., Alvarez-Ossorio, M. T., and Gonzalez-Gurriaran, E.: Influence of temperature on embryonic and larval development in *Necora puber* (Brachyura, Portunidae), *J. Mar. Biol. Assoc. UK*, 71, 787–789, <https://doi.org/10.1017/S0025315400053455>, 1991.
- van den Brink, A. M., McLay, C. L., Hosie, A. M., and Dunnington, M. J.: The effect of temperature on brood duration in three *Haliscarcinus anger* species (Crustacea: Brachyura: Hymenosomatidae), *J. Mar. Biol. Assoc. UK*, 92, 515–520, <https://doi.org/10.1017/S0025315411000579>, 2012.
- Vetrano, A., Napolitano, E., Iacono, R., Schroeder, K., and Gasparini, G. P.: Tyrrhenian Sea circulation and water mass fluxes in spring 2004: Observations and model results, *J. Geophys. Res.-Oceans*, 115, C06011, <https://doi.org/10.1029/2009JC005837>, 2010.
- Wolf, A. T. and Nugues, M. M.: Predation on coral settlers by the corallivorous fireworm *Hermodice carunculata*, *Coral Reefs*, 32, 227–231, <https://doi.org/10.1007/s00338-012-0969-x>, 2013.
- Wolf, A. T., Nugues, M. M., and Wild, C.: Distribution, food preference, and trophic position of the corallivorous fireworm *Hermodice carunculata* in a Caribbean coral reef, *Coral Reefs*, 33, 1153–1163, <https://doi.org/10.1007/s00338-014-1184-8>, 2014.
- Zenetos, A., Çinar, M. E., Pancucci-Papadopoulou, M. A., Harmelin, J. G., Furnari, G., Andaloro, F., Bellou, N., Streftaris, N., and Zibrowius, H.: Annotated list of marine alien species in the Mediterranean with records of the worst invasive species, *Mediterr. Mar. Sci.*, 6, 63–118, <https://doi.org/10.12681/mms.186>, 2005.