



1 Coastal ocean response during the unprecedented marine heat 2 waves in the western Mediterranean in 2022

3 Mélanie Juza¹(mjuza@socib.es), Marta de Alfonso², Ángels Fernandez-Mora¹

4 ¹ Balearic Islands Coastal Observing and Forecasting System (SOCIB), Palma, Spain

5 ² Puertos del Estado, Madrid, Spain

6 **Abstract**

7 The western Mediterranean Sea suffered unprecedented marine heat waves (MHWs) in 2022.
8 This study focuses on the response of coastal ocean, which is highly vulnerable to global
9 warming and extreme events that threaten the biodiversity, as well as goods and services that
10 humans rely on. Using remote sensing and in situ observations, strong spatio-temporal
11 variations of MHWs characteristics are observed in the coastal ocean over the last decade 2013-
12 2022. In 2022, shallow-water moorings in the western Mediterranean Sea detected between 23
13 and 131 days of MHWs. While the highest MHW mean and maximum intensities were detected
14 at surface in the French waters, the highest duration was observed near-shore at 17 m depth in
15 the Balearic Islands. As thermal stress indicators for marine ecosystems, the cumulative
16 intensity and total days were found the highest at surface at Tarragona, and MHW temperatures
17 warmer than 28°C were observed up to 58 days at Palma. Differences between satellite products
18 and moorings observations, as well as between daily and hourly in situ measurements are also
19 highlighted inviting us to continue the efforts in deploying and maintaining multi-platform
20 observing systems in both open and coastal ocean waters to better address the coastal adaptation
21 and mitigation in the context of climate change.

22 **Introduction**

23 The Mediterranean Sea is one of the most vulnerable regions to climate change and responds
24 rapidly to global warming with strong spatial variations (Giorgi, 2006; Lionello and Scarascia,
25 2018; Juza and Tintoré, 2021a; Juza et al., 2022). In 2022, the western Mediterranean Sea
26 (WMed) suffered extreme ocean temperatures and several marine heat waves (MHWs) in a
27 row from May to December 2022 (Juza and Tintoré, 2020, 2021b). These MHWs were
28 exceptional for their early occurrence, intensity, duration and spatial extent. In the Balearic
29 Islands region, the warmest spatially-averaged satellite sea surface temperature (SST) ever
30 registered since 1982 was observed on the 13th of August 2022 with a value of 29.2 °C,
31 corresponding to an anomaly of 3.3 °C with respect to the period 1982-2015, exceeding the
32 previous regional record in summer 2003 (Juza and Tintoré, 2020, 2021b). Warmer



33 temperatures and anomalies can be found locally than regionally due to their strong spatial
34 variations (Juza and Tintoré, 2021a). In summer 2022, ocean temperatures reaching more than
35 32°C were observed in the Mallorca Channel¹, while SST anomalies exceeded 5°C in French
36 waters, reaching historical records ever registered since 1982 (Guinaldo et al., 2023).

37 The Mediterranean Sea is the largest semi-enclosed sea, with 46.000 km of coastline
38 and many islands, being also considered a hot-spot of biodiversity with many endemic species
39 (Coll et al., 2010). Its coastal zone provides goods and services that humans rely on (Smith et
40 al., 2021; UNEP/MAP and Plan Bleu, 2020) but it concentrates and accumulates human
41 pressures (e.g. contamination, population in cities, overfishing, coastline artificialization,
42 marine traffic, offshore industry and tourism) (UNEP/MAP and Plan Bleu, 2020). In addition,
43 the coastal areas and ecosystems are highly vulnerable to global warming and extreme
44 temperature events that threaten the biodiversity in the Mediterranean Sea (Cerrano et al., 2000;
45 Garrabou et al., 2009, 2019, 2022; Bensoussan et al., 2019; Verdura et al., 2019). Recently,
46 Garrabou et al. (2022) have shown that MHWs drive recurrent mass mortalities of marine
47 organisms in the Mediterranean Sea. These mass mortality events affected thousands of
48 kilometres of coastline from the surface to 45m, across a range of marine habitats and taxa.
49 Also, *Posidonia Oceanica*, which is the dominant seagrass in the Mediterranean Sea living
50 between surface and 40m depth, is very sensitive to high temperatures above 27°C, particularly
51 in its early stage of development (Guerrero-Meseguer et al., 2017). Verdura et al. (2021) also
52 highlighted during the 2015 event high mortalities of habitat-forming seaweeds at temperatures
53 of 28°C with most severe implications for early life stage and fertility. In 2017, concomitant
54 with the thermal context, the large-scale and long-lasting mucilaginous benthic algal bloom
55 was observed along the coasts of the northern Catalan Sea affecting benthic coastal habitats
56 (Bensoussan et al., 2017).

57 The climate signal manifests differently from coastal areas to the open ocean and in the
58 different sub-regions due to the variety and complexity of coastal ocean processes (Juza et al.,
59 2022). Satellite products and *in situ* measurements are complementary ocean data sources.
60 There is a benefit of using *in situ* data as a complement of satellite products since they provide
61 a more accurate representation of the thermal characteristics in the near-shore environment
62 (Schlegel et al., 2017a). Satellite data are not always accurate close to the land and have a lower
63 temporal resolution. In this study, the coastal ocean response to the unprecedented MHWs that

¹ SOCIB news [10-08-2022]: https://www.socib.es/index.php?seccion=detalle_noticia&id_noticia=535, last access: 19 June 2023.



64 occurred in the WMed in 2022 is analysed using daily data from satellite observations and
65 coastal mooring measurements. Then, the events detected by moorings in 2022 are compared
66 to those observed over the last decade since 2013. In addition, since MHW events are addressed
67 in coastal areas where ecosystems are highly present and sensitive, the range of temperatures
68 reached during these events is also studied, in particular MHW temperatures exceeding 28°C,
69 when strongly altering marine habitat and accelerating species mortality. Finally, these extreme
70 temperature ranges are investigated through the analyses of daily and hourly data highlighting
71 differences in thermal stress estimations.

72 **Datasets and methodology**

73 Datasets

74 Daily reprocessed (REP) and near real-time (NRT) satellite products in the Mediterranean Sea
75 distributed by the Copernicus Marine Service² are used (products ref. no. 1 and 2, Table 1).
76 These products provide optimally interpolated estimates of SST into regular horizontal grids
77 of 1/20° and 1/16° spatial resolutions, respectively, covering the period 1982-2022 (Pisano et
78 al., 2016; Buongiorno Nardelli et al., 2013).

79 Hourly temperature timeseries from moorings in the WMed were uploaded from the
80 Copernicus Marine In Situ data portal³ (product ref. no. 3, Table 1) and the Balearic Islands
81 Coastal Observing and Forecasting System (SOCIB) data catalogue⁴ (products ref. no. 4 and
82 5, Table 1). Fixed stations with data covering the period 2013-2022 with limited temporal gaps
83 have been selected. In addition, focusing the study on the coastal response to extreme
84 temperature events, deep water stations (off the continental shelf) have been excluded. A total
85 of 10 coastal moorings located at depths shallower than 200 m are used in this study (Table 2,
86 Figure 1). Finally, all moorings data were post-processed removing spikes and erroneous data.

87 Methodology

88 The commonly used methodology for MHW identification and characterization from Hobday
89 et al. (2016) is applied. MHWs correspond to daily SSTs exceeding the daily 90th percentile of
90 the local SST distribution over a long-term reference period during at least five consecutive
91 days. In addition, two successive MHW events with 2-day or less time break are considered as
92 a continuous event. This also allows discarding the unrealistic jumps in SST time series due to
93 sparse erroneous daily interpolated data in the NRT satellite product or in temperature time

² Copernicus Marine Service: <https://marine.copernicus.eu/>, last access: 19 June 2023.

³ CMEMS In Situ TAC: <http://www.marineinsitu.eu/>, last access: 19 June 2023.

⁴ SOCIB thredds catalog: <https://thredds.socib.es/thredds/catalog.html>, last access: 19 June 2023.



94 series from *in situ* measurements. Finally, the daily climatological mean and threshold time
95 series are smoothed using a 30-day moving window to extract useful climatology from
96 inherently variable data.

97 First, daily SST from satellites are used to compute climatology over the period 1982-2015 and
98 to detect MHWs from 1982 to 2022, providing valuable information about the 2022 thermal
99 situation over the whole Mediterranean. The chosen reference period starts as early as possible,
100 covers at least a 30-year period as recommended (Hobday et al., 2016) and is aligned with the
101 methodology applied in recent publications in the Mediterranean Sea (Juza and Tintoré, 2021;
102 Juza et al., 2022). Then, the computation and detection are applied to the daily mean
103 temperature timeseries from mooring and the nearest satellite point when *in situ* data are
104 available, both over the commonly available period 2013-2022 for their direct comparison.
105 Although the *in situ* time series are shorter than the recommended 30-year minimum for the
106 calculation of climatology and characterization of MHWs, the calculation of MHWs using their
107 own climatology allows quantifying the amount they differ from their localities (Schlegel et
108 al., 2017b; Juza et al., 2022).

109 MHW indices are then calculated to characterize the 2022 MHW event and to estimate changes
110 over the last decade. For each year, the MHW mean and maximum intensities above the mean
111 climatology, mean duration and number of discrete events are computed. MHW cumulative
112 intensity and total days are also provided as interesting indicators for ecosystem stressor,
113 although they are an aggregation of MHW intensity and duration, and of duration and
114 frequency, respectively. Finally, ocean temperatures exceeding 28°C are also identified during
115 the detected MHW events. The combination of abnormal conditions (MHW) and stressful
116 threshold (temperature ranges) allows identifying high thermal stress situations that strongly
117 impact marine ecosystems. In this respect, these extreme temperatures are also investigated
118 through the use and analysis of hourly data as observed by the moorings.

119 **MHWs in the Mediterranean Sea**

120 MHWs are firstly detected using satellite SST with respect to the reference period 1982-2015.
121 MHW characteristics are quantitatively sensitive to the baseline period but remain qualitatively
122 consistent (Dayan et al., 2023). All MHW characteristics are substantially increasing in the
123 Mediterranean Sea over the last decades, as studied over 1982-2020 (Juza et al., 2022), 1987-
124 2019 (Dayan et al., 2023) and 1982-2021 (Pastor and Khodayar, 2023). Over the recent period
125 1982-2022, the local trend estimates with 95% confidence for the MHW characteristics have



126 reached maximum values of MHW mean and maximum intensities, mean duration, frequency
127 and total days of 0.18 and 0.65°C/decade, 12.4 days/decade, 2.4 events/decade and 42.2
128 days/decade, respectively (Juza and Tintoré, 2021b, Vargas-Yáñez et al., 2023). In 2022,
129 annual mean and maximum intensities, mean duration, frequency and total days in the whole
130 Mediterranean oscillate locally over 0.95-3.10 and 1.24-6.47°C, 5-235 days, 1-15 events and
131 5-291 days, respectively (Figure 2A for MHW total days). In 2022, there are strong differences
132 in MHW characteristics between the western and eastern sub-basins. In the WMed,
133 unprecedented MHWs occurred in 2022 which was the year with the highest annual total days
134 of MHWs over the period 1982-2022 reaching up to 291 days locally along the Spanish coast
135 in the Balearic Sea (Figure 2A). Spatially integrated in the WMed, annual MHW characteristics
136 reached records ever registered since 1982 during the year 2022 (Figure 2B for MHW total
137 days). In particular, mean and maximum intensities, mean duration and total days reached 2.25
138 and 4.36°C, 36.6 and 180 days, respectively.

139 **Coastal MHWs in 2022**

140 MHWs are then detected from daily temperature from mooring and satellite with respect to the
141 reference period 2013-2022, which is the longest common period available in the moorings of
142 study. The use of shorter time series for climatology induces errors in MHW detection and
143 characterization, in particular due to ocean warming trend (Juza et al., 2022; Izquierdo et al.,
144 2022). More precisely, MHW characteristics detected by satellites at the nearest point from
145 moorings differ according to the reference period used (not shown). Since the SST
146 climatologies have higher values over 2013-2022 than 1982-2015, fewer MHW events are
147 detected using the 2013-2022 reference period. More specifically, annual MHW total days,
148 maximum and cumulative intensities are underestimated by at least 21, 5 and 29%,
149 respectively, according to the year and mooring location over 2013-2022, and up to 100% some
150 years when MHWs are not detected with the recent and short reference period for climatology
151 (Table 3).

152 **Results from moorings**

153 In 2022, all moorings of the WMed detected MHWs over the last decade (Figure 3), although
154 MHWs were computed using the reference period 2013-2022. As mentioned above, the use of
155 recent baseline periods underestimates these extreme events (Table 3) due to ocean warming.
156 Different responses are highlighted between the moorings (Figure 3, Table 4), not only because
157 of the different depths of sensor installation but also because of their geographical location.



158 Indeed, results from satellite data at the nearest point also indicate the strong spatial variability.
159 In 2022, the highest mean and maximum intensities of MHWs detected by moorings are found
160 along the French coast (Sète and Leucate) and the southern Spanish coast (Malaga) up to 3.67
161 and 5.17°C, respectively. The highest mean duration is detected in the near-shore moorings at
162 Cala Millor (40 days) and Son Bou (31 days) installed at 17 m depth, as well as in the coastal
163 Balearic Sea (Tarragona, Dragonera and Palma) where the highest total days is observed with
164 values up to 131 days at Tarragona in 2022. Such responses have led to highest cumulative
165 intensity and possibly associated thermal stress on ecosystems in the moorings at Palma,
166 Dragonera, Tarragona, Sète and Leucate. Finally, MHW days with temperature exceeding 28°C
167 are found in the Balearic Sea, from Barcelona to Cala Millor and Son Bou, with the highest
168 numbers at Tarragona (47), Dragonera (53) and Palma (58). In addition, these highly stressful
169 thermal situations with temperatures higher than 28°C occurred several times during the
170 summer 2022 with long periods of consecutive days (up to 33 days at Palma). Moorings located
171 along the French coast (Leucate and Sète) and in the Alboran Sea (Malaga and Melilla) did not
172 face daily temperatures warmer than 28°C.

173 Differences with satellite

174 Differences between moorings and satellites are found in all locations although the satellite
175 points are very close to corresponding moorings (Table 4). In 2022, along the French coast,
176 moorings observed higher MHW mean intensity at Sète and Leucate (by 0.39 and 0.23°C,
177 respectively) and higher MHW maximum intensity at Leucate (by 1.47°C) than satellites. On
178 the contrary, satellites detected higher MHW mean and maximum intensity at Barcelona than
179 moorings, with differences around 0.5 and 1.07°C, respectively. Strong differences in MHW
180 maximum intensities are also found at Melilla, Palma and Son Bou (by 1.13, 0.53 and 0.52°C
181 respectively). The MHW mean duration is found longer in moorings than satellites particularly
182 at Cala Millor, Son Bou and Tarragona (by 15, 10.3 and 7.9 days, respectively) while it is
183 particularly longer in satellites than in moorings at Dragonera and Palma (by 8.3 and 13.4 days,
184 respectively). The MHW total days and cumulative intensity in 2022 are higher in moorings at
185 Sète and Tarragona than in satellites at the nearest point while they are found higher in satellites
186 at Leucate, Barcelona, Balearic Islands stations (particularly at Cala Millor and Son Bou) and
187 Melilla. Finally, where MHW days with temperatures warmer than 28°C are found (from
188 Barcelona to Son Bou), the number of days is higher in satellites than in mooring, except at
189 Tarragona.



190 Differences between MHWs detected by satellites and moorings may be explained by the depth
191 of measurements. While satellites provide SST, the selected moorings collected temperatures
192 at surface or subsurface (from 0.4 to 17 m depths, Table 2). However, even for moorings with
193 sensors installed near the surface (up to 0.5 m), strong differences with satellites are pointed
194 out as found at Sète, Leucate and Barcelona for MHW mean and maximum intensities (up to
195 0.5 and 1.47 °C, respectively), and at Tarragona for MHW mean duration (13.4 days). Also,
196 importantly, results at Cala Millor and Son Bou strongly differ between satellites at the surface
197 and moorings in subsurface (particularly in MHW total days and days with temperature warmer
198 than 28°C), as well as, between satellite locations and between moorings highlighting the
199 coastal ocean response differ from surface to subsurface and from one location to another at
200 both surface and subsurface even in the same sub-region (on each side of the Menorca Channel
201 in the Balearic Islands).

202 **Coastal MHWs from 2013 to 2022**

203 MHWs observed by the moorings are now analysed from 2013 to 2022 and the events in 2022
204 are compared with those over the last decade (Figure 4). All years over 2013-2022 suffered
205 MHWs in several locations of the coastal WMed. In 2020 and 2022, all moorings detected
206 MHWs. While 2020 events mostly happened in winter, 2022 MHWs mainly occurred in
207 summer reaching high ocean temperatures.

208 Time series of annual MHW characteristics from moorings show strong spatio-temporal
209 variability. Variations in MHW mean and maximum intensities are highlighted between years
210 while the increase in MHW frequency and duration in recent years leads to a clear increase in
211 MHW total days and cumulative intensity. In recent years, MHWs did not only occur during
212 their usual season over a longer period but also extended over more seasons. When one season
213 was concerned in 2013, MHW occurrences covered three seasons in 2022 (not shown).

214 The analysis over the period 2013-2022 highlights that many thermal records were reached in
215 2022. MHW total days reached the highest number in 2022 for the stations at Leucate,
216 Barcelona, Tarragona, Dragonera, Palma, Cala Millor, the second highest at Sète, Son Bou,
217 Melilla and the fourth highest at Malaga. The MHW cumulative intensity in 2022 is the
218 warmest observed since 2013 for the stations at Leucate, Barcelona, Tarragona, Dragonera,
219 Palma, Cala Millor, Melilla, the second warmest at Sète and Son Bou, and the third warmest at
220 Malaga. In addition, in 2022, the number of MHW days with temperatures exceeding 28°C is
221 the highest and can be considered as the unique year until now for the moorings at Barcelona,



222 Tarragona, Dragonera, Palma, Cala Millor, Son Bou, although Palma and Tarragona also
223 experienced 7 and 5 days, respectively, with such warm temperatures in 2015.

224 **Discussion**

225 Hourly measurements from moorings were averaged on a daily basis to be compared with the
226 daily satellite products. The associated standard deviations over 2013-2022 oscillate between
227 0.23 and 0.39 °C depending on the stations. In this section, the temporal resolution impact on
228 the estimation of thermal stress during MHW events is analysed, in particular when high
229 temperatures of 28°C or more are reached. As highlighted above, the MHW events concerned
230 are those in 2022 at the moorings from Barcelona to Son Bou.

231 Due to the diurnal cycle, maxima of MHW temperatures are found in the hourly datasets
232 (Figure 5). While the maxima from the daily datasets vary between 28.37°C (Barcelona) and
233 29.95°C (Palma), in the hourly datasets they oscillated between 28.96°C (Cala Millor) and
234 31.36°C (Dragonera), this latter being the record ever registered by the Spanish mooring
235 network from Puertos del Estado. The difference between the daily and hourly data maxima is
236 the highest at Dragonera (1.52°C) and the smallest at Palma (0.05°C). The distribution of the
237 temperatures higher than 28°C is schematically represented by the median, as well as the 5 and
238 95th percentiles whose difference allows estimating the width (Figure 5). This latter is larger in
239 the hourly than daily datasets due to the diurnal cycle. Comparing the moorings between
240 themselves, the width is larger in both daily and hourly datasets at Dragonera (1.34 and 1.56°C,
241 respectively), Palma (1.33 and 1.42°C, respectively) and Tarragona (1.07 and 1.30°C,
242 respectively) where warmer temperatures were reached.

243 At Palma, the daily and hourly data provide similar results on the maxima reached and
244 distribution characteristics of extreme ocean temperatures in summer. At the moorings located
245 further off the coast of peninsula (Barcelona, Tarragona and Dragonera), the temporal
246 resolution of *in situ* data clearly impacts the extreme temperature observations. Such findings
247 are also highlighted in the two near-shore stations although their sensors are located at 17m
248 depth.

249 **Conclusions**

250 Society is facing unprecedented challenges arising from climate change impacts. Among them,
251 marine heat waves (MHWs) are becoming more frequent, longer and more intense worldwide
252 (Frölicher et al., 2018, Oliver et al., 2018) and particularly in the Mediterranean Sea (Juza et



253 al., 2022; Dayan et al., 2023; Pastor and Khodayar, 2023). Such physical changes have major
254 ecological impacts with socio-economic implications and compromising carbon storage,
255 particularly in coastal ocean waters (Smith et al., 2021, 2023). Although MHWs are mainly
256 induced by large-scale anomalous atmospheric conditions in the Mediterranean Sea (Holbrook
257 et al., 2019; Guinaldo et al., 2023; Hamdeno and Alvera-Azcarate, 2023), the ocean response
258 strongly differs from the open ocean to near-shore areas, and from one coastal location to
259 another.

260 In this study, MHWs in the coastal and shallow waters of the western Mediterranean Sea
261 (WMed) have been investigated during the year 2022 and the period 2013-2022. Satellite and
262 moorings observed MHWs along the coast of the WMed whose characteristics strongly vary
263 in time and space. Coastal MHWs were observed almost every year over the last decade, and
264 they were exceptional in 2022 in intensity, duration and geographical extension. In 2022,
265 although the coastal MHW events have a strong spatial variation, all moorings - from northern
266 to southern WMed, from surface to subsurface - observed MHWs registering records in
267 intensity (in French waters), duration (in subsurface in the Balearic Islands), total days,
268 cumulative intensity (at Tarragona), and number of days with temperature warmer than 28°C
269 (at Dragonera and Palma).

270 Although the satellite products have the great benefit to monitor all the ocean surface,
271 differences with the moorings have been detected in the characterization of MHWs in coastal
272 areas and shallow waters. Compared with mooring measurements at surface (between 0 and
273 3m depth) in 2022, satellites underestimate MHW intensities in French waters and MHW
274 duration at Tarragona while they overestimate MHW intensities at Barcelona, Palma and
275 Melilla, as well as MHW duration at Dragonera and Palma. The thermal stress on the physical
276 and biological oceans is also minimized with the use of daily data compared to hourly
277 measurements. Finally, the ocean response to extreme warm events strongly differs from one
278 location to another even in the same region, from surface to subsurface even in very shallow
279 waters. Such findings assert the importance of multi-platform, multi-sensor and sustainable
280 ocean observing systems from open to coastal and near-shore waters and from surface to
281 subsurface to continue the investigation concerning MHWs and impact assessment.

282 **References**

283 Bensoussan, N., Chiggiato, J., Buongiorno Nardelli, B., Pisano, A., and Garrabou, J. (2019).
284 Insights on 2017 marine heat waves in the Mediterranean Sea. Copernicus marine service



- 285 ocean state report: issue 3. *J. Operational Oceanogr.* 12, 101–108,
286 [doi:10.1080/1755876X.2019.1633075](https://doi.org/10.1080/1755876X.2019.1633075)
- 287 Buongiorno Nardelli, B., Tronconi, C., Pisano, A., Santoleri, R. (2013). High and Ultra-High
288 resolution processing of satellite Sea Surface Temperature data over Southern European Seas
289 in the framework of MyOcean project. *Rem. Sens. Env.*, 129, 1-16,
290 [doi:10.1016/j.rse.2012.10.012](https://doi.org/10.1016/j.rse.2012.10.012)
- 291 Cerrano, C., Bavestrello, G., Bianchi, C. N., Cattaneo-Vietti, R., Bava, S., Morganti, C., et al.
292 (2000). A catastrophic mass-mortality episode of gorgonians and other organisms in the
293 Ligurian Sea (North-western Mediterranean), summer 1999. *Ecol. Lett.* 3, 284–293,
294 [doi:10.1046/j.1461-0248.2000.00152.x](https://doi.org/10.1046/j.1461-0248.2000.00152.x)
- 295 Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Lasram, F. B. R., Aguzzi, J., et al. (2010).
296 The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One*
297 5:e11842, [doi:10.1371/journal.pone.0011842](https://doi.org/10.1371/journal.pone.0011842)
- 298 Coma, R., Ribes, M., Serrano, E., Jiménez, E., Salat, J., and Pascual, J. (2009). Global
299 warming-enhanced stratification and mass mortality events in the Mediterranean. *Proc. Natl.*
300 *Acad. Sci.* 106, 6176–6181, [doi:10.1073/pnas.0805801106](https://doi.org/10.1073/pnas.0805801106)
- 301 Dayan, H., McAdam, R., Juza, M., Masina, S., & Speich, S. (2023). Marine heat waves in the
302 Mediterranean Sea: An assessment from the surface to the subsurface to meet national needs.
303 *Frontiers in Marine Science*, 10, 142, [doi:10.3389/fmars.2023.1045138](https://doi.org/10.3389/fmars.2023.1045138)
- 304 EU Copernicus Marine Service Product: Mediterranean Sea High Resolution and Ultra High
305 Resolution Sea Surface Temperature Analysis, Mercator Ocean International [data set],
306 <https://doi.org/10.48670/moi-00172>, 2022a.
- 307 EU Copernicus Marine Service Product: Mediterranean Sea - High Resolution L4 Sea
308 Surface Temperature Reprocessed, Mercator Ocean International [data set],
309 <https://doi.org/10.48670/moi-00173>, 2022b.
- 310 EU Copernicus Marine Service Product: Mediterranean Sea - In-Situ Near Real Time
311 Observations, Mercator Ocean International [data set], <https://doi.org/10.48670/moi-00044>,
312 2022c.
- 313 Fernández-Mora, À., Juza, M., Tintoré, J. (2021). Hourly in situ data of ocean temperature in
314 the near-shore Balearic Islands over 2012-2020 (Version 1.0.0) [Data set]. Balearic Islands
315 Coastal Observing and Forecasting System, SOCIB. <https://doi.org/10.25704/ra9h-5127>
- 316 Frölicher, T. L., Fischer, E. M., and Gruber, N. (2018). Marine heatwaves under global
317 warming. *Nature* 560, 360–364, [doi:10.1038/s41586-018-0383-9](https://doi.org/10.1038/s41586-018-0383-9)
- 318 Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonné, P., Cigliano, M., et al.
319 (2009). Mass mortality in North-western Mediterranean rocky benthic communities: effects
320 of the 2003 heat wave. *Glob. Change Biol.* 15, 1090–1103, [doi:10.1111/j.1365-
321 2486.2008.01823.x](https://doi.org/10.1111/j.1365-2486.2008.01823.x)



- 322 Garrabou, J., Gómez-Gras, D., Ledoux, J. B., Linares, C., Bensoussan, N., López-Sendino,
323 P., et al. (2019). Collaborative database to track mass mortality events in the Mediterranean
324 Sea. *Frontiers in Marine Science*, 707, [doi:10.3389/fmars.2019.00707](https://doi.org/10.3389/fmars.2019.00707)
- 325 Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Ponti, M., Schlegel, R., ... &
326 Harmelin, J. G. (2022). Marine heatwaves drive recurrent mass mortalities in the
327 Mediterranean Sea. *Global Change Biology*, 28(19), 5708-5725, [doi:10.1111/gcb.16301](https://doi.org/10.1111/gcb.16301)
- 328 Giorgi, F. (2006). Climate change hot-spots. *Geophysical research letters*, 33(8),
329 [doi:10.1029/2006GL025734](https://doi.org/10.1029/2006GL025734)
- 330 Guerrero-Meseguer, L., Marín, A., & Sanz-Lázaro, C. (2017). Future heat waves due to
331 climate change threaten the survival of *Posidonia oceanica* seedlings. *Environmental*
332 *Pollution*, 230, 40-45, [doi:10.1016/j.envpol.2017.06.039](https://doi.org/10.1016/j.envpol.2017.06.039)
- 333 Guinaldo, T., Voldoire, A., Waldman, R., Saux Picart, S., & Roquet, H. (2023). Response of
334 the sea surface temperature to heatwaves during the France 2022 meteorological summer.
335 *Ocean Science*, 19(3), 629-647, [doi:10.5194/os-19-629-2023](https://doi.org/10.5194/os-19-629-2023)
- 336 Hamdeno, M., & Alvera-Azcaráte, A. (2023). Marine heatwaves characteristics in the
337 Mediterranean Sea: Case study the 2019 heatwave events. *Frontiers in Marine Science*,
338 [doi:10.3389/fmars.2023.1093760](https://doi.org/10.3389/fmars.2023.1093760)
- 339 Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., ...
340 & Wernberg, T. (2016). A hierarchical approach to defining marine heatwaves. *Progress in*
341 *Oceanography*, 141, 227-238, [doi:10.1016/j.pocean.2015.12.014](https://doi.org/10.1016/j.pocean.2015.12.014)
- 342 Holbrook, N. J., Scannell, H. A., Gupta, A. S., Benthuyssen, J. A., Feng, M., Oliver, E. C., et
343 al. (2019). A global assessment of marine heatwaves and their drivers. *Nat. Commun.*
344 10:2624. [doi:10.1038/s41467-019-10206-z](https://doi.org/10.1038/s41467-019-10206-z)
- 345 In Situ TAC partners (2022): EU Copernicus Marine Service Product User Manual for In situ
346 Products, Issue 1.14, Mercator Ocean International,
347 [https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-INS-PUM-013-030-](https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-INS-PUM-013-030-036.pdf)
348 [036.pdf](https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-INS-PUM-013-030-036.pdf) [Accessed on 23 May 2023]
- 349 Izquierdo P., Taboada F. G., González-Gil R., Arrontes J., Rico J. M. (2022). Alongshore
350 upwelling modulates the intensity of marine heatwaves in a temperate coastal sea. *Sci. Total*
351 *Environ.* 835, 155478, [doi:10.1016/j.scitotenv.2022.155478](https://doi.org/10.1016/j.scitotenv.2022.155478)
- 352 Juza, M., & Tintoré, J. (2020). Sub-regional Mediterranean Sea Indicators [Web app].
353 Balearic Islands Coastal Observing and Forecasting System, SOCIB.
354 <https://apps.socib.es/subregmed-indicators>, last access on 1 June 2023.
- 355 Juza, M., & Tintoré, J. (2021a). Multivariate Sub-regional ocean indicators in the
356 Mediterranean Sea: From event detection to climate change estimations. *Frontiers in Marine*
357 *Science*, 8, 610589, [doi:10.3389/fmars.2021.610589](https://doi.org/10.3389/fmars.2021.610589)



- 358 Juza, M., & Tintoré, J. (2021b). Sub-regional Mediterranean Marine Heat Waves [Web app].
359 Balearic Islands Coastal Observing and Forecasting System, SOCIB.
360 <https://apps.socib.es/subregmed-marine-heatwaves>, last access on 1 June 2023.
- 361 Juza, M., Fernández-Mora, A., and Tintoré, J. (2022). Sub-regional marine heat waves in the
362 Mediterranean Sea from observations: long-term surface changes, sub-surface and coastal
363 responses, *Front. Mar. Sci.*, [doi:10.3389/fmars.2022.785771](https://doi.org/10.3389/fmars.2022.785771)
- 364 Lionello, P., & Scarascia, L. (2018). The relation between climate change in the
365 Mediterranean region and global warming. *Regional Environmental Change*, 18, 1481-1493,
366 [doi:10.1007/s10113-018-1290-1](https://doi.org/10.1007/s10113-018-1290-1)
- 367 Oliver, E. C., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V., et
368 al. (2018). Longer and more frequent marine heatwaves over the past century. *Nat. Commun.*
369 9:1324, [doi:10.1038/s41467-018-03732-9](https://doi.org/10.1038/s41467-018-03732-9)
- 370 Pastor, F., and Khodayar, S. (2023). Marine heat waves: Characterizing a major climate
371 impact in the Mediterranean. *Science of The Total Environment*, 861, 160621,
372 [doi:10.1016/j.scitotenv.2022.160621](https://doi.org/10.1016/j.scitotenv.2022.160621)
- 373 Pisano, A., Nardelli, B.B., Tronconi, C., Santoleri, R. (2016). The new Mediterranean
374 optimally interpolated pathfinder AVHRR SST Dataset (1982–2012). *Remote Sens. Environ.*,
375 176, 107-116, [doi:10.1016/j.rse.2016.01.019](https://doi.org/10.1016/j.rse.2016.01.019)
- 376 Pisano, A., Fanelli, C., Buongiorno Nardelli, B., Tronconi, C., La Padula, F., Cesarini, C.
377 (2022a): EU Copernicus Marine Service Quality Information Document for Mediterranean
378 Sea and Black Sea Surface Temperature NRT data, Issue 3.0, Mercator Ocean International,
379 <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-004-006-012-013.pdf> [Accessed on 23 May 2023]
- 381 Pisano, A., Fanelli, C., Buongiorno Nardelli, B., Tronconi, C., Cesarini, C., La Padula, F.
382 (2022b): EU Copernicus Marine Service Product User Manual for SST products over the
383 Mediterranean and Black Seas, Issue 3.0, Mercator Ocean International,
384 <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-004-006-012-013.pdf> [Accessed on 23 May 2023]
- 386 Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., Buongiorno Nardelli, B.
387 (2022c): EU Copernicus Marine Service Quality Information Document for Mediterranean
388 Sea and Black Sea Sea Surface Temperature Reprocessing, Issue 2.0, Mercator Ocean
389 International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-021-022-041-042.pdf> [Accessed on 23 May 2023]
- 391 Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., Buongiorno Nardelli, B.
392 (2022d): EU Copernicus Marine Service Product User Manual for Mediterranean Sea and
393 Black Sea L3S and L4 SST Reprocessed Products, Issue 2.0, Mercator Ocean International,
394 <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-021-022-041-042.pdf> [Accessed on 23 May 2023]



- 396 Schlegel, R. W., Oliver, E. C., Perkins-Kirkpatrick, S., Kruger, A., and Smit, A. J. (2017a).
397 Predominant atmospheric and oceanic patterns during coastal marine heatwaves. *Front. Mar.*
398 *Sci.* 4:323. [doi:10.3389/fmars.2017.00323](https://doi.org/10.3389/fmars.2017.00323)
- 399 Schlegel, R. W., Oliver, E., Wernberg, T., and Smit, A. J. (2017b). Nearshore and offshore
400 co-occurrence of marine heatwaves and cold-spells. *Prog. Oceanogr.* 151, 189–205.
401 [doi:10.1016/j.pocean.2017.01.004](https://doi.org/10.1016/j.pocean.2017.01.004)
- 402 Smith, K. E., Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., et al.
403 (2023). Biological impacts of marine heatwaves. *Annual review of marine science*, 15, 119-
404 145. [doi:10.1146/annurev-marine-032122-121437](https://doi.org/10.1146/annurev-marine-032122-121437)
- 405 Tintoré, J., Vizoso, G., Casas, B., Heslop, E., Pascual, A., Orfila, A., et al. (2013). SOCIB:
406 the Balearic Islands coastal ocean observing and forecasting system responding to science,
407 technology and society needs. *Mar. Technol. Soc. J.* 47, 101–117. [doi:10.4031/MTSJ.47.1.10](https://doi.org/10.4031/MTSJ.47.1.10)
- 408 Tintoré, J., Pinardi, N., Álvarez-Fanjul, E., Aguiar, E., Álvarez-Berastegui, D., Bajo, M., ... &
409 Zodiatis, G. (2019). Challenges for sustained observing and forecasting systems in the
410 Mediterranean Sea. *Frontiers in Marine Science*, 568. [doi:10.3389/fmars.2019.00568](https://doi.org/10.3389/fmars.2019.00568)
- 411 Tintoré, J. (2022). Buoy BahiaDePalma data (Version 1.0.0) [Data set]. Balearic Islands
412 Coastal Observing and Forecasting System, SOCIB, <https://doi.org/10.25704/S6JB-CK61>
- 413 Vargas-Yáñez, M., Moya, F., Serra, M., Juza, M., Jordà, G., Ballesteros, E., Alonso, C.,
414 Pascual, J., Salat, J., Moltó, V., Tel, E., Balbín, R., Santiago, R., Piñeiro, S., García-Martínez,
415 M.C. (2023). Observations in the Spanish Mediterranean waters: A review and update of results
416 from 30-year monitoring. *Submitted to Journal of Marine Science and Engineering, in revision.*
- 417 Verdura, J., Linares, C., Ballesteros, E., Coma, R., Uriz, M. J., Bensoussan, N., et al. (2019).
418 Biodiversity loss in a Mediterranean ecosystem due to an extreme warming event unveils the
419 role of an engineering gorgonian species. *Sci. Rep.* 9:5911, [doi:10.1038/s41598-019-41929-0](https://doi.org/10.1038/s41598-019-41929-0)
- 420 Verdura, J., Santamaría, J., Ballesteros, E., Smale, D. A., Cefalì, M. E., Golo, R., ... &
421 Cebrian, E. (2021). Local-scale climatic refugia offer sanctuary for a habitat-forming species
422 during a marine heatwave. *Journal of Ecology*, 109(4), 1758-1773, [doi:10.1111/1365-](https://doi.org/10.1111/1365-2745.13599)
423 [2745.13599](https://doi.org/10.1111/1365-2745.13599)
- 424 United Nations Environment Programme/Mediterranean Action Plan and Plan Bleu (2020).
425 State of the Environment and Development in the Mediterranean. Nairobi,
426 https://planbleu.org/wp-content/uploads/2021/04/SoED_full-report.pdf [accessed on 1 June,
427 2023]
- 428 Wehde, H., von Schuckmann, K., Pouliquen, S., Grouazel, A., Bartolome, T., Tintoré, J., De
429 Alfonso Alonso-Munoyerro, M., Carval, T., Racapé V., and the INSTAC team (2022): EU
430 Copernicus Marine Service Quality Information Document for In situ Products,
431 INSITU_MED_PHYBGCWAV_DISCRETE_MYNRT_013_035, Issue 2.2, Mercator Ocean



432 International, [https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-INS-QUID-](https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-INS-QUID-013-030-036.pdf)
433 [013-030-036.pdf](https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-INS-QUID-013-030-036.pdf) [Accessed on 23 May 2023]



434 **Tables**

Product ref. no.	Product ID & type	Data access	Documentation
1	SST_MED_SST_L4_NRT_OBSERVATIONS_010_004 (1982-2021); Satellite observations	EU Copernicus Marine Service Product, 2022a	Quality Information Document (QUID): Pisano al., (2022a) Product User Manual (PUM): Pisano et al., (2022b)
2	SST_MED_SST_L4_REP_OBSERVATIONS_010_021 (2022); Satellite observations	EU Copernicus Marine Service Product, 2022b	Quality Information Document (QUID): Pisano al., (2022c) Product User Manual (PUM): Pisano et al., (2022d)
3	INSITU_MED_PHYBGCWAV_DISCRETE_MYNRT_013_035 (2013-2022); In Situ observations	EU Copernicus Marine Service Product, 2022c	Quality Information Document (QUID): Wehde et al., (2022) Product User Manual (PUM): In Situ TAC partners (2022)
4	Buoy Bahía de Palma Physico-chemical parameters of sea water data (2013-2022); In situ observations	Balearic Islands Coastal Observing and Forecasting System (SOCIB) product, 2022	Tintoré, J. (2022)
5	Two nortek AWACs in near-shore Balearic Islands ; In situ observations (extended until 2022)	Balearic Islands Coastal Observing and Forecasting System (SOCIB) data, 2022	Fernández-Mora et al., (2021)

435 *Table 1: Product Table describing data products used in this study.*

Mooring	Nº	Location Mooring	Location Satellite	Distance (km)	Sensor depth (m)	Bathymetry (m)
Sète	1	43.37°N-3.78°E	43.35°N-3.77°E	1.8 (SSW)	0.0, 0.4 (since 2019-04-16)	32.4
Leucate	2	42.92°N-3.12°E	42.94°N-3.10°E	2.4 (NW)		38.2
Barcelona	3	41.32°N-2.21°E	41.31°N-2.23°E	2.1 (SEE)	0.5	76.8
Tarragona	4	41.07°N-1.19°E	41.06°N-1.19°E	0.8 (SW)	0.5	18.2
Dragonera	5	39.56°N-2.10°E	39.56°N-2.10°E	0.5 (NE)	3	183.4
Palma Bay	6	39.49°N-2.70°E	39.48°N-2.69°E	1.9 (SW)	1	31.8
Cala Millor	7	39.59°N-3.40°E	39.60°N-3.40°E	1.5 (NW)	17	17
Son Bou	8	39.90°N-4.06°E	39.90°N-4.06°E	0.5 (SW)	17	17
Málaga	9	36.66°N-4.44°W	36.65°N-4.44°W	1.4 (SSE)	0.5	21.3
Melilla	10	35.32°N-2.94°W	35.35°N-2.94°W	3.4 (NNE)	0.5	16.2



436 *Table 2: Characteristics of the study moorings in the western Mediterranean Sea (name,*
 437 *coordinates of the station and the nearest satellite point, their distance, sensor depth and*
 438 *bathymetry) as displayed in Figure 1. The distance is the one to the nearest satellite point and*
 439 *its orientation from the mooring.*

	Maximum Intensity	Cumulative Intensity	Total days
Sète	5-69	54-95	53-93
Leucate	15-100	52-100	50-100
Barcelona	17-100	64-100	65-100
Tarragona	16-100	58-100	56-100
Dragonera	19-100	51-100	42-100
Palma	26-100	51-100	37-100
CalaMillor	20-100	55-100	43-100
Son Bou	16-100	48-100	34-100
Málaga	8-100	29-100	21-100
Melilla	14-100	49-100	35-100

440 *Table 3. Underestimation error (in %) of annual MHW characteristics (maximum and*
 441 *cumulative intensities, total days) as detected by the nearest satellite points (products ref. no.*
 442 *1 and 2, Table 1) from moorings (products ref. no. 3, 4 and 5, Table 1) over 2013-2022 with*
 443 *respect to the reference periods 2013-2022 and 1982-2015 (reference for error estimation).*

	Mean Intensity	Maximum Intensity	Cumulative Intensity	Duration	Frequency	Total days	Total days with T>28°C [consecutive days]
Sète	<u>3.67</u> 3.28	<u>5.11</u> 5.35	146.68 118.16	10 9	4 4	40 36	- -
Leucate	<u>2.72</u> 2.49	<u>5.17</u> 3.70	212.07 221.64	9.8 14.8	<u>8</u> 6	78 89	- -
Barcelona	1.80 2.30	2.64 3.71	108.07 188.23	15 16.4	4 5	60 82	8 [6-2] 17 [1-16]
Tarragona	2.10 2.18	4.21 4.22	<u>274.48</u> 242.01	<u>21.8</u> 13.9	6 8	<u>131</u> 111	47 [11-19-11-1-4-1] 22 [2-4-15-1]
Dragonera	1.87 1.87	3.34 3.19	209.58 253.11	18.7 27	6 5	<u>112</u> 135	53 [1-9-17-26] 56 [7-24-9-10-6]
Palma	1.80 1.91	2.45 2.98	221.27 237.14	17.6 31	<u>7</u> 4	<u>123</u> 124	58 [33-25] 59 [43 10 6]
Cala Millor	1.85 1.90	3.09 3.24	147.76 237.71	<u>40</u> 25	2 5	80 125	20 [3-4-5-1-6-1] 55 [40-6-3-6]
Son Bou	1.90 1.90	2.65 3.17	117.91 235.27	<u>31</u> 20.7	2 6	62 124	8 [5-1-2] 45 [4-29-4-3-3-2]




Málaga	<u>3.51</u> 3.34	<u>4.38</u> 4.51	80.69 76.82	7.7 7.7	3 3	23 23	- -
Melilla	1.66 1.71	2.75 3.82	77.90 168.37	9.4 12.5	5 8	47 98	1 1

444 *Table 4. Annual MHW characteristics (mean, maximum and cumulative intensities, mean*
445 *duration, frequency and total days) and number of MHW days with temperature warmer than*
446 *28°C as detected by moorings (products ref. no. 3, 4 and 5, Table 1, in black) and satellite*
447 *nearest point (product ref. no. 1, Table 1, in red) in 2022.*




448 **Figures**




449


450 *Figure 1. Bathymetry (in m) in the western Mediterranean Sea with contour at 200m (grey line)*
451 *and locations of selected mooring for the study (pink points) as listed in Table 2.*




452
453 *Figure 2: (A) MHW total days in 2022 from satellite (product ref. no. 1, Table 1) with respect*
454 *to the historical data (product ref. no. 2, Table 1) over the period 1982-2015. (B) Time series*
455 *of annual MHW total days averaged in the western, central and eastern Mediterranean sub-*
456 *basins from 1982 to 2022.*



457
458 *Figure 3: (A) Daily SST and MHWs from mooring at Dragonera in 2022 with respect to the*
459 *reference period 2013-2022 (product ref. no. 3, Table 1). (B) MHW days from study moorings*
460 *(black) and satellites at the nearest point (red) during the year 2022 (products ref. no. 3, 4 and*
461 *5, Table 1).*



462
463 *Figure 4. Annual MHW characteristics (mean, maximum and cumulative intensities, mean*
464 *duration, frequency and total days) and number of MHW days with temperatures exceeding*
465 *28°C as detected by moorings (products ref. no. 3, 4 and 5, Table 1) from 2013 to 2022.*



466
467 *Figure 5: The 5, 50 and 95th percentiles and maxima of the distribution of MHW temperatures*
468 *warmer than 28°C as detected with the daily (black) and hourly (blue) data from moorings*
469 *(products ref. no. 3, 4 and 5, Table 1).*