

Forecasting the Mediterranean Sea Marine Heatwave of summer 2022

Ronan McAdam¹, Giulia Bonino¹, Emanuela Clementi¹, Simona Masina¹

¹CMCC Foundation – Euro-Mediterranean Center on Climate Change, Italy

Correspondence to: Ronan McAdam (ronan.mcadam@cmcc.it)

Abstract. Early-warning of marine heatwaves requires short-term forecasts to provide precise information on timings, local-scale coverage and intensities of coming events. Here, we describe our successful efforts to track the onset, peak and decay of the Mediterranean Sea marine heatwave of summer 2022 with the Copernicus MedFS short-term (10-day) forecast system. First, we show that the 2022 event eclipses the economically and ecologically damaging event of 2003 in terms of MHW activity (a measure of intensity and duration). Forecasts of MHW area and activity provide a means of basin-wide validation, highlighting the capability of MedFS to capture regional behaviour. On local scales, we found that the MHW occurrence in the Ligurian Sea and Gulf of Taranto, two regions of economic and ecological importance, was also reliably forecast. Encouragingly, we note that the forecast has demonstrated skill in capturing not just the season-long MHW cycle but also breaks in MHW persistence and abrupt changes in local activity. Subseasonal forecasts do not yet demonstrate the capacity to predict MHW response to short-lived weather patterns, but this study confirms that short-term forecasts, at least in the Mediterranean Sea, can fill this gap.

Short Summary

In the summer of 2022, a regional short-term forecasting system was able to predict the onset, spread, peaks and decay of a record-breaking marine heatwave in the Mediterranean Sea, up to 10 days in advance. Satellite data shows that the event was record-breaking in terms of basin-wide intensity and duration. This study demonstrates the potential of state-of-the-art forecasting systems to provide early-warning of marine heatwaves to marine activities (e.g. conservation and aquaculture).

1 Introduction

Disease outbreaks, mass mortality events and the redistribution of species induced by marine heatwaves (MHWs) lead to economic losses to fisheries and aquaculture farms, and hamper conservation efforts (Smith et al., 2021; Garrabou et al., 2022; Smith et al., 2022). The need to prepare for and mitigate these MHW-induced impacts has driven developments in understanding the drivers and predictability of MHWs (Holbrook et al., 2019; Rodrigues et al., 2019; Sen Gupta et al., 2020; Li et al., 2020; Schlegel et al., 2021; McAdam et al., 2023), and in quantifying the skill of forecasts of MHWs (Benthuyssen et

Formatted: Numbering: Continuous

Deleted: Bonino¹

Deleted: Clementi²

Deleted: ¹

Deleted: Climate Simulations & Predictions Division, Fondazione Centro Euro-Mediterraneo sui Cambiamenti Climatici -

Deleted: , Bologna,

Deleted: ²Ocean Predictions & Applications Division, Fondazione Centro Euro-Mediterraneo sui Cambiamenti Climatici - CMCC, Bologna, Italy...

Deleted: ¶

Deleted: the system

Deleted: Z

Deleted: y

al., 2021; Jacox et al., 2022; McAdam et al., 2023). The Mediterranean Sea is a particular “hot-spot” for MHWs, with much literature documenting the increases in intensity, duration, frequency and impacts (Darmaraki et al., 2019; Ibrahim et al., 2019; Juza et al., 2022; Dayan et al., 2023). Despite this, there is currently less information on forecasting capability and event predictability in the Mediterranean region than in others (e.g. the North Pacific: Jacox et al., 2019; de Boissesson et al., 2022). With marine services projected to play an ever-increasing role in global sustainability and economic security (Rayner et al., 2019), early-warning systems of heat extremes can aid their planning and day-to-day management (Hartog et al., 2023).

While inter-annual variability of MHW occurrence and characteristics is derived from ocean warming and preconditioning (de Boissesson et al., 2022), ENSO (Jacox et al., 2022) and atmospheric teleconnections (e.g. Rossby wave trains, Rodrigues et al., 2019), short-lived atmospheric processes and weather systems can disrupt MHW persistence or halt their continuation completely (Benthuyssen et al., 2021). The definition of MHWs assumes persistent conditions are harmful to marine life if the duration is 5 days or longer (Hobday et al., 2016), although this number is quite arbitrary and in principle should be species-dependent. The average duration of MHWs across most of the global ocean, as well as in the Mediterranean Sea, falls within the definition of short-term forecasting (< 2 weeks) (Oliver et al., 2021; Dayan et al., 2023). A short-term view of MHWs is therefore crucial to understanding their predictability and their impacts.

Short-term forecasting of MHWs has a range of potential roles in marine activities. While some contingency plans for extreme heat events in the aquaculture and fishing industries require several months notice (e.g. relocating or switching species), others should be performed at the latest possible moment in order to avoid minimise losses (e.g. early harvesting, or cooling of farm water) (Holsman et al., 2019; Galappaththi et al., 2020). In these cases, accurate information on daily timescales is crucial. Short-term forecasts are also useful for marine protected areas (MPAs), allowing them to prepare to monitor ecosystem damage (e.g. coral bleaching) and recovery, which in turn helps assess the effectiveness of their conservation efforts (McLeod et al., 2008). Forecasts of SST can also, in theory, be coupled to distribution models to forecast changes in species habitat for highly-mobile species (Abrahms et al., 2019). “Early-warnings” are a key means of climate resilience for marine services (Galappaththi et al., 2020); an assessment of their ability to track MHWs will contribute to further uptake by these services and unlock potential socio-economic benefits.

During the summer of 2022, the Copernicus Marine Service Mediterranean Physical Forecasting system (MedFS) was employed to monitor and forecast sea surface temperature (SST) increases which eventually evolved into a record-breaking MHW for the region. MedFS has already demonstrated skill in detecting past extreme events in the Mediterranean Sea: the “aqua alta” flooding in Venice in 2019 (Giesen et al., 2020), and Medicane Ianos (Clementi et al., 2022) and Storm Gloria (Alvarez-Fanjul et al., 2022) in 2020, providing evidence that the forecast system has ability to capture a wide range of concurrent conditions (e.g. high surface air temperatures, moisture, atmospheric instability for medicanes; Cavicchia et al., 2014). Here, we provide a basin-wide description of the event and demonstrate the ability of MedFS to accurately predict many

Deleted: y

Deleted: and

Deleted: t

Deleted: the Mediterranean Physical Forecasting system

84 facets of the event (e.g. the onset, spread, persistence and decay). First, we introduce the high-resolution regional forecast
85 system and the satellite-derived SST data used to identify MHWs. Then, the record-breaking characteristics (intensity,
86 geographic extent) of the 2022 event are described. We demonstrate the system's ability to predict the MHW spread across the
87 basin and daily temperature variability in regions of key economic and ecological importance. Finally, we explore the potential
88 role of short-term forecasting in the early-warning of MHWs compared to other forecasting time scales.

89 2. Dataset & Methods

90 Here, MHWs are detected with a 0.05° resolution reprocessing of a blend of satellite-derived products provided by the ESA
91 Climate Change Initiative (CCI) and the Copernicus Climate Change Service (C3S) initiatives, including AVHRR Pathfinder
92 dataset version 5.3 to increase the input observation coverage (Product ref. no. 1). The dataset provides daily SST of the
93 Mediterranean Sea from January 1st 1982 to present (currently, up to six months before real time).

94
95 The Mediterranean Near Real Time Analysis and Forecast is a 3D coupled hydrodynamic-wave modelling system
96 implemented at 1/24° (~4 km) horizontal spatial resolution, which produces analysis and 10-day forecasts of the main ocean
97 essential variables (Product ref. no. 2). The analysis system assimilates satellite sea level anomalies and in-situ temperature
98 and salinity observations, and nudges SST towards an ultra-high resolution satellite product. The same model framework is
99 used to provide a multi-decadal reanalysis of the ocean, extending from 1987 to the present (Product ref. no. 3). Forecasts are
100 made daily; once a week (on Tuesdays) an analysis is used to initialise forecasts, while on other days a hindcast is used. A
101 schematic of the provision of forecast and analysis data is found in the QUID/PUM of the product.

102
103 Forecasts of 2m temperature (T2M) and wind speed are obtained from the European Centre for Medium-Range Weather
104 Forecasts (ECMWF) operational forecast and analysis distributed by the Italian National Meteorological Service
105 (USAM/CNMA). Variables are available at 1 hourly resolution for the first 3 days of forecast, 3 hours for the following 3 days
106 and 6 hourly for the following 4 days. The horizontal resolution is 0.1o. The T2M anomaly is calculated using the same
107 variables in the ERA5 reanalysis, over the period 1987-2021.

108
109 MHWs are defined as SSTs which persist above the 90th-percentile for 5-days or longer (Hobday et al., 2016). Here, the 90th-
110 percentile threshold corresponds to the 33 year baseline period 1987-2021 calculated individually for satellite derived and
111 model-derived data, and smoothed with an 11-day moving window. MHWs in MedFS are defined relative to the climatology
112 of the physical reanalysis. Although there are benefits of detrending SST prior to detecting MHWs (Amaya et al., 2023), we
113 chose not to detrend in order to present the true values of temperature because they are of more relevance to species impacts
114 (e.g. Galli et al., 2017).

115

Deleted:

Formatted: Superscript

Deleted: the

Deleted: forecast system

Deleted:

120 We use the MHW activity as a means to describe the event on a basin-scale, and to validate the forecast ability to capture the
 121 spatial scale of the event. Previously, activity has been defined as the product of event intensity, duration and area over a target
 122 period (Simon et al., 2022). Here, in order to study basin-wide spread at daily resolution, we define activity as the sum of the
 123 intensity over the area undergoing a MHW in the Mediterranean Basin. We assume that all MHW activity in the basin
 124 corresponds to the same event, unlike more novel methods of MHW tracking which employ spatial clustering (Bonino et al,
 125 2023). Nonetheless, the activity metric used here identifies very similar phases of MHW activity during the 2003 event as the
 126 more advanced clustering method (Fig 1a; Bonino et al., 2023).

127

128

129 **Product Table**

Product ref. no	Product ID & type	Data Access	Documentation
1	SST_MED_SST_L4_REP_OBSERVATIONS_010_021; Satellite observations	EU Copernicus Marine Service Product (2022a)	PUM: Pisano et al. (2022a) QUID: Pisano et al. (2022b)
2	MEDSEA_ANALYSISFORECAST_PHY_006_013; Numerical models	EU Copernicus Marine Service Product (2022c)	PUM: Lecci et al. (2022b) QUID: Goglio et al. (2022)
3	MED_MULTIYEAR_PHYS_006_004; Numerical models	EU Copernicus Marine Service Product (2022b)	PUM: Lecci et al. (2022a) QUID: Escudier et al. (2022)
<u>4</u>	ECMWF Integrated Forecast System (IFS) Forecast & Analysis		
<u>5</u>	ERA5		

Formatted: English (US)

130

131 **3. Results**

132 First, we describe the event on a basin scale using satellite observations. We remind the reader that references to specific dates
133 are for indication only, as the precise timings of peaks and onsets may differ when using different datasets and climatologies.
134 In 2022, the onset of summer MHW conditions began in mid-May; by 23rd May, 35% of the area of the Mediterranean Sea
135 was already experiencing MHW conditions (Fig. 1c). Maps of temperature anomaly confirm that the onset occurred mostly in
136 the western regions and the Adriatic Sea (Fig. 2a). The geographic extent of MHW extended into the central and eastern parts
137 of the basin (e.g. Fig. 2b), and MHW area remained above a third of the total basin area until the decay at the end of September.
138 Peak area (70%) was reached on June 6th, while notable peaks of activity occurred later, on June 29th, July 6th and July 27th
139 (Fig. 1a). The peak temperature anomaly of 6.45 °C (above the 1987-2019 average) was reached in the Gulf of Lion on July
140 18th.

141
142 Prior to 2022, the MHWs of the summers of 2003, 2015 and 2018 had been found to have the highest activities on record
143 (using a slightly different definition of activity, but which is still based on intensity and duration; Simon et al., 2022). Here,
144 we find that the activity in 2022 clearly eclipses that of 2015 and 2018, in terms of both maxima and persistence of activity
145 (Fig. 1a). Though the summer of 2003 reached similarly high peaks of activity (twice, in mid-June and at the end of August),
146 the total activity during the summer (defined as the area under the curve) is lower (82 x 10⁶ °C.km²) than for 2022 (139 x 10⁶
147 °C.km²). While in 2003 the MHW activity returned to zero in late May and mid-July, in 2022 it persisted throughout the
148 summer above at least 0.5 °C.km² each day. Using the total activity as a measure, the summer of 2022 now holds the record
149 for MHW activity.

150
151 Using the MHW activity provides an efficient, if not complete, means of validating MedFS on the basin-scale. It is important
152 to remember that activity time series cannot identify where and when MHWs are occurring (we study forecast ability in specific
153 regions later). Here, we show both the activity (Fig. 1b) and the area (Fig. 1c) predicted, to infer whether forecast inaccuracies
154 are caused by an inability to capture the geographic extent or the temperature intensity. Overall, we find that MedFS was able
155 to forecast the evolution of basin-wide MHW activity (Fig. 1b). In particular, we highlight the accurate predictions of the
156 timings of the May onset, the various peaks throughout the summer, the two stages of the decay and the September rebound.
157 On several occasions, MHW activity rapidly increases, often doubling or tripling over the period of less than a week; such
158 increases in activity are predicted by the forecasts in mid-May, early-June, mid-June (twice) and mid-July. Timings of declines
159 in activity are also correctly forecast on all occasions, in some cases even with lead times of 5 days or more (e.g., early June).
160 Similar capabilities are found for the forecasts of area of MHW conditions. Ability to capture both the activity and the area
161 implies accuracy in capturing the intensity as well, although this analysis does not yet determine the geographic distribution
162 of MHW intensity.

163

Deleted: conditions

Formatted: Superscript

Deleted: overall

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Deleted: By this measure

Formatted: Superscript

Deleted: the forecast system

Deleted: the forecast system

Deleted: this

Deleted: is

Deleted: accurately

Deleted: D

Deleted: accurately

Deleted: . Moreover, the start of declines are often accurately predicted even

Deleted: The

Deleted: was also well predicted;

Deleted: a

Deleted: ccuracy

Deleted: in

Deleted: ing

Deleted: ¶

183 There are indeed forecast inaccuracies to highlight. Firstly, there are instances of peaks of activity being overestimated (e.g.,
184 by up to a third in early June) and of false alarms about growth being raised (e.g. early August; Fig. 1b). Overestimations of
185 both activity and area occur throughout the summer, except for the onset in May. Then, there are instances in which MHW
186 area tendencies follow the activity but are overestimated (late July to early August); given that activity continues to increase
187 while area decreases (e.g. early September), there is an implied overestimation of the sea surface temperatures. Elsewhere, we
188 see fluctuations in activity on daily timescales which dominate over the longer-term growth tendencies (e.g., during the growth
189 period beginning at the end of June). Throughout the summer of 2022, we see various examples of the activity forecasts being
190 unable to detect this higher-frequency variability. Variability in area, on the other hand, is lower and the forecasts generally
191 follow the observed tendencies. In summary, the forecasts sometimes persist or increase temperature anomalies for too long,
192 suggesting that they fail to capture sporadic cooling. Candidate drivers for short term cooling mechanisms not captured by the
193 forecasts include cloud cover changes or winds.

194
195 It is important also to consider the ability to capture the spread of MHW occurrence. The geographical distribution of intensity
196 in key phases of the MHW life cycle in forecasts agrees well with observations (Fig. 2). During the onset, forecasts capture
197 the basin-wide patterns, with MHW occurrence at this stage correctly forecast in the Tyrrhenian Sea, Gulf of Lion and parts
198 of the Adriatic Sea. The spread of the MHW conditions during the peak was correctly predicted to cover the south part of the
199 Alboran Sea, the Ionian and southern regions of the Levantine Basin. Meanwhile, the Aegean Sea was predicted to be shielded
200 from MHWs and instead experience cold anomalies, most likely caused by cooling related to the Etesian winds (Poupkou et
201 al., 2011). Lastly, the first decay phase at the end of August produced very inhomogeneous MHW conditions across the basin.
202 This “patchiness”, indicative of local-scale processes acting to cool the ocean such as increased cloud cover or weak winds,
203 was indeed predicted, but how well the forecast matches observations depends greatly on the local regions of interest and the
204 exact day and lead time considered. Although it is not possible to draw rigorous conclusions from snapshots, the accuracy of
205 basin-wide occurrence (Fig. 1b) suggests that forecast ability to capture MHW patterns and spread was generally high across
206 the entire summer.

Deleted: ¶

Deleted: growth

Deleted:

Deleted: Generally, these are found for longer lead times, and it is expected for a forecast to experience a decay in skill with forecast time.

Deleted: In the first half of the summer, area is accurately forecast while activity is overestimated (e.g. at the end of June), implying an overestimation of intensity. In the latter half of the summer, area is typically overestimated, partially explaining the overestimation of activity.

Deleted: ¶

¶

Deleted: forecast

Deleted: ;

Deleted: a

Deleted: , instead,

Deleted: activity

225

226

227 While basin-scale analysis allows an overview of forecasting skill, local-scale testing is imperative as forecasting tools are
 228 expected to be used on local-scale analysis (Dayan et al., 2023). Here, we also provide MHW forecasts for two key areas of
 229 maritime activity in the Mediterranean Sea: the Ligurian Sea and the Gulf of Taranto (Fig. 3). Each region experienced MHW
 230 conditions at different times during the summer, and in each case the forecasts accurately predicted the onset, persistence,
 231 intensity and decays. The Ligurian Sea, bordered by Italy and France, is a crucial location for marine conservation; it doubles
 232 as a marine protected area (the Pelagos Sanctuary for Mediterranean Marine Mammals) which is home to unique species of
 233 fin whales and striped dolphins, amongst other species (Notarbartolo-de-Sciara et al., 2008). The Ligurian Sea experienced
 234 115 days of MHWs throughout the summer, and temperature anomalies reached a maximum of 4.46^ΔC above the 1987-2019
 235 average at the end of July, coinciding with the peak temperature of the summer (28.74^ΔC). ~~The forecasts of SST were highly~~
 236 ~~accurate: root-means-squared difference (RMSE) of the forecasts shown in Fig. 3a (across lead times) was 0.12^Δ C, lower than~~
 237 ~~the reported RMSDs averaged over the entire Mediterranean Sea between the analysis and independent observations (Goglio~~
 238 ~~et al., 2023).~~ For an indication of forecast reliability, we highlight the false alarms (MHW days forecast but not observed) and
 239 misses (MHW days observed but not forecast). First we note that in the 89% of days the correct conditions are forecast with
 240 few, sporadic exceptions. For example, the forecast made on May 10th captured the sharp rise in SST but not the MHW
 241 conditions at the end of the week. However, reducing the lead time (i.e. checking forecasts made on the 12th or 13th) correctly
 242 forecast the MHW state.

243

244 The Gulf of Taranto, situated in the Northern Ionian Sea, is one of the most productive areas of shellfish (mussels) farming in
 245 Italy (Prioli, 2004) but there is not yet data on MHW-induced mass-mortality or economic loss in this region (Garrahou et al.,
 246 2019). Unlike the Ligurian Sea, the Gulf of Taranto experienced three short but intense periods of MHW ~~occurrence~~, in June
 247 and July, adding up to 61 days of MHWs in total. The peak temperature anomaly was 4.76^ΔC on 6th June, though peak
 248 temperatures occurred later in the season. As in the Ligurian Sea, the forecasts ~~were highly accurate (demonstrating a RMSD~~
 249 ~~of 0.08^ΔC).~~ ~~Regarding the reliability of MHW forecasts, the continuation of the start of the heatwave in early May was missed~~
 250 ~~by the forecast of May 17th, while the forecast of 2nd August missed several days of MHW occurrence. However, in both~~
 251 ~~cases, SST increases were predicted. None of the forecasts shown in Figure 3 raised false alarms.~~

252

253 So far we have studied accuracy of the entire forecast period but, in some applications, it might be necessary or of more interest
 254 to have a specific warning time (e.g. 4 days). As ~~MedFS~~ produces forecasts every day, we now study forecast accuracy for the
 255 summer of 2022 at different lead times; ~~we focus on the Western Mediterranean Sea, as opposed to the entire basin, in order~~
 256 ~~to investigate local forcings as reasons for poor skill (Fig. 4).~~ ~~Table 1 quantifies the error of forecasts over the Western~~
 257 ~~Mediterranean Sea.~~ The overestimation of MHW activity in July and August occurred ~~in forecasts with~~ lead time of one day.
 258 In many instances, lead time 1 and lead time 4 are similarly far from the observed values, while lead time 7 further

Formatted: Superscript

Deleted: This activity is indicative of the conditions experienced by the rest of the western part of the Mediterranean basin.

Formatted: Superscript

Formatted: Superscript

Deleted: vast majority

Deleted: activity

Formatted: Superscript

Deleted: reliably captured the MHW state, with few false alarms and misses...

Deleted: T

Formatted: Superscript

Deleted: activity

Deleted: In both cases, however, the observations show the temperature was only very slightly over the MHW threshold, and upon visual inspection the forecast temperature was very similar to the observed. ...

Deleted: 3

Deleted: the forecasts system

Deleted:

Deleted: after

Deleted: a

276 overestimates the peaks in activity (Table 1). MHW area, on the other hand, while predicted more accurately than activity on
277 all lead times, is typically underestimated by the forecasts. This implies an overestimation of SST (i.e. MHW intensity) during
278 the activity peaks. The RMSD normalised by the standard deviation, indicates where errors fall within the range of natural
279 variability (normalised RMSD < 1); in all lead times, this is true for both area and activity, suggesting their skill is similar to
280 atmospheric variables (Table 1).

281
282 The decreases in skill with lead time can partly be explained by the decrease in skill of the ECMWF atmospheric forecasts
283 used to force MedFS. T2M and wind speed correlate strongly and significantly with the MHW activity (correlation values of
284 0.89 and 0.50 with the ECWMF analysis respectively), evidencing their role in MHW formation. Errors of forecasts of T2M
285 and wind speed grow with time but do not exceed natural variability at lead time 7 (Table 1). In the first half of the summer,
286 forecasts at lead time 7 of both T2M and wind speed are frequently out of phase with the observed changes. In fact, the
287 underestimations of MHW area in this period occur simultaneously with underestimations of T2M. For example, the
288 underestimation of MHW area at the end of May, by an area of roughly 30% of the western Mediterranean, corresponds to
289 overestimations of wind speed by up to 1 m/s and temperature anomalies roughly 1°C weaker than observed. However, the
290 overestimation of activity in July and August, found to be linked to overestimations of SST, does not correspond to
291 overestimations of T2M, implying that other phenomena are not well represented. It should be noted that the use of area-
292 averaged atmospheric variables may hide sub-regional scale processes which impact the MHW location and intensity.

293 4. Discussion & Summary

294
295 The MHW of summer 2022 in the Mediterranean Sea was record-breaking, eclipsing 2003 in terms of basin-wide activity
296 (defined as the integral of intensity, duration and area). Other contributions to the Ocean State Report 8 also define the MHW
297 of 2022 as a record-breaking event, using other definitions (e.g. local SST records). Here, we provide a basin-wide view of the
298 MHW conditions. The Copernicus Mediterranean Physical forecasting system was used to track this event, serving as the first
299 validation of MHW prediction for this system. Forecasts captured the full life cycle of the MHWs several days in advance:
300 onset (mid-May) in the Western part of the basin; spread into the Adriatic and Ionian Sea; sporadic local-scale occurrences in
301 the Levantine Basin; persistence of peak conditions throughout July and August; breaks in MHW persistence and abrupt
302 changes in local occurrence; and the gradual decay (September). The forecasts also identified regions shielded from MHWs
303 e.g. during cooling in the Aegean Seas. Subseasonal forecasts do not yet demonstrate the capacity to predict MHW response
304 to weather patterns (Benthuyssen et al., 2018), but this study confirms that short-term forecasts, at least in the Mediterranean
305 Sea, can fill this gap.

Formatted: Superscript

Deleted: Other

Deleted: is report

Deleted:

Deleted: various other

Deleted: accurately

Moved (insertion) [1]

312 ~~A full analysis of potential drivers and attribution of forecast skill to certain processes was outside the scope of this study, but~~
313 ~~the dependence on accurate atmospheric forcings (here provided by ECMWF forecasts) has been shown to be crucial for~~
314 ~~accurate forecasts of the 2022 event.~~ Unlike the other **extreme** events, the common drivers of MHWs in the Mediterranean are
315 yet to be identified. The MHW of 2022, as well as the concurrent and record-breaking atmospheric heatwave which occurred
316 over western Europe, appears to be linked to the northward extension of the subtropical ridge (Barriopedro et al., 2023), **while**
317 ~~model studies have suggested that mid-latitude MHWs in summer typically arise from reduced ocean heat loss to the~~
318 ~~atmosphere and reduced vertical diffusion (Vogt et al, 2022).~~ **Here, the decay in skill of MHW forecasts match the decay in**
319 **skill of key atmospheric conditions (T2M and wind speed). Erroneous forecast of early summer, in the Western Mediterranean**
320 **Sea, are explained by inaccurate forecasts of these atmospheric conditions. Peak summer conditions, such as the overestimation**
321 **of MHW activity, are not yet understood, meaning further studies of short-term forecasting of MHWs are necessary.**

322
323 The time scale of forecasting determines the information that can be provided and the type of response to that information.
324 Here, we make the case for using short-term forecasting in MHW tracking tools and studies. Seasonal forecasting informs
325 management decisions and contingency plans, while subseasonal forecasting can update these plans (White et al, 2017). Short-
326 term forecasting, on the other hand, can then be used to determine the precise timings of events and instruct users on when to
327 implement urgent response actions. Longer-term forecasts are typically global in scale and have a relatively low model
328 resolution, while short-term forecasting centres, benefitting from the reduced time scale, can put more computational power
329 towards regional-scale forecasting at a finer scale more relevant to stakeholders. In principle, for MHWs, this means the
330 following: seasonal forecasts forewarn of extreme summer temperatures (e.g. seasonal averages above the 90th percentile,
331 identification of ocean basins affected); sub-seasonal forecasts then update this to forewarn of MHW occurrence (e.g. daily
332 temperatures persisting above 90th percentile, greater detail on geographic spread); finally, short-term forecasts can provide
333 key details such as the start date, onset rate and breaks in **occurrence**, on a local-to-regional scale. Currently, more effort is
334 being placed on seasonal forecasting of MHWs (Liu et al., 2018; Jacox et al., 2022). With the level of accuracy for local-scale
335 MHW indicators shown here, such tools should be complemented with daily, short-term updates.

336
337 In particular, we found that the MHW occurrence in the Ligurian Sea and Gulf of Taranto, two regions of economic and
338 ecological importance, was also reliably forecast. There is, though, a need to include subsurface temperatures or heat content
339 to report MHWs occurring at depth (Dayan et al., 2023; McAdam et al., 2023). For example, caged fish have been observed
340 to avoid the top of cages when surface temperatures increase (Gamperl et al., 2021), meaning truly stakeholder-relevant
341 tracking tools need a 3D view. The near-real-time analysis, as well as the forecast system, provides 3D temperatures and can
342 track subsurface propagation of MHWs (unlike satellite observations). The MHW record in the analysis aligns exceptionally
343 well with satellite observations for the two target regions shown (Fig. 3), suggesting **a high level of accuracy** (the same is
344 found for the basin-wide MHW activity; not shown). However, a subsurface validation with in-situ data **should** be performed
345 in the near-future, before using the analysis and forecast to track subsurface MHWs.

Deleted: The system has already demonstrated skill in detecting past extreme events in the Mediterranean Sea: the “aqua alta” flooding in Venice in 2019 (Giesen et al., 2020), and Mediane Ianos (Clementi et al., 2022) and Storm Gloria (Alvarez-Fanjul et al., 2022) in 2020. While these and MHWs are all considered “extreme events”, their natures and formations are quite different, meaning that, once again, the forecast system has demonstrated the ability to capture a wide range of concurrent conditions (e.g. high surface air temperatures, moisture, atmospheric instability for medicanes; Cavicchia et al, 2014).

Deleted: . Meanwhile,

Moved up [1]: Subseasonal forecasts do not yet demonstrate the capacity to predict MHW response to weather patterns (Benthuysen et al., 2018), but this study confirms that short-term forecasts, at least in the Mediterranean Sea, can fill this gap.

Deleted: ,

Deleted: but the link between these mechanisms has yet to be demonstrated in the Mediterranean. Encouragingly, however, we note that the forecast has demonstrated skill in capturing not just the season-long MHW cycle but also breaks in MHW persistence and abrupt changes in local activity.

Deleted: activity

Deleted: its

Deleted: scale

Deleted: wil

Deleted: l

372 **References**

- 373 Abrahms, B, Welch, H, Brodie, S, et al. Dynamic ensemble models to predict distributions and anthropogenic risk exposure
374 for highly mobile species. *Divers Distrib.* 2019; 25: 1182– 1193. <https://doi.org/10.1111/ddi.12940>
375
- 376 Alvarez-Fanjul E, Pérez Gómez B, de Alfonso Alonso-Muñoyerro M, et al. 2022. Western Mediterranean record-breaking
377 storm Gloria: An integrated assessment based on models and observations. In: Copernicus Ocean State Report, Issue 6, *Journal*
378 *of Operational Oceanography*, 15:sup1, s151-s159; <https://doi.org/10.1080/1755876X.2022.2095169>
379
- 380 Amaya, D.J., Miller, A.J., Xie, S.P. and Kosaka, Y., 2020. Physical drivers of the summer 2019 North Pacific marine heatwave.
381 *Nature communications*, 11(1), p.1903.
382
- 383 [Amaya, D. J., Jacox, M. G., Fewings, M. R., Saba, V. S., Stuecker, M. F., Rykaczewski, R. R., ... & Powell, B. S. \(2023\).
384 *Marine heatwaves need clear definitions so coastal communities can adapt. Nature*, 616\(7955\), 29-32.](#)
385
- 386 Barbeaux, S.J., Holsman, K. and Zador, S., 2020. Marine heatwave stress test of ecosystem-based fisheries management in the
387 Gulf of Alaska Pacific cod fishery. *Frontiers in Marine Science*, 7, p.703.
388
- 389 Barriopedro, D., García-Herrera, R., Ordóñez, C., Miralles, D.G. and Salcedo-Sanz, S., 2023. Heat waves: Physical
390 understanding and scientific challenges. *Reviews of Geophysics*, p.e2022RG000780.
391
- 392 Benthuyzen, J. A., Smith, G. A., Spillman, C. M., & Steinberg, C. R. (2021). Subseasonal prediction of the 2020 Great Barrier
393 Reef and Coral Sea marine heatwave. *Environmental Research Letters*, 16(12), 124050. doi: 10.1088/1748-9326/ac3aa1
394
- 395 Bonino, G., Masina, S., Galimberti, G. and Moretti, M., 2023. Southern Europe and western Asian marine heatwaves (SEWA-
396 MHWs): a dataset based on macroevents. *Earth System Science Data*, 15(3), pp.1269-1285.
397
- 398 Cavicchia, L., von Storch, H. and Gualdi, S., 2014. A long-term climatology of medicanes. *Climate dynamics*, 43, pp.1183-
399 1195.
400
- 401 Darmaraki, S., Somot, S., Sevault, F. and Nabat, P., 2019. Past variability of Mediterranean Sea marine heatwaves.
402 *Geophysical Research Letters*, 46(16), pp.9813-9823.
403

404 de Boisseson, E., Balmaseda, M., Mayer, M., Zuo H. 2022. Monitoring and predictions of the series of marine heatwave events
405 impacting the Northeast Pacific in 2020. In: Copernicus Ocean State Report, issue 6. Journal of Operational Oceanography,
406 15:sup1, s168-s175; <https://doi.org/10.1080/1755876X.2022.2095169>
407

408 de Burgh-Day Catherine O., Spillman Claire M., Smith Grant, Stevens Craig L. (2022) Forecasting extreme marine heat events
409 in key aquaculture regions around New Zealand. Journal of Southern Hemisphere Earth Systems Science 72, 58-72.
410 <https://doi.org/10.1071/ES21012>
411

412 Dayan, H., McAdam, R., Juza, M., Masina, S. and Speich, S., 2023. Marine heat waves in the Mediterranean Sea: An
413 assessment from the surface to the subsurface to meet national needs. *Frontiers in Marine Science*, 10, p.142.
414

415 Escudier, R., Clementi, E., Nigam, T., Aydogdu, A., Fini, E., Pistoia, J., Grandi, A., Miraglio, P.: EU Copernicus Marine
416 Service Quality Information Document for the Mediterranean Sea Physics Reanalysis,
417 MEDSEA_MULTIYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International,
418 catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-004.pdf, last access: 23 June 2023, 2022.
419

420 EU Copernicus Marine Service Product: Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed,
421 Mercator Ocean International, [data set], <https://doi.org/10.48670/moi-00173>, 2022a.
422

423 EU Copernicus Marine Service Product: Mediterranean Sea Physics Reanalysis, Mercator Ocean International, [data set],
424 https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_PHY_006_004_E3R1, 2022b.
425

426 EU Copernicus Marine Service Product: Mediterranean Sea Physics Analysis and Forecast, Mercator Ocean International,
427 [data set], https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS7, 2022c.
428

429 [Galli, G., Solidoro, C., & Lovato, T. \(2017\). Marine heat waves hazard 3D maps and the risk for low motility organisms in a](#)
430 [warming Mediterranean Sea. *Frontiers in Marine Science*, 4, 136.](#)
431

432 Gamperl, A.K., Zrini, Z.A. and Sandrelli, R.M., 2021. Atlantic salmon (*Salmo salar*) cage-site distribution, behavior, and
433 physiology during a Newfoundland heat wave. *Frontiers in Physiology*, 12, p.719594.
434

435 Galappaththi, E.K., Ichien, S.T., Hyman, A.A., Aubrac, C.J. and Ford, J.D. (2020), Climate change adaptation in aquaculture.
436 *Rev. Aquacult.*, 12: 2160-2176. <https://doi.org/10.1111/raq.12427>
437

Formatted: Italian

438 Giesen, R., Clementi, E., Bajo, M., Federico, I., Stoffelen, A., Santoleri, R. 2021. The November 2019 record high water levels
439 in Venice, Italy. Copernicus Ocean State Report, ssue 5, Journal of Operational Oceanography. 14:sup1, s156-s162,
440 <https://doi.org/10.1080/1755876X.2021.1946240>
441
442
443 Goglio, A.C., Clementi, E., Grandi, A., Mariani, A., Giurato, M., Aydogdu., A: EU Copernicus Marine Service Quality
444 Information Document for the Mediterranean Sea Physics Analysis and Forecast,
445 MEDSEA_ANALYSISFORECAST_PHY_006_013, Issue 2.3, Mercator Ocean International,
446 <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-013.pdf>, last access: 23 June 2023,
447 2022.
448
449 Hartog, J.R., Spillman, C.M., Smith, G. and Hobday, A.J., 2023. Forecasts of marine heatwaves for marine industries:
450 Reducing risk, building resilience and enhancing management responses. Deep Sea Research Part II: Topical Studies in
451 Oceanography, 209, p.105276.
452
453 Hobday, A.J., Alexander, L.V., Perkins, S.E., Smale, D.A., Straub, S.C., Oliver, E.C., Benthuyesen, J.A., Burrows, M.T., Donat,
454 M.G., Feng, M. and Holbrook, N.J., 2016. A hierarchical approach to defining marine heatwaves. Progress in Oceanography,
455 141, pp.227-238.
456
457 Holbrook, N.J., Scannell, H.A., Sen Gupta, A., Benthuyesen, J.A., Feng, M., Oliver, E.C., Alexander, L.V., Burrows, M.T.,
458 Donat, M.G., Hobday, A.J. and Moore, P.J., 2019. A global assessment of marine heatwaves and their drivers. Nature
459 communications, 10(1), p.2624.
460
461 Ibrahim, O., Mohamed, B. and Nagy, H., 2021. Spatial variability and trends of marine heat waves in the eastern mediterranean
462 sea over 39 years. Journal of Marine Science and Engineering, 9(6), p.643.
463
464 Jacox, M.G., Tommasi, D., Alexander, M.A., Hervieux, G. and Stock, C.A., 2019. Predicting the evolution of the 2014–2016
465 California Current System marine heatwave from an ensemble of coupled global climate forecasts. Frontiers in Marine Science,
466 6, p.497.
467
468 Jacox, M.G., Alexander, M.A., Amaya, D., Becker, E., Bograd, S.J., Brodie, S., Hazen, E.L., Pozo Buil, M. and Tommasi, D.,
469 2022. Global seasonal forecasts of marine heatwaves. Nature, 604(7906), pp.486-490.
470

Formatted: Italian

- 471 Juza, M., Fernández-Mora, À. and Tintoré, J., 2022. Sub-Regional marine heat waves in the Mediterranean Sea from
472 observations: Long-term surface changes, Sub-surface and coastal responses. *Frontiers in Marine Science*, 9, p.785771.
473
- 474 Holsman, K.K., Hazen, E.L., Haynie, A., Gourguet, S., Hollowed, A., Bograd, S.J., Samhouri, J.F. and Aydin, K., 2019.
475 Towards climate resiliency in fisheries management. *ICES Journal of Marine Science*, 76(5), pp.1368-
476 1378. <https://doi.org/10.1093/icesjms/fsz031>
477
- 478 Lecci, R., Drudi, M., Grandi, A., Creti, S., Clementi, E.: EU Copernicus Marine Service Product User Manual for the
479 Mediterranean Sea Physics Reanalysis, MEDSEA_MULTITYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International,
480 <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-004.pdf>, last access: 23 June 2023, 2022a.
481
482
483
- 484 Lecci, R., Drudi, M., Grandi, A., Creti, S., Clementi, E.: EU Copernicus Marine Service Product User Manual for the
485 Mediterranean Sea Physics Analysis and Forecast, MEDSEA_ANALYSISFORECAST_PHY_006_013, Issue 2.2, Mercator
486 Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-013.pdf>, last access:
487 23 June 2023, 2022b.
488
- 489 Lee, K.H., Noh, J. and Khim, J.S., 2020. The Blue Economy and the United Nations' sustainable development goals:
490 Challenges and opportunities. *Environment international*, 137, p.105528.
491
- 492 Liu, G., Eakin, C.M., Chen, M., Kumar, A., De La Cour, J.L., Heron, S.F., Geiger, E.F., Skirving, W.J., Tirak, K.V. and Strong,
493 A.E., 2018. Predicting heat stress to inform reef management: NOAA Coral Reef Watch's 4-month coral bleaching outlook.
494 *Frontiers in Marine Science*, 5, p.57.
495
- 496 McLeod, E., Salm, R., Green, A. and Almany, J. (2009), Designing marine protected area networks to address the impacts of
497 climate change. *Frontiers in Ecology and the Environment*, 7: 362-370. <https://doi.org/10.1890/070211>
498
- 499 Notarbartolo-di-Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T. and Van Klaveren, P., 2008. The Pelagos sanctuary for
500 Mediterranean marine mammals. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18(4), pp.367-391.
501
- 502 Oliver, E.C., Benthuisen, J.A., Darmaraki, S., Donat, M.G., Hobday, A.J., Holbrook, N.J., Schlegel, R.W. and Sen Gupta, A.,
503 2021. Marine heatwaves. *Annual Review of Marine Science*, 13, pp.313-342.
504

505 Poupkou, A., Zanis, P., Nastos, P., Papanastasiou, D., Melas, D., Tourpali, K. and Zerefos, C., 2011. Present climate trend
506 analysis of the Etesian winds in the Aegean Sea. *Theoretical and applied climatology*, 106, pp.459-472.
507

508 Li, Z., Holbrook, N.J., Zhang, X., Oliver, E.C. and Coughon, E.A., 2020. Remote forcing of Tasman Sea marine heatwaves.
509 *Journal of Climate*, 33(12), pp.5337-5354.
510

511 Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., Buongiorno Nardelli, B.: EU Copernicus Marine Service
512 Quality Information Document for the Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed
513 Product, SST_MED_SST_L4_REP_OBSERVATIONS_010_021, Issue 2.0, Mercator Ocean International,
514 <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-021-022-041-042.pdf>, last access: 23
515 June 2023, 2022b.
516

517 Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., Buongiorno Nardelli, B.: EU Copernicus Marine Service
518 Product User Manual for the Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed Product,
519 SST_MED_SST_L4_REP_OBSERVATIONS_010_021, Issue 2.0, Mercator Ocean International,
520 <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-021-022-041-042.pdf>, last access: 23 June
521 2023, 2022a.
522

523 Prioli, G., 2004. Shellfish farming: technologies and production. *Veterinary research communications*, 28, p.51.
524

525 Rayner, R., Jolly, C. and Gouldman, C., 2019. Ocean observing and the blue economy. *Frontiers in Marine Science*, 6, p.330.
526

527 Rodrigues, R.R., Taschetto, A.S., Sen Gupta, A. and Foltz, G.R., 2019. Common cause for severe droughts in South America
528 and marine heatwaves in the South Atlantic. *Nature Geoscience*, 12(8), pp.620-626.
529

530 Schlegel, R.W., Oliver, E.C. and Chen, K., 2021. Drivers of marine heatwaves in the Northwest Atlantic: The role of air–sea
531 interaction during onset and decline. *Frontiers in Marine Science*, 8, p.627970.
532

533 Sen Gupta, A., Thomsen, M., Benthuisen, J.A., Hobday, A.J., Oliver, E., Alexander, L.V., Burrows, M.T., Donat, M.G., Feng,
534 M., Holbrook, N.J. and Perkins-Kirkpatrick, S., 2020. Drivers and impacts of the most extreme marine heatwave events.
535 *Scientific reports*, 10(1), p.19359.
536

537 Simon A., Plecha SM, Russo A, Teles-Machado A, Donat MG, Auger P-A and Trigo RM (2022) Hot and cold marine extreme
538 events in the Mediterranean over the period 1982-2021. *Front. Mar. Sci.* 9:892201. doi: 10.3389/fmars.2022.892201

539
540 Smith, K.E., Burrows, M.T., Hobday, A.J., Sen Gupta, A., Moore, P.J., Thomsen, M., Wernberg, T. and Smale, D.A., 2021.
541 Socioeconomic impacts of marine heatwaves: Global issues and opportunities. *Science*, 374(6566), p.eabj3593.
542
543 Smith, K.E., Burrows, M.T., Hobday, A.J., King, N.G., Moore, P.J., Sen Gupta, A., Thomsen, M.S., Wernberg, T. and Smale,
544 D.A., 2023. Biological impacts of marine heatwaves. *Annual review of marine science*, 15, pp.119-145.
545
546 Vogt, L., Burger, F.A., Griffies, S.M. and Frölicher, T.L., 2022. Local drivers of marine heatwaves: a global analysis with an
547 earth system model. *Frontiers in climate*, p.49.
548
549 Wang, C., Li, Z., Wang, T. et al. Intelligent fish farm—the future of aquaculture. *Aquacult Int* 29, 2681–2711 (2021).
550 <https://doi.org/10.1007/s10499-021-00773-8>
551
552 White, C.J., Carlsen, H., Robertson, A.W., Klein, R.J.T., Lazo, J.K., Kumar, A., Vitart, F., Coughlan de Perez, E., Ray, A.J.,
553 Murray, V., Bharwani, S., MacLeod, D., James, R., Fleming, L., Morse, A.P., Eggen, B., Graham, R., Kjellström, E., Becker,
554 E., Pegion, K.V., Holbrook, N.J., McEvoy, D., Depledge, M., Perkins-Kirkpatrick, S., Brown, T.J., Street, R., Jones, L.,
555 Remenyi, T.A., Hodgson-Johnston, I., Buontempo, C., Lamb, R., Meinke, H., Arheimer, B. and Zebiak, S.E. (2017), Potential
556 applications of subseasonal-to-seasonal (S2S) predictions. *Met. Apps*, 24: 315-325. <https://doi.org/10.1002/met.1654>
557
558
559
560
561
562







563 **Code Availability**
564 Codes used to analyse data and produce figures in this study are available at <https://github.com/RJMcAdam>.

565 **Author Contributions**

566 R.M., G.B., S.M. and E.C. conceived the study. R.M. and G.B. performed the analysis and prepared the figures. R.M wrote
567 the manuscript. G.B., S.M., E. C contributed to the interpretation of the results and to the paper writing. R.M., G.B., S.M. and
568 E.C reviewed the manuscript.


569 **Competing Interests**

570 The authors no competing interests.



Deleted:


573 **Figure 1: MHW activity across the Mediterranean Sea.** (a) MHW activity defined by reprocessed satellite observations for 2022 and the
 574 three previous record years according to Simon et al, (2022). (b) Comparison between satellite observations and forecasts of 2022 MHW
 575 activity. (c) Area of Mediterranean Sea experiencing a MHW (as a percentage of total basin area). Activity is defined as the sum of the
 576 intensity over the area undergoing a MHW. Shown here are the first 8 days of forecasts initiated on Tuesdays. Forecast start dates are
 577 shown by the blue dots.





578
 579 **Figure 2: Snapshots of SST anomalies and MHW occurrence during the different stages of the 2022 MHW.** Left: reprocessed satellite
 580 observations. Right: forecasts with a lead time of 4 days. Areas in which SST is above the 90th-percentile threshold is indicated by the dark
 581 red contour. The 13th May highlights the MHW onset, the 9th July highlights the peak activity, and the 26th August highlights the (first)
 582 decay. Regions used in Figures 3 & 4 are highlighted: Ligurian Sea (magenta), Gulf of Taranto (yellow) and the Western Mediterranean
 583 (blue).

Deleted:

Deleted: Activity is defined as the daily product of area and intensity. ...





Deleted:



Deleted:

588
 589 **Figure 3: Time series of SST and MHW occurrence in summer 2022.** Orange (yellow) shading highlights MHW (MHS) occurrence in
 590 reprocessed satellite observations. Forecast start dates are shown by the blue dots. Definitions of the Ligurian Sea (a) and Gulf of Taranto
 591 (b). Note that the climatology lines (red) do not correspond to the satellite data, not to the model output (analysis and forecasts). Crosses
 592 correspond to misses (red) and false alarms (blue) in the forecast output.

593



Deleted:

Formatted: Centred

597 **Figure 4: Effect of lead time and atmospheric forcings on forecasts of MHW activity and area.** Comparison between reprocessed
 598 satellite observations and forecasts of 2022 MHW activity (a) and area (b). Each forecast time series corresponds to a different lead time
 599 (i.e. how many days in advance the forecast was made). Forecasts of MHW activity was calculated for forecasts initiated every day; the lead
 600 time from each forecast was extracted to construct the time series. Area-averaged 2m temperature anomaly (c) and wind speed (d) from the
 601 ECMWF analysis and forecasts used to force the MedFS system. All time series correspond to the Western Mediterranean Sea (Fig. 2).

603 **Table 1: Root-Mean-Square Differences of forecasts of summer 2022 MHW activity and atmospheric conditions (Fig. 4).** Values in
 604 parenthesis are RMSD values normalised by standard deviation over the summer. Differences in MHW activity and area are relative to
 605 reprocessed satellite observations, while differences in T2M anomaly and windspeed are relative to ECMWF analysis. Each column
 606 corresponds to a different lead time.

RMSD (Normalised)	Lead: 1 day	Lead: 4 days	Lead: 7 days
MHW Activity	0.16 (0.48)	0.20 (0.59)	0.28 (0.82)
MHW Area	8.88 (0.33)	11.65 (0.43)	16.50 (0.61)
T2M Anomaly	0.18 (0.21)	0.31 (0.38)	0.52 (0.62)
Wind Speed	0.22 (0.18)	0.52 (0.42)	0.94 (0.76)

Deleted: (Fig. 1)

Deleted:

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Formatted Table