

The role of air-sea heat flux for marine heatwaves in the Mediterranean Sea

Dimitra Denaxa^{1,2}, Gerasimos Korres¹, Giulia Bonino³, Simona Masina³, Maria Hatzaki²

¹ Hellenic Centre for Marine Research (HCMR), Greece

5 ² National and Kapodistrian University of Athens, Department of Geology and Geoenvironment, Greece

³ Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Italy

Correspondence to: Dimitra Denaxa (ddenaxa@hcmr.gr)

Abstract.

Recent studies have significantly contributed in understanding physical mechanisms associated with the occurrence of Marine heatwaves (MHWs). Building upon prior research, this study investigates the relative role of air-sea heat exchange and oceanic processes during the onset and decline phases of surface MHWs in the Mediterranean Sea, based on a joint analysis of remote sensing data and reanalysis outputs over the period 1993-2022. Results show that air-sea heat flux is the major driver in 44% of the onset and only 17% of the decline MHW phases. Thus, these findings suggest that oceanic processes play a key role in driving SST anomalies during MHWs, particularly during declines. The role of surface flux becomes more important during warmer months and onset periods. Spatially, the heat flux contribution is greater in the Adriatic and Aegean sub-basins, where it becomes the major driver of most onset phases. Latent heat emerges as the most significant heat flux component in forming the SST evolution, across all seasons. Onset/decline phases lasting less than 5 days experience a weaker contribution of heat flux, compared to longer phases (lasting 5-10 or more than 10 days). Moreover, an inverse relationship between MHW severity and the contribution of heat flux is observed. At the subsurface, mixed layer shoaling is found over the entire duration of most MHWs, particularly for those of shorter duration. Therefore, the surface cooling right after the peak day is likely not associated with vertical mixing in such cases. These findings suggest that other oceanic processes, potentially horizontal advection, have a key role in modulating SST at the beginning of most MHW declines. In turn, further dissipation of heat is commonly driven by vertical mixing, as indicated by a significant mixed layer deepening after the MHW end day in most cases. This study emphasizes the need for considering subsurface information for future studies of MHWs and highlights the importance of accounting for limitations associated with the definitions employed for MHW phases.

1. Introduction

Marine heatwaves (MHWs) are extreme events, characterized by prolonged periods of anomalously high water temperature, lasting for at least five consecutive days (Hobday et al., 2016). These events have gathered increased attention due to their detrimental effects on marine life, especially given the increase observed in their frequency, intensity and duration over the recent decades, at global and Mediterranean scale (Oliver et al., 2018, Holbrook et al., 2019; Darmaraki et al., 2019a; Juza et al., 2022; Dayan et al.,

2023, Pastor and Khodayar, 2023). Mass mortality events and local extinctions, coral bleaching and massive shifts of marine species have been extensively reported (Wernberg et al., 2016; Frölicher and Laufkotter, 2018; Smale et al., 2019; Garrabou et al., 2022; Smith et al., 2023), along with socioeconomic impacts on fishery and aquaculture industries (Mills K.E., et al., 2013; Cavole L., 2016). The intensification of MHW conditions has been attributed mostly to ocean warming (Oliver et al., 2018, Ciappa, 2022), while further intensification is expected in the future (Oliver et al., 2019; Darmaraki et al., 2019b; Plecha & Soares, 2019; Hayashida et al., 2020), driven by anthropogenic forcing and particularly pronounced under high-emission future scenarios (Oliver et al., 2019).

Given these concerns, it is crucial to enhance our understanding of the driving factors behind MHWs at regional scale, particularly within the framework of exploring predictability options and facilitating marine decision-making (Holbrook et al., 2020; Spillman et al., 2021). Recent research has significantly contributed in identifying physical drivers and MHW-favoring conditions (e.g., Holbrook et al., 2019; Sen Gupta et al., 2020; Oliver et al., 2021; Vogt et al., 2022; Marin et al., 2022). Individual events in the Mediterranean Sea have also been explored, such as the widely known MHW in 2003 in the western Mediterranean Sea (e.g., Sparnocchia et al., 2006; Olita et al. 2007; Bonino et al., 2023), the short-lasting, record-breaking MHW in May 2020 in the southeastern Mediterranean Sea (Ibrahim et al., 2021; Denaxa et al., 2022) and the most recent long-lasting MHW in summer 2022 (Marullo et al., (2023); McAdam et al., 2024; Pirro et al., 2024).

However, a limited number of studies have assessed physical drivers separately for the buildup and decay of MHW events, employing different methodologies and datasets. Schlegel et al. (2021) demonstrated that nearly 50% of surface MHWs in the northwest Atlantic are heat flux-driven but less than 20% decay due to heat flux, suggesting that oceanic processes are mainly responsible for the MHW decline. Marin et al. (2022) investigated upper-ocean MHWs based on global ocean circulation model output and found that heat advection, followed by anomalous air-sea heat flux, explain most of the upper ocean temperature anomalies during both MHW onset and decline phases. Within this context, the present study utilizes high resolution observational Sea Surface Temperature (SST) and modelled heat flux data to assess the driving role of air-sea heat exchange during onset and decline phases of surface MHWs in the Mediterranean Sea. Furthermore, it provides insights into the concurrent subsurface conditions by examining the mixed layer dynamics during MHWs.

2. Data and Methods

MHWs in this study were identified based on high resolution gridded satellite SST data in the Mediterranean Sea. Daily SST values from the Reprocessed and near-real-time datasets from the Copernicus Marine (products ref. no. 01 and 02 (Table 1), $0.05^\circ \times 0.05^\circ$ and $0.01^\circ \times 0.01^\circ$ horizontal resolution, respectively) were used to cover the period Jan 1993 - Dec 2022. To study the net surface heat budget (Q_{net}), turbulent and radiative surface flux were obtained from the ECMWF ERA5 Reanalysis dataset at hourly frequency and $0.25^\circ \times 0.25^\circ$ horizontal resolution (product ref. no. 03 (Table 1)). Finally, daily values of mixed layer depth (MLD) for the period Jan 1993 - Dec 2022 were obtained from the Mediterranean Sea Physics Reanalysis and the Mediterranean Sea Physics

Analysis and Forecast (products ref. no. 04 and no. 05 (Table 1), 0.042 ° x 0.042 ° horizontal resolution). Collocation of SST and MLD with the coarser ERA5 data was performed by using the nearest neighboring value to each ERA5 grid point. The paired values at daily frequency were used in the heat budget analysis.

MHW detection was performed based on the definition and detection methodology of Hobday et al. (2016), using the matlab toolbox provided by Zhao and Marin (2019). The reference period used in this study to create the daily climatology required for the event detection is the same as the 30-year study period (1993-2022). To compute the climatology, a time-window of 11 days was employed, centered on the day when each daily climatological value was computed. For the computation of the daily threshold time-series, the 90th percentile was selected, being widely used in MHW studies, thus allowing for a more direct inter-comparison with literature. Additionally, a 30 day-window was applied for smoothing the threshold time-series.

Basic properties were computed for each identified event (e.g., start/end day, mean/max intensity, duration). Next, events were split in their onset and decline phase. The onset phase was considered to last from the 1st day until the day of maximum intensity (I_{\max}), and the decline phase from I_{\max} -day until the last day of the event. As in Schlegel et al. (2021), for each phase, an ocean mixed layer heat budget analysis was applied to derive the change in SST attributed to Q_{net} , based on the following equation:

$$SST'_{t_2} - SST'_{t_1} = \int_{t_1}^{t_2} \frac{Q'}{\rho_o c_p h} dt + R \quad (1)$$

The left-hand side of Eq.1 represents the observed change in SST anomaly ($DSST'_{\text{obs}}$) relative to climatology, during a specific phase. Each phase starts at day t_1 and ends at day t_2 , being the start day and I_{\max} -day for onset phases, or I_{\max} -day and end day for decline phases, respectively. On the right-hand side of Eq.1, Q' is the daily anomaly of Q_{net} . The latter consists of the latent and sensible heat flux and net short-wave and net long-wave radiation (LH, SH, SWR and LWR, respectively), as follows:

$$Q_{\text{net}} = LH + SH + SWR + LWR \quad (2)$$

It should be noted that a simplified approach was used that does not account for the penetration of solar radiation below the mixed layer. Considering the Jerlov Water Type IA for relatively clear sea water, 77% of the solar radiation is expected to be absorbed within the upper 10 m of the ocean, based on the solar radiation attenuation equation in Paulson and Simpson (1977). Taking into account the MLD values obtained from the utilized reanalysis dataset, we have considered that the followed approach does not significantly affect our conclusions.

To compute heat flux anomalies, first a daily climatology was computed following the same methodology as with SST for the period 1993-2022. Daily heat flux anomalies were then constructed relative to the mean climatological value over the phase duration (e.g.,

as in Fewings and Brown, 2019). The time integral of Q' divided by the product of the constant values ρ (seawater density), C_p (specific heat capacity) and h (mixed layer thickness) represents the part of the $DSST'_{obs}$ explained by Q' during this phase ($DSST_Q$). The second term of the right-hand side in Eq.1 stands for any contribution to $DSST'_{obs}$ from other mechanisms affecting SST: horizontal advection, vertical mixing processes, horizontal diffusion of heat flux, radiative heat loss below the mixed layer.

100 Therefore, the role of air-sea heat flux in this study is assessed in relation to a single residual term (i.e., the non-heat flux terms merged into a single one), representing the cumulative effect of all other (oceanic) factors influencing the SST tendency during a MHW phase.

The contribution of Q' in driving a MHW onset/decline phase N is then quantified through the following proportion of change:

$$P(N) = \frac{DSST_Q(N)}{DSST'_{obs}(N)} \quad (3)$$

Therefore, a positive heat flux contribution value during a MHW phase indicates a favoring role of heat flux in the corresponding SST evolution, i.e., a warming (cooling) effect of heat flux during onset (decline). Analogously, a negative contribution value during either an onset or a decline phase, indicates that heat flux opposes the corresponding SST tendency. Finally, to examine the evolution of MLD during MHWs, time-series of cumulative MLD anomalies (MLDA) were constructed for onset/decline phases. These time-series were computed by adding, at each MHW day, the daily anomaly of MLD of the previous day, in order to account for longer time-scales associated with the mixed layer evolution (as in Schlegel et al. (2021)). To explore the correlation between MLDA and SST anomalies (SSTA), Pearson correlation coefficients were computed for onset and decline phases, separately.

Table 1

Product ref. no	Product ID & type	Data Access	Documentation
1	SST_MED_SST_L4_REP_OBSERVATIONS_010_021; Satellite observations	EU Copernicus Marine Service Product (2022a)	Product User Manual (PUM): Pisano et al. (2022a) Quality Information Document (QUID): Pisano et al. (2022b)
2	SST_MED_SST_L4_NRT_OBSERVATIONS_010_004; Satellite observations	EU Copernicus Marine Service Product (2022b)	PUM: Pisano et al. (2022c) QUID: Pisano et al. (2022d)

3	ERA5 hourly data on single levels; Numerical models	Copernicus Climate Data Store	Hersbach et al. (2023) https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview
4	MED_MULTIYEAR_PHY S_006_004; Numerical models	EU Copernicus Marine Service Product (2022c)	PUM: Lecci et al. (2022a) QUID: Escudier et al. (2022)
5	MEDSEA_ANALYSISFO RECAST_PHY_006_013; Numerical models	EU Copernicus Marine Service Product (2022d)	PUM: Lecci et al. (2022b) QUID: Goglio et al. (2022)

3. Results

3.1 MHW detection

Properties of MHWs exhibit high variability throughout the Mediterranean Sea (Fig. 1a-c). The northwestern part of the basin, along with the northern Adriatic and northern Aegean Seas, experienced on average the highest event intensity over the period 1993-2022, exceeding 2.5 and 2 °C, respectively (Fig. 1b). Events tend to last longer in the eastern part of the basin (Aegean and Levantine Seas) and in the central-western Mediterranean Sea, while the shortest durations are mostly found in the Ionian and Alboran Seas (Fig. 1c). The mean event frequency over the study period shows some similarities with the mean intensity spatial distribution, suggesting that the most (least) intense and most (least) frequent MHWs are encountered in the northernmost (southernmost) flanks of the Mediterranean Sea (Fig. 1a,b). Results generally agree with literature on MHWs in the Mediterranean Sea during the recent decades (Darmaraki et al., 2019a; Ibrahim et al., 2021; Juza et al., 2022; Dayan et al., 2023), despite some differences in event detection methods (e.g., choice for percentile-based threshold, accounting for MHW spatial extent) or study periods (choice of climatological period, period for event detection).

MHW frequency has been increasing over the past 30 years, with a trend of 1 ± 0.6 events per decade for the entire basin (Fig. 1d). Also, MHW duration has increased particularly in the eastern basin, with trend values locally exceeding ten days per decade (1.7 ± 1.5 days per decade for the entire basin) (Fig. 1f). Notably, intensity has not increased over the entire basin during the study period (0.08 ± 0.2 °C per decade for the basin, not significant at the 95% confidence level) (Fig. 1e). The northernmost regions, which are characterized by higher MHW intensity, present small decreasing MHW intensity trends (though not statistically significant in most cases) which is in agreement with Dayan et al. (2023), and Ibrahim et al. (2021) for the eastern basin.




Figure 1 Left: Mean annual frequency, intensity and duration of MHWs for the period 1993-2022 (a, b, c, respectively); Right: Linear trends of annual values of frequency, intensity and duration (d, e, f, respectively). Black dots superimposed on trend fields correspond to statistically non-significant trends (Mann Kendall test, 95% confidence level)

3.2 The role of heat flux during MHW onset and decline

140 Surface heat flux contributes to the observed surface warming during the onset phase of the majority of the events (92%) detected in the Mediterranean Sea within 1993-2022 (Fig. 2a). In particular, in 44% of the events, heat flux presents a major role during the development of MHWs, in terms of explaining more than half of the observed change in SST anomaly (Fig. 2b). During the rest of the events (8%), heat flux opposes the surface warming over the onset phases; therefore, other mechanisms compensate for their cooling effect and are responsible for the observed warming in such cases.




Figure 2 a) Contribution of heat flux to the observed change in SSTA during MHW onset and decline phases, for the period 1993-2022; b) Percentage of events primarily driven by heat flux (i.e., with more than half of the observed warming (cooling) attributed to heat flux during onset (decline)) for the entire Mediterranean Sea and sub-regions mapped in (c); d-e) Boxplots for the contribution of heat flux during onset (red) and decline (blue) for the entire Mediterranean Sea and the sub-regions; f-g) Boxplots for heat flux contribution per season, for onset (red) and decline (blue); h-i) Boxplots per season, for the contribution of heat flux components during onset (red) and decline (blue); From left to right: Latent heat flux, sensible heat flux, net short-wave radiation, net long-wave radiation. Notes: Positive heat-flux contribution during onset (decline) means that heat-flux warm (cool) the sea surface; boxes in boxplots define the interquartile range from the 25th up to the 75th percentile and whisker bars correspond to values falling within 1.5 times the interquartile range.

Air-sea heat exchange contributes to surface cooling during decline phases in a much smaller percentage of MHWs (58%), while it is the primary driver only in 17% of the decline phases (Fig. 2a,b). In other words, almost half (42%) of the observed MHWs in the basin decay under non-favorable heat flux conditions (i.e., heat flux opposing the SST decrease), while oceanic processes are the dominant driver of most (83%) MHW declines.

150

Heat flux exchange explains to a great extent the MHW onset in all examined sub-regions (Fig. 2d). The Adriatic and the Aegean Seas stand out with most of the observed heating during onset attributed to heat flux in more than half of the identified events (Fig. 2b). In 6 out of the 7 sub-regions, the percentage of events primarily driven by heat flux during their onset ranges between 39%-53%. The Alboran Sea presents a much lower percentage of events attributed to heat flux (22%), potentially suggesting an enhanced
155 role of advective processes in regulating SST in the area, likely associated with its proximity to the Gibraltar Strait. In contrast, the corresponding percentages for the decline phases show less spatial variability across all sub-regions, with only 15-20% of the examined declines found to be mainly driven by heat flux exchange (Fig. 2b).

160

Seasonal analysis was performed considering winter as the period from December to February, while events spanning different seasons were assigned to the season when their intensity maximizes. Most of the events were found to occur during summer, followed by spring, autumn and then winter. As expected, the magnitude of SST anomalies during MHWs varies among the seasons (e.g., Thorat et al., 2022), but the contribution of heat flux to their formation is consistently weaker during decline compared to onset phases throughout the year (Fig. 2f,g).

165

Results suggest a greater contribution of heat flux exchange to the MHW evolution within warmer seasons (from an ocean perspective, i.e., summer and autumn) (Fig. 2f,g). Autumn shows the highest percentage of events driven by Q_{net} in both onset and decline. LH flux is mainly regulating the contribution of Q_{net} during both phases and throughout the year, followed by SWR and then SH, while LWR exhibits less clear behavior in all seasons (Fig. 2h,i). With the exception of SH, the relative contribution of each heat flux component presents higher variability during decline compared to onset phases, across all seasons. Specifically,
170 approximately an equal number of decline cases are associated with positive and negative contributions of the heat flux components to the observed cooling, indicating a less predictable role of Q_{net} during decline compared to onset periods throughout the year. (Fig. 2i). The lowest percentage of heat flux-driven events is observed during the winter onset and spring decline phases. Particularly for spring, heat flux during most declines act against the observed surface cooling (mainly through suppressed LH losses) suggesting that MHW dissipation in spring is commonly driven by oceanic factors.

175

Results show that the air-sea interaction, with a dominant role of LH flux, plays a major role in the development of nearly half (44%) of the MHWs in the Mediterranean Sea. This finding suggests that oceanic processes play a key role during 56% of the onset cases. A further weakened role of heat flux is found during decline periods (being the major contributor in only 17% of declines), indicating that MHWs decay is also primarily driven by oceanic processes.

180 3.3 Links with mixed layer depth and MHW characteristics

To gain insight into subsurface conditions during MHWs, we examined how MLD evolves in relation to SST. Negative (positive) correlation between MLDA and SSTA that was found for a large number of events during onset (decline) implies mixed layer shoaling over the entire MHW duration in these cases (Fig. 3a-top). While a MHW event develops, a reduction in MLD is commonly expected, as surface warming may strengthen the stratification of the water column (D’Ortenzio & Prieur, 2012). Given that heat
185 flux is found to contribute to the warming phases in most cases, the concurrent mixed layer shoaling found during most onsets may be interpreted as an effect of the warming driven by the atmosphere. Nevertheless, a thinner than usual mixed layer may also exist before the event occurrence and act as a pre-conditioning factor (e.g., Lee et al., 2023). The greatest MLDA-SSTA correlation is found during spring and summer events, while no significant correlation is observed in winter (not shown), as the deeper mixed layer during colder months is expected to be less responsive to surface SST variations (D’Ortenzio & Prieur, 2012). High positive
190 MLDA-SSTA correlation observed during declines implies that the mixed layer in such cases continues to shoal while SST decreases (Fig. 3a), similar to the mixed layer analysis by Schlegel et al. (2021) for the northwest Atlantic. This finding further supports that oceanic processes play a significant role during MHW decline periods, as also indicated by the weaker contribution of heat flux during declines. It also suggests that the surface cooling occurring right after the peak intensity day is likely not due to mixing in the vertical (further discussed below).

195

To gain a better understanding of the relationship between subsurface conditions and the contribution of heat flux exchange at the air-sea interface during the evolution of MHWs, this information is integrated into Fig. 3a. Whereas during onset, MHWs are largely driven by heat flux exchange and most of them are accompanied by mixed layer shoaling, there are onset cases where MLDA are strongly positively correlated with SSTA, indicating surface warming evolves while the mixed layer deepens (Fig. 3a-top).
200 Regarding decline periods, almost equal numbers of cases show positive and negative heat flux contributions (Fig. 3a-bottom). While a decrease in MLD is evident in most decline phases (as indicated by correlation coefficient close to one), a significant MLDA-SSTA correlation is absent in a considerable number of decline phases. Additionally, there are cases during declines where a high negative correlation between MLDA and SSTA is observed, indicating mixed layer deepening while SST decreases. Such cases are encountered when heat flux contributes to surface cooling (Fig. 3a-bottom), suggesting that vertical mixing (most probably
205 wind-induced), works in the same direction.

MHW onset and decline phases were also examined in relation to their duration. They were grouped in short, medium and long duration, considering phases lasting less than 5 days, from 5 to 10 days and more than 10 days, respectively (Fig. 4a). Shorter onset/decline periods, being the most prevalent category, slightly overshadow the contribution of heat flux during longer lasting
210 MHW phases (Fig. 4a-c vs Fig. 2a). In particular, as compared to medium, followed by long durations, short durations (both for





Figure 3 a) Heatmap relating the net heat-flux contribution with Pearson correlation coefficient (CC) between time-series of SST anomalies and time-series of mixed layer depth cumulative anomalies computed for each MHW onset and decline phase (upper and lower heatmap, respectively) for the period 1993-2022. Colors correspond to the number of events falling in each bin. b) Boxplots for the distribution of CC values for onset/decline phases lasting less than 5 days, between 5-10 days and more than 10 days (short, medium and long duration phases, respectively); c) Percentage change (%) of MLD over a fixed 7-days period after the MHW end day compared to MLD during the onset period. Notes: Positive heat-flux contribution during onset (decline) means that heat-flux warm (cool) the sea surface; Negative (positive) CC values close to -1 (+1) during onset (decline) correspond to reduction of the mixed layer depth while SST increases (decreases); Boxes in boxplots define the interquartile range from the 25th up to the 75th percentile and whisker bars correspond to values falling within 1.5 times the interquartile range.

onset and decline) tend to exhibit a smaller contribution of heat flux exchanges triggering the SST evolution (Fig. 4a-c). For long-

duration onset phases, the contribution of heat flux surpasses the contribution of oceanic factors, as the former explain more than half of the surface warming, in more than half of the events (Fig. 4c). Similarly, it is mostly during shorter declines that heat flux does not present a systematically positive or negative contribution to the SST decrease, while a shift towards higher contributions of heat flux is found for longer phases (Fig. 4a-c). Importantly, the relationship between MLDA and SSTA becomes less clear during onset/decline phases of medium and, in turn, long duration, as indicated by weaker correlation (Fig. 3b). In the case of decline periods, mixed layer shoaling is more frequently observed during short-lasting declines, suggesting a continuation of mixed layer shoaling already present during the onset in such cases.

These findings suggest that the definition followed for a MHW evolution phase needs to be cautiously taken into account when interpreting results, especially when surface and sub-surface conditions are examined for the same phases. This is due to the time needed for a surface warming signal to penetrate below as well as due to the longer time-scales associated with processes at deeper layers. Notably, at the end day of a surface MHW, being the end day of the decline phase (as defined in this study), SST is still above the event-detection threshold and will fall below this value by the following day. Therefore, events further dissipate after the decline phase ends. Given such considerations, and depending on the specific purpose of a study, different approaches for defining a decline period may be followed. For instance, Darmaraki (2019c) considered the entire MHW duration as the event development phase and treated the following period as decline, Schlegel et al. (2021) used the same definitions as the present study, while Marin et al. (2022) considered a dynamic criterion for defining the decay period, including a larger portion of the temperature change during the event dissipation after the MHW end day.

On these grounds, we additionally examine how MLD evolves after the MHW end day, to shed light on what follows the progressively increasing stratification we observed during MHWs. To this aim, we compute for each event the percentage change in the mean MLD between the onset period and a fixed 7 days-period following the MHW end day (Fig. 3c). Results reveal the existence of a deeper mixed layer after the decline of most events (83.5%) compared to the onset period. In these cases, the magnitude of change in MLD is also significantly greater, occasionally exceeding 100%, compared to instances when the mixed layer is thinner during this period than during the onset (changing the predetermined length of the examined post-decline period does not significantly alter these conclusions). This analysis shows that a significant mixed layer deepening most likely occurs after the end of MHW decline phases in the Mediterranean Sea, suggesting that vertical mixing eventually contributes to the heat dissipation.

Considering the continuation of mixed layer shoaling found mostly during shorter declines, results suggest that the oceanic factor regulating the SST decrease after the peak intensity day of most events is probably heat advection, while vertical mixing in these cases becomes important over the following days. In line with our findings, Marin et al. (2022) found that the principal driver of the upper ocean temperature changes during MHW onset and decline phases in mid latitudes is horizontal heat advection. However,

245 their results cannot be directly inter-compared to ours, as they use modelled depth-integrated upper ocean temperatures (thus not detecting surface events), they focus only on extreme cases and they employ a different definition of onset/decay period.

Finally, results were examined in relation to MHW intensity and severity, to account for the perspective of MHW extremity in terms of absolute and normalized SST anomalies, respectively. Events were split based on their severity category following the categorization scheme proposed by Hobday et al., 2018 (definitions included therein). Results suggest a smaller contribution of heat flux in the evolution of more severe events (Fig. 4d-f). Half of the moderate events were found to be mainly driven by heat flux, while this percentage decreases for strong, severe and extreme events, as illustrated in Fig. 4d-f. This may be associated with the enhanced role of heat flux during longer events, as was previously discussed, and the fact that events of higher severity categories tend to present shorter duration. Similar results were found when examining the contribution of air-sea heat flux in relation to the mean intensity of MHWs (not included). Nevertheless, further investigations are needed to unravel how our methodological choices affect these findings.

250

255




Figure 4 Left: Contribution of heat flux to the observed change in SSTA during MHW onset and decline phases, for the period 1993-2022, for phases lasting: less than 5 days (a), between 5-10 days (b) and more than 10 days (c) (short, medium and long duration phases, respectively). Right: Same as left but for different MHW severity categories: moderate (d), strong (e) and severe/extreme (f), based on the categorization scheme by Hobday et al. (2018).

4. Conclusions

This study investigates the role of air-sea heat exchange during MHWs in the Mediterranean Sea, using satellite and reanalysis data within 1993-2022. An ocean mixed layer heat budget analysis is performed to derive the change in SST attributed to the net surface
260 heat budget during onset and decline phases.

Air-sea heat flux is the primary driver in 44% of the onset and only in 17% of the decline phases in the basin. Heat flux acts in favor of the development of most MHWs, across all seasons and especially during warmer months and onset phases. Moreover, its contribution is greater in the Adriatic and Aegean sub-basins, where it becomes the major driver of most onset phases. Among the
265 heat flux components, LH emerges as the most significant contributor to SST anomalies, in line with prior studies (Sen Gupta et al., 2020; Oliver et al. 2021; Schlegel et al., 2021; Marin et al., 2022). Short-lasting onset/decline phases (shorter than 5 days) tend to experience a smaller contribution of heat flux in forming the SST evolution, compared to longer phases (lasting 5-10 or more than 10 days). Furthermore, there is an inverse relationship between MHW severity and the contribution of heat flux. As less than half of the events are primarily driven by air-sea heat flux, our results imply a key role of oceanic processes in regulating SST during
270 most events and particularly during decline periods.

Examining the mixed layer during MHWs revealed a progressively decreasing MLD over the entire event-duration, particularly for shorter-lasting events. In turn, a significant mixed layer deepening was found to occur after the end of the decline period of most events in the basin. In cases of smaller contribution of heat flux (e.g. during shorter compared to longer events, or during declines
275 compared to onsets), a stronger correlation between SSTA and MLDA is also found, further supporting the key-role of oceanic processes in such cases. These findings suggest that the surface cooling occurring right after the peak intensity day is likely not associated with vertical mixing. Moreover, this potentially suggests that horizontal advection is the oceanic factor playing the most significant role by the time the decline period begins, especially for shorter declines. This hypothesis finds support in the results of Marin et al. (2022), who highlighted the role of horizontal heat advection during the MHW evolution, and Schlegel et al. (2021)
280 who suggested that advection and mixing should drive the MHW decline, based on similar indications.

Nevertheless, authors suggest taking into account potential limitations associated with the definition of MHW phases followed within a study, especially while examining concurrent sub-surface conditions. These considerations concern the longer time-scales below the sea surface, as well as the complexity associated with long-lasting events which are not expected to be adequately
285 described by two single phases. Specifically, subsequent warming and cooling periods may occur within a long onset or decline phase, complicating their representation by the definition employed in this study for onset and decline. These concerns are part of the broader discussion on challenges related to the lack of a standardized framework for analyzing MHW drivers, as many other methodological approaches (e.g., the choice of integration depths in heat budget analyses), may potentially influence findings on MHW drivers. For this reason, alongside recommending the use of definitions and methods aligned with the specific contexts of

290 individual studies, we highlight that clearly articulating the employed methods within a study is vital both for the precise
interpretation of results and for meaningful comparisons across different studies on MHW drivers. Despite the aforementioned
caveats, this study provides useful insights into the role of surface heat flux and mixed layer dynamics during MHWs in the
Mediterranean Sea. Importantly, following the methodology applied by Schlegel et al. (2021) allowed for a fair inter-comparison of
295 is largely attributed to the same methodology employed to assess drivers of SST-defined events, and especially to the same definition
followed for onset and decline periods. The striking similarities in the results for the two regions may also imply inherent
characteristics of the interplay of air-sea interaction and oceanic processes during anomalous SST fluctuations over similar
latitudinal zones, which is observed to a certain extent also in Marin et al. (2022).

300 This study highlights the need for considering subsurface information in MHW studies, to gain insight into ocean internal dynamics
throughout their evolution. In this context, combining observations and ocean reanalysis systems, such as in Dayan et al. (2023),
and considering MHW evolution periods aligned with the objectives and specific characteristics of a study appears to be a promising
direction towards understanding physical drivers, improving monitoring, and therefore enabling early warning of MHWs.

Data availability

305 Information on the products used in this paper is included in Table 1.

Author contribution

DD defined the research problem. DD conducted the analysis and wrote the manuscript, with contributions from GK and MH. All
authors contributed to the interpretation of results.

Competing interests

310 The contact author has declared that none of the authors has any competing interests.

References

Bonino, G., Masina, S., Galimberti, G., and Moretti, M.: Southern Europe and western Asian marine heatwaves (SEWA-MHWs): a
dataset based on macroevents, *Earth Syst. Sci. Data*, 15, 1269–1285, <https://doi.org/10.5194/essd-15-1269-2023>, 2023.

Cavole, L. M., Demko, A. M., Diner, R. E., Giddings, A., Koester, I., Pagniello, C. M. L. S., Paulsen, M. L., Ramirez-Valdez, A.,
315 Schwenck, S. M., Yen, N. K., Zill, M. E., and Franks, P. J. S.: Biological impacts of the 2013–2015 warm-water anomaly in the
northeast Pacific: Winners, Losers, and the Future, *Oceanography*, 29, 273–285, <https://doi.org/10.5670/oceanog.2016.32>, 2016.

- Ciappa, A. C.: Effects of Marine Heatwaves (MHW) and Cold Spells (MCS) on the surface warming of the Mediterranean Sea from 1989 to 2018, *Prog. Oceanogr.*, 205, 102828, <https://doi.org/10.1016/j.pocean.2022.102828>, 2022.
- D’Ortenzio, F. and Prieur, L.: The upper mixed layer, *Life Mediterr. Sea A Look Habitat Chang.*, 127–156, 2012.
- 320 Darmaraki, S., Somot, S., Sevault, F., and Nabat, P.: Past Variability of Mediterranean Sea Marine Heatwaves, *Geophys. Res. Lett.*, 46, 9813–9823, <https://doi.org/10.1029/2019GL082933>, 2019a.
- Darmaraki, S., Somot, S., Sevault, F., Nabat, P., Cabos Narvaez, W. D., Cavicchia, L., Djurdjevic, V., Li, L., Sannino, G., and Sein, D. V.: Future evolution of Marine Heatwaves in the Mediterranean Sea, *Clim. Dyn.*, 53, 1371–1392, <https://doi.org/10.1007/s00382-019-04661-z>, 2019b.
- 325 Darmaraki S., Mediterranean marine heatwaves: detection, past variability and future evolution, Ph.D. thesis, Université Paul Sabatier - Toulouse III, ⟨NNT : 2019TOU30072⟩, 2019c.
- Dayan, H., McAdam, R., Juza, M., Masina, S., and Speich, S.: Marine heat waves in the Mediterranean Sea: An assessment from the surface to the subsurface to meet national needs, *Front. Mar. Sci.*, 10, 1–21, <https://doi.org/10.3389/fmars.2023.1045138>, 2023.
- Denaxa, D., Korres, G., Sotiropoulou, M., and Perivoliotis L.: Extreme Marine Heatwave in the eastern Mediterranean in May 2020.
- 330 In: Copernicus Ocean State Report, Issue 6, *Journal of Operational Oceanography*, 15:sup1, s119–s126. <https://doi.org/10.1080/1755876X.2022.2095169>, 2022.
- Escudier, R., Clementi, E., Nigam, T., Aydogdu, A., Fini, E., Pistoia, J., Grandi, A., and Miraglio, P.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea Physics Reanalysis Product, MEDSEA_MULTIYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International, [https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-](https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-004.pdf)
- 335 [004.pdf](https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-004.pdf), last access: 13 July 2023, 2022.
- EU Copernicus Marine Service Product: Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed, Mercator Ocean International [data set], <https://doi.org/10.48670/moi-00173>, 2022a.
- EU Copernicus Marine Service Product: Mediterranean Sea High Resolution and Ultra High Resolution Sea Surface Temperature Analysis, Mercator Ocean International [data set], <https://doi.org/10.48670/moi-00172>, 2022b.
- 340 EU Copernicus Marine Service Product: Mediterranean Sea Physics Reanalysis, Mercator Ocean International [data set], https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_PHY_006_004_E3R1, 2022c.
- EU Copernicus Marine Service Product: Mediterranean Sea Physics Analysis and Forecast, Mercator Ocean International [data set], https://doi.org/10.25423/CMCC/MEDSEA_ANALYSISFORECAST_PHY_006_013_EAS7, 2022d.
- Fewings, M. R. and Brown, K. S.: Regional Structure in the Marine Heat Wave of Summer 2015 Off the Western United States,
- 345 *Front. Mar. Sci.*, 6, 1–14, <https://doi.org/10.3389/fmars.2019.00564>, 2019.
- Frölicher, T. L. and Laufkötter, C.: Emerging risks from marine heat waves, *Nat. Commun.*, 9, 2015–2018, <https://doi.org/10.1038/s41467-018-03163-6>, 2018.
- Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Ponti, M., Schlegel, R., Bensoussan, N., Turicchia, E., Sini, M., Gerovasileiou, V., Teixido, N., Mirasole, A., Tamburello, L., Cebrian, E., Rilov, G., Ledoux, J. B., Souissi, J. Ben, Khamassi, F.,
- 350 Ghanem, R., Benabdi, M., Grimes, S., Ocaña, O., Bazairi, H., Hereu, B., Linares, C., Kersting, D. K., la Rovira, G., Ortega, J.,

- Casals, D., Pagès-Escolà, M., Margarit, N., Capdevila, P., Verdura, J., Ramos, A., Izquierdo, A., Barbera, C., Rubio-Portillo, E., Anton, I., López-Sendino, P., Díaz, D., Vázquez-Luis, M., Duarte, C., Marbà, N., Aspillaga, E., Espinosa, F., Grech, D., Guala, I., Azzurro, E., Farina, S., Cristina Gambi, M., Chimienti, G., Montefalcone, M., Azzola, A., Mantas, T. P., Frascchetti, S., Ceccherelli, G., Kipson, S., Bakran-Petricioli, T., Petricioli, D., Jimenez, C., Katsanevakis, S., Kizilkaya, I. T., Kizilkaya, Z., Sartoretto, S.,
355 Elodie, R., Ruitton, S., Comeau, S., Gattuso, J. P., and Harmelin, J. G.: Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea, *Glob. Chang. Biol.*, 28, 5708–5725, <https://doi.org/10.1111/gcb.16301>, 2022.
- Goglio, A.C., Clementi, E., Grandi, A., Mariani, A., Giurato, M., and Aydogdu, A.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea Physics Analysis and Forecast Product, MEDSEA_ANALYSISFORECAST_PHY_006_013, Issue 2.3, Mercator Ocean International,
360 <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MED-QUID-006-013.pdf>, last access: 13 July 2023, 2022.
- Hayashida, H., Matear, R. J., Strutton, P. G., and Zhang, X.: Insights into projected changes in marine heatwaves from a high-resolution ocean circulation model, *Nat. Commun.*, 11, 1–9, <https://doi.org/10.1038/s41467-020-18241-x>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J-N.: ERA5 hourly data on single levels from 1940 to present.
365 Copernicus Climate Change Service (C3S) Climate Data Store (CDS), <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C. J., Benthuisen, J. A., Burrows, M. T., Donat, M. G., Feng, M., and Holbrook, N. J.: A hierarchical approach to defining marine heatwaves, *Prog. Oceanogr.*, 141, 227–238, 2016.
- Hobday, A. J., Oliver, E. C. J., Sen Gupta, A., Benthuisen, J. A., Burrows, M. T., Donat, M. G., Holbrook, N. J., Moore, P. J., Thomsen, M. S., Wernberg, T., and Smale, D. A.: Categorizing and naming marine heatwaves, *Oceanography*, 31, 162–173, 2018.
370 Holbrook, N. J., Scannell, H. A., Sen Gupta, A., Benthuisen, J. A., Feng, M., Oliver, E. C. J., Alexander, L. V., Burrows, M. T., Donat, M. G., Hobday, A. J., Moore, P. J., Perkins-Kirkpatrick, S. E., Smale, D. A., Straub, S. C., and Wernberg, T.: A global assessment of marine heatwaves and their drivers, *Nat. Commun.*, 10, 1–13, <https://doi.org/10.1038/s41467-019-10206-z>, 2019.
- Holbrook, N. J., Sen Gupta, A., Oliver, E. C. J., Hobday, A. J., Benthuisen, J. A., Scannell, H. A., Smale, D. A., and Wernberg, T.: Keeping pace with marine heatwaves, *Nat. Rev. Earth Environ.*, 1, 482–493, <https://doi.org/10.1038/s43017-020-0068-4>, 2020.
- 375 Ibrahim, O., Mohamed, B., and Nagy, H.: Spatial Variability and Trends of Marine Heat Waves in the Eastern Mediterranean Sea over 39 Years, *J. Mar. Sci. Eng.*, 9, 643, <https://doi.org/10.3390/jmse9060643>, 2021.
- Juza, M., Fernández-Mora, A., and Tintoré, J.: Sub-Regional Marine Heat Waves in the Mediterranean Sea From Observations: Long-Term Surface Changes, Sub-Surface and Coastal Responses, *Front. Mar. Sci.*, 9, <https://doi.org/10.3389/fmars.2022.785771>, 2022.
- Lecci, R., Drudi, M., Grandi, A., Cretì, S., and Clementi, E.: EU Copernicus Marine Service Product User Manual for the
380 Mediterranean Sea Physics Reanalysis Product, MEDSEA_MULTIYEAR_PHY_006_004, Issue 2.3, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-004.pdf>, last access: 13 July 2023, 2022a.
- Lecci, R., Drudi, M., Grandi, A., Cretì, S., and Clementi, E.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea Physics Analysis and Forecast Product, MEDSEA_ANALYSISFORECAST_PHY_006_013, Issue 2.2,

- 385 Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MED-PUM-006-013.pdf>, last access: 13 July 2023, 2022b.
- Lee, E. Y., Lee, D. E., Park, Y. G., Kang, H., and Baek, H.: The local stratification preconditions the marine heatwaves in the Yellow Sea, *Front. Mar. Sci.*, 10, 1–9, <https://doi.org/10.3389/fmars.2023.1118969>, 2023.
- Marin, M., Feng, M., Bindoff, N. L., and Phillips, H. E.: Local Drivers of Extreme Upper Ocean Marine Heatwaves Assessed Using
390 a Global Ocean Circulation Model, *Front. Clim.*, 4, 1–16, <https://doi.org/10.3389/fclim.2022.788390>, 2022.
- Marullo, S., Serva, F., Iacono, R., Napolitano, E., di Sarra, A., Meloni, D., Monteleone, F., Sferlazzo, D., De Silvestri, L., de Toma, V., Pisano, A., Bellacicco, M., Landolfi, A., Organelli, E., Yang, C., and Santoleri, R.: Record-breaking persistence of the 2022/23 marine heatwave in the Mediterranean Sea, *Environ. Res. Lett.*, 18, <https://doi.org/10.1088/1748-9326/ad02ae>, 2023.
- McAdam, R., Bonino, G., Clementi, E., and Masina, S. Forecasting the Mediterranean Sea Marine Heat Wave of summer 2022. In:
395 Copernicus Ocean State Report, Issue 8, State of the Planet, 2024
- Mills, K. E., Pershing, A. J., Brown, C. J., Chen, Y., Chiang, F. S., Holland, D. S., Lehuta, S., Nye, J. A., Sun, J. C., Thomas, A. C., and Wahle, R. A.: Fisheries management in a changing climate: Lessons from the 2012 ocean heat wave in the Northwest Atlantic, *Oceanography*, 26, <https://doi.org/10.5670/oceanog.2013.27>, 2013.
- Olita, A., Sorgente, R., Natale, S., Gaberšek, S., Ribotti, A., Bonanno, A., and Patti, B.: Effects of the 2003 European heatwave on
400 the Central Mediterranean Sea: Surface fluxes and the dynamical response, *Ocean Sci.*, 3, 273–289, <https://doi.org/10.5194/os-3-273-2007>, 2007.
- Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V., Benthuyesen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Straub, S. C., and Wernberg, T.: Longer and more frequent marine heatwaves over the past century, *Nat. Commun.*, 9, 1–12, <https://doi.org/10.1038/s41467-018-03732-9>, 2018.
- 405 Oliver, E. C. J., Burrows, M. T., Donat, M. G., Sen Gupta, A., Alexander, L. V., Perkins-Kirkpatrick, S. E., Benthuyesen, J. A., Hobday, A. J., Holbrook, N. J., Moore, P. J., Thomsen, M. S., Wernberg, T., and Smale, D. A.: Projected Marine Heatwaves in the 21st Century and the Potential for Ecological Impact, *Front. Mar. Sci.*, 6, 1–12, <https://doi.org/10.3389/fmars.2019.00734>, 2019.
- Oliver, E. C. J., Benthuyesen, J. A., Darmaraki, S., Donat, M. G., Hobday, A. J., Holbrook, N. J., Schlegel, R. W., and Sen Gupta, A.: Marine Heatwaves, *Ann. Rev. Mar. Sci.*, 13, 313–342, <https://doi.org/10.1146/annurev-marine-032720-095144>, 2021.
- 410 Pastor, F., Valiente, J. A., and Palau, J. L.: Sea Surface Temperature in the Mediterranean: Trends and Spatial Patterns (1982–2016), *Pure Appl. Geophys.*, 175, 4017–4029, <https://doi.org/10.1007/s00024-017-1739-z>, 2018.
- Pastor, F. and Khodayar, S.: Marine heat waves: Characterizing a major climate impact in the Mediterranean, *Sci. Total Environ.*, 861, 160621, <https://doi.org/10.1016/j.scitotenv.2022.160621>, 2023.
- Pastor, F., Valiente, J. A., and Estrela, M. J.: Sea surface temperature and torrential rains in the Valencia region: Modelling the role
415 of recharge areas, *Nat. Hazards Earth Syst. Sci.*, 15, 1677–1693, <https://doi.org/10.5194/nhess-15-1677-2015>, 2015.
- Paulson, C. A., and J. J. Simpson: Irradiance Measurements in the Upper Ocean, *J. Phys. Oceanogr.*, 7(6), 952–956. Payne, [https://doi.org/10.1175/1520-0485\(1977\)007<0952:IMITUO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1977)007<0952:IMITUO>2.0.CO;2), 1977

- Pirro, A., Martellucci, R., Gallo, A., Kubin, E., Mauri, E., Juza, M., Notarstefano, G., Pacciaroni, M., Bussani, A., and Menna, M. Subsurface warming derived by Argo floats during the 2022 Mediterranean marine heatwave. In: Copernicus Ocean State Report, Issue 8, State of the Planet, 2024
- 420
- Pisano, A., Fanelli, C., Buongiorno Nardelli, B., Tronconi, C., Cesarini, C., and La Padula, F.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea High Resolution and Ultra High Resolution Sea Surface Temperature Analysis Product, SST_MED_SST_L4_NRT_OBSERVATIONS_010_004, Issue 3.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-004-006-012-013.pdf>, last access: 13 July 2023,
- 425 2022c.
- Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., and Buongiorno Nardelli, B.: EU Copernicus Marine Service Product User Manual for the Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed Product, SST_MED_SST_L4_REP_OBSERVATIONS_010_021, Issue 2.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-SST-PUM-010-021-022-041-042.pdf>, last access: 13 July 2023,
- 430 2022a.
- Pisano, A., Fanelli, C., Cesarini, C., Tronconi, C., La Padula, F., and Buongiorno Nardelli, B.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed Product, SST_MED_SST_L4_REP_OBSERVATIONS_010_021, Issue 2.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-021-022-041-042.pdf>, last access: 13 July
- 435 2023, 2022b.
- Pisano, A., Fanelli, C., Buongiorno Nardelli, B., Tronconi, C., La Padula, F., and Cesarini, C.: EU Copernicus Marine Service Quality Information Document for the Mediterranean Sea High Resolution and Ultra High Resolution Sea Surface Temperature Analysis Product, SST_MED_SST_L4_NRT_OBSERVATIONS_010_004, Issue 3.0, Mercator Ocean International, <https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-SST-QUID-010-004-006-012-013.pdf>, last access: 13 July
- 440 2023, 2022d.
- Plecha, S. M. and Soares, P. M. M.: Global marine heatwave events using the new CMIP6 multi-model ensemble: From shortcomings in present climate to future projections, *Environ. Res. Lett.*, 15, 124058, <https://doi.org/10.1088/1748-9326/abc847>, 2019.
- Schlegel, R. W., Oliver, E. C. J., and Chen, K.: Drivers of Marine Heatwaves in the Northwest Atlantic: The Role of Air–Sea Interaction During Onset and Decline, *Front. Mar. Sci.*, 8, 1–18, <https://doi.org/10.3389/fmars.2021.627970>, 2021.
- 445 Sen Gupta, A., Thomsen, M., Benthuisen, J. A., Hobday, A. J., Oliver, E., Alexander, L. V., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Perkins-Kirkpatrick, S., Moore, P. J., Rodrigues, R. R., Scannell, H. A., Taschetto, A. S., Ummenhofer, C. C., Wernberg, T., and Smale, D. A.: Drivers and impacts of the most extreme marine heatwave events, *Sci. Rep.*, 10, 19359, <https://doi.org/10.1038/s41598-020-75445-3>, 2020.
- Smale, D. A., Wernberg, T., Oliver, E. C. J., Thomsen, M., Harvey, B. P., Straub, S. C., Burrows, M. T., Alexander, L. V., Benthuisen, J. A., Donat, M. G., Feng, M., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Sen Gupta, A., Payne, B.
- 450

- L., and Moore, P. J.: Marine heatwaves threaten global biodiversity and the provision of ecosystem services, *Nat. Clim. Chang.*, 9, 306–312, <https://doi.org/10.1038/s41558-019-0412-1>, 2019.
- Smith, K. E., Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., Thomsen, M. S., Wernberg, T., and Smale, D. A.: Biological Impacts of Marine Heatwaves, *Ann. Rev. Mar. Sci.*, 15, 119–145, <https://doi.org/10.1146/annurev-marine-032122-455121437>, 2023.
- Sparnocchia, S., Schiano, M. E., Picco, P., Bozzano, R., and Cappelletti, A.: The anomalous warming of summer 2003 in the surface layer of the Central Ligurian Sea (Western Mediterranean), *Ann. Geophys.*, 24, 443–452, <https://doi.org/10.5194/angeo-24-443-2006>, 2006.
- Spillman, C. M., Smith, G. A., Hobday, A. J., and Hartog, J. R.: Onset and Decline Rates of Marine Heatwaves: Global Trends, Seasonal Forecasts and Marine Management, *Front. Clim.*, 3, 1–13, <https://doi.org/10.3389/fclim.2021.801217>, 2021.
- Thoral, F., Montie, S., Thomsen, M. S., Tait, L. W., Pinkerton, M. H., and Schiel, D. R.: Unravelling seasonal trends in coastal marine heatwave metrics across global biogeographical realms, *Sci. Rep.*, 12, 1–13, <https://doi.org/10.1038/s41598-022-11908-z>, 2022.
- Vogt, L., Burger, F. A., Griffies, S. M., and Frölicher, T. L.: Local Drivers of Marine Heatwaves: A Global Analysis With an Earth System Model, *Front. Clim.*, 4, 1–18, <https://doi.org/10.3389/fclim.2022.847995>, 2022.
- Wernberg, T., Bennett, S., Babcock, R. C., De Bettignies, T., Cure, K., Depczynski, M., Dufois, F., Fromont, J., Fulton, C. J., Hovey, R. K., Harvey, E. S., Holmes, T. H., Kendrick, G. A., Radford, B., Santana-Garcon, J., Saunders, B. J., Smale, D. A., Thomsen, M. S., Tuckett, C. A., Tuya, F., Vanderklift, M. A., and Wilson, S.: Climate-driven regime shift of a temperate marine ecosystem, *Science (80-.)*, 353, 169–172, <https://doi.org/10.1126/science.aad8745>, 2016.
- Zhao, Z. and Marin, M.: A MATLAB toolbox to detect and analyze marine heatwaves, *J. Open Source Softw.*, 4, 1124, <https://doi.org/10.21105/joss.01124>, 2019.