



Sea Level Rise in Europe: impacts and consequences

5 Coordinating lead authors: Roderik van de Wal^{1,6} (RvdW), Angélique Melet² (AM)

Authors: Debora Bellafiore³ (DB), Michalis Vourdoukas⁴ (MV), Paula Camus⁵ (PC), Christian Ferrarin³ (CF), Gualbert Oude Essink^{6,7} (GOE), Ivan D. Haigh⁸ (IH), Piero Lionello⁹ (PL), Arjen Luijendijk⁷ (AL), Alexandra Toimil¹⁰ (AT), Joanna Staneva¹¹ (JS)

¹ Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, the Netherlands

10 ² Mercator Ocean International, Toulouse, 31400, France

³ CNR - National Research Council of Italy, ISMAR – Institute of Marine Sciences, Venice, 30122, Italy

⁴ Department of Marine Sciences, University the Aegean, Mithilene, 81100, Greece

⁵ Geomatics and Ocean Engineering Group, Department of Sciences and Technologies of Water and Environment, University of Cantabria, 39005, Santander, Spain

15 ⁶ Department of Physical Geography, Utrecht University, Princetonlaan 8a, 3584 CB Utrecht, The Netherlands

⁷ Deltares, Boussinesqweg 1, 2629 HV Delft, The Netherlands

⁸ School of Ocean and Earth Science, University of Southampton, European Way, Southampton, SO14 3ZH, UK

⁹ Department of Biological and Environmental Sciences and Technologies, University of Salento, Lecce, 73100, Italy

¹⁰ IHCantabria – Instituto de Hidráulica Ambiental de la Universidad de Cantabria, Isabel Torres 15, 39011, Santander, Spain

20 ¹¹ Department on Hydrodynamics and Data Assimilation, Helmholtz Zentrum Hereon, GMBH, Geesthacht, 21502, Germany

Correspondence to: Roderik van de Wal (R.S.W.vandeWal@uu.nl), Angélique Melet (amelet@mercator-ocean.fr)

25 **Abstract.**


Sea level rise has major impacts in Europe which vary from place to place and in time depending on the source of the impacts. Flooding, erosion and saltwater intrusion lead via different pathways to various consequences in coastal regions across Europe. Flooding leads via overflow, overtopping and breaching to damage to assets, environment and people. Erosion leads via cliff failure along a different pathway also to damage and saltwater intrusion affects ecosystems and surface waters and salinizes existing fresh water resources diminishing fresh water availability causing salt damage to crops and health issues to people. This paper provides an overview of the various impacts in Europe.



1 Introduction

Sea-level rise (SLR) is a major threat for coastal zones, inducing hazards such as coastal flooding (mild and chronic at high-tides or intense and episodic during storms), permanent submersion of coastal zones, coastal erosion, salt intrusion in surface and ground water (with adverse impacts on drinkable water and agriculture), problems with water management, and coastal ecosystems degradation or loss (affecting coastal wetlands contributing to coastal zone protection, biodiversity conservation and carbon storage). As sea level is committed to rise over the next centuries (Melet et al., in prep.), coastal zones and communities are expected to be increasingly threatened by sea level changes at various timescales, ranging from episodic extreme events to interannual-to-centennial changes and trends linked to climate change and modes of variability (Bednar-Friedl et al., 2022; Glavovic et al., 2022; Hallegatte et al., 2013; Le Cozannet et al., 2022; Oppenheimer et al., 2019). Locally, these threats are reinforced by subsidence caused by human activities like groundwater pumping or groundwater fluid extraction for energy purposes (Carbognin & Tosi, 2002).

Impacts of SLR result from the combination of sea-level changes, exposure, and vulnerability (Cardona et al., 2012). Sea level changes with a focus on Europe are discussed in Sea Level Rise in Europe: observations and projections. The increased hazards posed by SLR in response to climate change have been identified as the main driver of future rise in coastal flood losses, with the relative importance of trends in exposure (related to coastward migration, urbanization and rising asset values) diminishing over time (Vousdoukas et al., 2018b). However, population and economic activities are expected to change as well, in Europe and worldwide possibly also increasing the exposure. Coastal zones are increasingly more densely populated than the hinterland (Small & Nicholls, 2003), exhibit higher rates of population growth and urbanization, which are concentrating economic assets and critical infrastructures. Coastal migration is driven by the combinations of specific economic, geographic and historical conditions, and includes the concentration of densely settled agricultural areas in well-watered, fertile deltas and coastal plains (Hugo, 2011; McGranahan et al., 2007). The exposure of people and assets to SLR hazards is therefore widespread and increasing; and can thereby extend to a change in the vulnerability to SLR (e.g., urbanization changes the imperviousness of flood-prone areas, Andreadis et al., 2022). Coastal zones in Europe are highly urbanized; more than 30 million people live in the 100-year event flood coastal plain and 50 million in the contiguous and hydrologically connected zone of land along the coast and below 10 m of elevation (sometimes referred to as low-elevation coastal zone (LECZ), (Neumann et al., 2015a) (Figure 1), whether the number of people exposed changes over time depends on assumption on future developments of fertility, mortality and migration.



60

Figure 1: Mean sea level rise (SLR, in m) and coastal population density in Europe for global warming levels of +1.5°C and +3°C. SLR data consider the long-term period (2081-2100) and SSP1-2.6 for (a) and SSP3-7.0 for (b). Adapted from Fig 13.4 of Chapter 13 of AR6 WG2 (Bednar-Friedl et al., 2022).

65 Land use planning strategy in coastal lowlands can reduce exposure of EU population to SLR (see Sea Level Rise in Europe: adaptation measures and decision-making principles). For instance, the establishment of coastal setback zones, which are buffer spaces defined by a specific distance from the shoreline's highest water mark where new developments in potentially exposed coastal regions are restricted. Sanò et al. (2011) can reduce the exposure of new urban development by at least 50% in most EU countries by 2100 (Wolff et al., 2023).

70 Due to the large economic value of coastal zones, economic losses due to coastal flood risks are huge (Abadie et al., 2020; Hallegatte et al., 2013). Presently, the expected annual damage from coastal flooding for Europe alone is around €1.25 billion euros, but could increase by 2-3 orders of magnitudes if coastal adaptation is only maintained to its current level (Vousdoukas et al., 2018a). The impacts of floods (both marine and riverine/ pluvial) on people, the built environment and the economy is one of the four key climate change induced risks identified for Europe (Bednar-Friedl et al., 2022). Deltas are particularly
75 vulnerable to SLR. In Europe, the main deltas are those of the Rhine-Meuse-Escault (NL), Rhone (FR), Pô (IT), and Ebro (SP) rivers. Rotterdam and London are amongst the world's most exposed cities in terms of population living in the 100-year event flood plain, if there was no flood protection (Hallegatte et al., 2013). In addition to human fatalities, economic losses due to coastal flood risks are considerable. European cities where the annual average losses due to coastal flooding by 2050 will increase particularly assuming present-day defense standards or flood probability, tend to be concentrated along the
80 Mediterranean coast (Hallegatte et al., 2013). These cities were built close to the shore since historical sea level variability has been low (e.g., small tidal range and interannual variability), as a consequence changes in the mean are felt earlier than in regions where the sea level variability is higher.

Therefore, SLR creates risks for people, ecosystems, land uses, the built environment, and human activities (Le Cozannet et al., 2022). In this paper, a summary of the 6th cycle of the IPCC reports is provided (Section 2). Then, various impacts of SLR
85 are discussed with a European perspective, using the Source, Pathway, Receptor and Consequence (SPRC) framework



introduced in Section 3. SLR impacts are discussed for coastal flooding (Section 4), coastal erosion (Section 5) and salt-water intrusion (Section 6).

2 Summary of previous assessments

An updated assessment of the impact and risks for natural and human systems by SLR is provided by the recent 6th Assessment Report of the Intergovernmental Panel on Climate change (IPCC AR6), whose Working Group II (WG2, Pörtner et al., 2022) assessed impacts, adaptation and vulnerabilities related to climate change. SLR is considered in many chapters, particularly in chapter 3 “Ocean and coastal ecosystems and their services” (Cooley et al., 2022, see Cross Chapter Box 3 for an overall summary), and chapter 6 “Cities, settlements and key infrastructure” (Dodman et al., 2022), with more focussed material presented in the Cross Chapter Paper 2 “Cities and settlements by the sea” (Glavovic et al., 2022). Material directly addressing European regional issues is included in chapter 13 “Europe” (Bednar-Friedl et al., 2022) and in the Cross Chapter Paper 4 “Mediterranean region” (Ali et al., 2022). The material on SLR covered by AR6 WG2 (Dodman et al., 2022) and the AR6 Synthesis Report (Calvin et al., 2023)) builds on and updates the former IPCC assessment in the “Special Report on the Ocean and Cryosphere in a Changing Climate” published in 2019, particularly in its Chapter 4 “Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities” (Oppenheimer et al., 2019).

There is extensive evidence that at global scale, relative SLR is already impacting ecosystems, human livelihoods, infrastructure, food security and climate mitigation potential at the coast (Pörtner et al., 2022). Observed impacts include chronic flooding at high tides, more frequent episodic flooding during storms, wetland, and underground water salinisation and ecosystem transitions, increased erosion and coastal flood damages. SLR poses risks for cities, settlements and populations in low-elevation coastal zones, cultural heritage along coasts, and threatens the very existence of some island nations. The large exposure caused by the disproportional concentration of population and economic activities in coastal areas lead to high risks, which are very likely to increase further with future SLR. In addition, risks in the agriculture sector and nature conservation are induced by the salinisation of groundwater, estuaries, wetlands and soils. The IPCC AR6 WG2 shows that these risks will generally increase with SLR. At global scale and in the absence of further adaptation and mitigation action, they will be very likely one order of magnitude larger in 2100 than during present-day. Along many European coastlines, extreme water levels, coastal floods and sandy coastline recession are projected to increase during the 21st century, because of the increase in relative SLR.

According to the IPCC AR6 WG2, there is high confidence that acceleration of SLR will increase risks to people and infrastructures from inundation and extreme floods along European low-lying coasts and estuaries. Related damages will increase at least tenfold even before the end of the 21st century assuming present levels of adaptation and mitigation measures. Annual damage (which today is 1.3 billion EUR) is expected to increase disproportionately with global warming. It will be in the range 13–39 billion EUR by 2050 at global warming levels between 2°C and 2.5°C, and 93–960 billion EUR by 2100




between 2.5° and 4.4°C temperature increase. Assuming present distribution of population and protection levels, the increase of people at risk depends on the emission scenario: with respect to present, an extra 10 million people will be at risk of a 100-
120 year flood event under a very high-emission scenario (RCP8.5) by 2100, whereas just below 10 million people under a low emission scenario (RCP2.6) by 2150. Along low-lying coasts and estuaries flood risks might further increase because of compounding storm surges, waves, rainfall, and river runoff events, but to date this has been poorly quantified (see also section 4). Port operations may be negatively affected by SLR in northern and western Europe. In Mediterranean ports negative effects are to be expected by a combined change of wave regimes and sea-level.

125 In the IPCC AR6 WG2 it is shown that soft cliffs and beaches in Europe are most affected by coastal erosion. Observations suggest that 27–40% of Europe’s sandy coast are already eroding today. Though there is no convincing evidence that past erosion of sandy shorelines can be attributed to climate change or SLR. At the same time there is high confidence that SLR will increase sandy shoreline retreat in the future. Actual rates are uncertain, with large spatial variability and a strong dependence on local geometry, but it is suggested that by 2100 coastal retreat could reach approximately 100 m for a 4°C
130 temperature increase, being reduced to less than 60 m under 3°C of global surface warming. On a centennial time-scale coastal erosion and flooding will become an existential threat for some coastal communities and UNESCO World Heritage sites. This is particularly the case in the Mediterranean region, where seawater intrusion in coastal aquifers is projected to increase by the combination of overexploitation and SLR, with pronounced impacts on agricultural productivity.

The IPCC AR6 WG2 shows that warming is the main climate hazard for European coastal ecosystems. However, rapid SLR
135 (potentially aggravated by human induced subsidence) is also expected to have negative impacts by reducing the surface area of intertidal flats (e.g. the Wadden Sea) that cannot always be compensated by sediment accumulation. By 2100 under intermediate scenarios coastal erosion will cause the loss of 4.2–5.1% of present values of ecosystem services and reduce their contribution to shoreline protection across Europe. The vulnerability of Europe’s coastal subtidal seagrass meadows and intertidal salt marshes to SLR is particularly high in the microtidal areas of the Baltic and Mediterranean coasts, with a potential
140 loss of 75% of Posidonia oceanic seagrass habitats in the Mediterranean Sea.

3 The Source, Pathways, Receptors and Consequences framework

To describe the main impacts of SLR across Europe, we use the concept of Source-Pathway-Receptor-Consequence (SPRC), proposed by Sayers et al. (2002) as an alternative approach to the traditional, exposure vulnerability approach. The ‘source’ describes the origin of a hazard. The ‘pathway’ is the route that a hazard takes to reach the ‘receptors’, and includes processes
145 and characteristics of the coastline influencing or mediating the hazard. The ‘receptor’ is the entity (e.g., people, property, environment) that may be harmed by the hazard, and the ‘consequences’ are the corresponding social, economic and environmental effects on the receptors. The concept is illustrated in Figure 2. The SPRC concept will be applied for coastal flooding (Section 4), erosion (Section 5), and salt-water intrusion (Section 6).



150

Figure 2: Source-Pathway-Receptor-Consequence (SPRC) conceptual model (after Sayers et al., 2002)

4 Coastal flooding and compounding flood events

The first pronounced impact of SLR we consider is coastal flooding. A schematic overview of the SPRC concept for coastal flooding is provided in Figure 3. Coastal floods are amongst the most impactful hazards both in Europe and globally, with widely ranging social, economic and environmental consequences. Many severe flooding events have affected European coastlines throughout history (Ferrarin et al., 2022; Haigh et al., 2015, 2017; Melet et al., in prep.; Paprotny et al., 2018). An increase in coastal flooding frequency is one of the most certain and costly consequences of SLR (R. J. Nicholls & Cazenave, 2010). Flood-defense standards in many European countries are among the highest in the world (Galluccio et al., in prep.). However, significant populations and assets are in coastal flood plains and are threatened when defense infrastructure fails or is exposed to flooding exceeding the protection standard. Furthermore, impacts of coastal flooding are likely to increase as flood defense infrastructure is ageing, as coastal population continues to grow, and urbanization and development in low-elevation coastal zones continue (Stevens et al., 2016; McMichael et al., 2020). In addition, further decline of the extent of natural habitats (i.e., saltmarshes, mudflats, shingle beaches and sand dunes), which act as a natural buffer to flooding, has the potential to increase coastal flood risks (Campbell & Keddy, 2022).

165



Coastal Flooding




Figure 3: The Source-Pathway-Receptor concept visualised for flooding.

Multiple drivers, some related to climate change and others not, influence coastal flood risk and its future change. These different drivers can be considered using the SPRC conceptual model as explained in Section 3. The ‘source’ of coastal floods (describing the origin of governing hazards) is formed by extreme coastal water levels (including the contributions from tides, storm surge, wave runup, superimposed on relative mean sea level). The influence of climate change on the source component of coastal flooding is discussed in Section 4.1, and in sections 4.3 and 5.3 in (Melet et al., in prep.). In estuaries, the compounding influence of rainfall and fluvial input can be important leading to compound flooding, which is described as a special case in Section 4.2. The ‘pathway’ represents how seawater makes its way onto normally dry land to cause flooding. Climate change, especially SLR, can significantly affect this pathway in addition to modifying the source profile. This influence of SLR on pathways is discussed in Section 4.3. Receptors and consequences are closely linked and are discussed jointly in Section 4.4. Initiatives to develop flood-related climate services in Europe are discussed in Section 4.5.

180 4.1 Source of flooding

Coastal floods are governed by anomalously high-water levels exceeding a site-specific threshold. Extreme coastal water levels (ECWLs) arise as combinations of different drivers: (1) astronomical tides; (2) storm surges and associated seiches; (3) waves, including setup, infragravity waves and swash (e.g., (Dodet et al., 2019)); and (4) relative mean sea level (including SLR and land subsidence) (see also Sea Level Rise in Europe: observations and projections). These four components exhibit



185 considerable intra- and inter-annual variability (e.g., driven by tidal cycles (Haigh et al., 2011) or climate variability, such as
the North Atlantic Oscillation (Boucharel et al., 2023; Hurrell, 1995; Mentaschi et al., 2017); or changes in wave climate
(Morim et al., 2019; Melet et al., 2020). In addition, there are non-linear interactions between the four components (Horsburgh
& Wilson, 2007; Idier et al., 2019; Arns et al., 2020). Long-term changes in any or all of the four components can modify the
variability of frequency and magnitude of ECWL, and thus affect coastal flooding. The additional influence of rainfall and
190 fluvial input on water levels can be important in estuaries, discussed in Section 4.2 in relation to compound flood events.
ECWLs, and hence the frequency of coastal flooding events, are impacted by climate change in three main ways (Pugh &
Woodworth, 2014) as also argued in Sea level rise in Europe: observations and projections: (1) SLR affects the ECWL directly,
by raising the baseline mean water level leading to lower storm surge and/or wave elevations necessary to cause flooding; (2)
rising mean sea levels alter water depths and therefore modify the propagation and dissipation of the tide and storm surge
195 components, or alter wave processes in shallow water (e.g., Arns et al., 2017; Chaigneau et al., 2023); and (3) climate variations
in the tracks, speed, strengths and frequency of weather systems may alter the intensity and/or duration and frequency of storm
surges and waves (and variations in rainfall and river discharge in estuaries).

As discussed in Sea level rise in Europe: observations and projections (Sections 4.3, 5.3), direct changes in mean sea levels
appear to have been the main driver of observed changes in ECWL in the past (Ferrarin et al., 2022; Marcos et al., 2015;
200 Menendez & Woodworth, 2010) and are projected to dominate changes in extremes along the European coastline in the future,
increasing the likelihood of coastal flooding (Vousdoukas et al., 2016, 2017). However, changes in storm surges (Calafat et
al., 2022; Calafat & Marcos, 2020; Muis et al., 2020) and wave height (Aarnes et al., 2017; Benetazzo et al., 2022; Vousdoukas
et al., 2018b) have and may also play a substantial role in the changes in ECWL in some European regions in the future.
Coastal flooding could also be influenced by changes in tides, especially in regions with a shallow water depth. Regionally
205 coherent changes (positive and negative) in tidal range have been observed in historic sea level records around both European
and global coastlines, and are projected to occur in the future with changes in water depth, driven by SLR and factors such as
ice sheet extent and ocean warming (Ferrarin et al., 2015; Haigh et al., 2020; Idier et al., 2017; Pickering et al., 2012). Changes
in tidal range are likely to be smaller than $\pm 15\%$ of mean SLR along most coastlines.

The relative contribution of SLR to changes in coastal flooding depends on different factors. Areas with tides below 2 meter
210 amplitude will be primarily affected by mean sea level changes. For example, in the Mediterranean Sea, extreme sea levels
that are now occurring once in a century are expected to occur at higher frequency in future depending on climate scenarios
but in some places even more frequently than annually. This intensification is stronger than the overwash and flooding in the
South of Portugal, where current return periods of 1-in-100-year can reduce to lower than 1-in-20-year by 2055 and 1-in-10-
year by 2100 (Ferreira et al., 2021). Venice stands as an exception for the Mediterranean Sea, as the existing protection
215 measures are higher than for most of the south European coastlines and could potentially protect the monumental city during
the next few decades (Lionello et al., 2021; Mel et al., 2021).



4.2 Compound Flooding

Many coastal settlements along the European coastline are in estuaries and lagoons (e.g., London, Rotterdam, Hamburg, Venice, Lisbon). In these coastal regions, floods can arise not only from climate driven changes in oceanographic sources (e.g., tides, storm surges and waves), but also via river discharge (fluvial) and direct surface runoff (pluvial). These mainly arise from heavy precipitation, but are also incurred via snow melt. Many cities and towns located on the open coastline can also experience flooding during heavy rainfall, because of insufficient drainage during high tides (Van Den Hurk et al., 2015a). In the past, flood risk assessments typically considered the oceanographic, fluvial and pluvial drivers of flooding separately. However, in coastal regions floods are often caused by more than just one factor, which can be physically correlated (e.g., with storms). Furthermore, the adverse consequences of a flood can be greatly exacerbated when the oceanographic, fluvial, and/or pluvial drivers occur concurrently or in close succession (i.e., a few hours to days apart), depending on local characteristics which influence lag times between variables. This can result in disproportionately extreme events, referred to as ‘compound flood events’. (Zscheischler et al., 2018) define compound events as ‘a combination of multiple drivers and/or hazards that contributes to societal or environmental risk’. Flood drivers are typically causally related through associated weather patterns (the modulator, Zscheischler et al., 2020) and therefore, it is assumed that stronger dependence between drivers increases the impact of compound floods (Wahl et al., 2015).

In recent years there has been a large increase in the number of studies that have started to investigate compound flood events in Europe. Many studies have been undertaken for specific localised regions in Europe, such as the Rhine delta, Netherlands (Kew et al., 2013; Khanal et al., 2018); Brest, France (Mazas & Hamm, 2017); Santander, Spain (Rueda et al., 2016); Ravenna, Italy (Bevacqua et al., 2017); Venice, Italy (Ferrarin et al., 2022); and the river Trent, the Yare basin, the river Ancholme, and the rivers Taff and Lewes in East Sussex in the UK (Granger, 2001; Mantz & Wakeling, 1979; Thompson & Law, 1983; Samuels & Burt, 2002; C. J. White, 2007). Larger-scale assessments of compound flood events have been undertaken more recently for the UK (Hendry et al., 2019; Svensson & Jones, 2002, 2004) and for Europe (Bevacqua et al., 2019; Bevacqua et al., 2020a; Camus et al., 2021, 2022; Paprotny et al., 2018; Petroliaqkis et al., 2016). On a quasi-global scale, which obviously includes Europe, Ward et al. (2018) and Couasnon et al. (2020) assessed the dependence between coastal and river flooding, using observational datasets and reanalysis, respectively.

Most of these studies quantified the statistical dependence between flooding sources as an indirect measure of the flooding hazard called compound flooding potential. The analysis of the interdependencies has been primarily limited to storm surge and precipitation (compound surge-rain events, Wahl et al., 2015; Wu et al., 2018), or to surge and river discharge (compound surge-discharge events, (Couasnon et al., 2020; Hendry et al., 2019; Ward et al., 2018). In those studies, precipitation is considered as a fluvial proxy, which can be assumed as an equivalent driver to discharge in small to medium-sized river catchments (Bevacqua et al., 2020b). Waves can be included as part of the total sea level, by adding it linearly to the storm surge and/or astronomical tide components (Bevacqua et al., 2019). Statistical studies have notably shown that: (a) the joint exceedance probability of compound surge-discharge events is on average a factor of 2-4 higher when the dependence is



considered (Santos et al., 2021; Van Den Hurk et al., 2015b; Ward et al., 2018); (b) ignoring the dependence between precipitation and surge can overestimate the flooding return period considerably (Bevacqua et al., 2019; Figure 4a); (c) the river discharge of the 50-year compound flood is up to 70% larger, conditional on the occurrence of extreme water levels (Ganguli & Merz, 2019a). Simulations of the non-linear interactions of these flood drivers at local scale using coupled
255 modelling approaches demonstrate a rise of extreme sea levels at some locations along the estuaries (e.g., in the Netherlands, Van Den Hurk et al., 2015; and in northwestern Spain, Bermúdez et al., 2021). Recently, a global analysis of simulated river flood levels using a global coupled river-coast flood model framework, showed that surge exacerbates 1-in-10 years flood levels at 64% of the river mouths analysed, with a mean increase of 11 cm (Eilander et al., 2020). Furthermore, 55% of the world coastlines face compound storm surge and wave extremes which increases the potential coastal flooding (Marcos et al.,
260 2019).

Although different sampling methods to identify compound events and dependence measures to quantify compound flooding potential have been used, several hotspots have been identified along the European coastlines. High joint occurrences of extreme river discharge and storm surge events are found on the coasts of Portugal, the Strait of Gibraltar, the western facing coasts of North and Central Europe (Heinrich et al., 2023), and along the south-west coast of the UK (Hendry et al., 2019).
265 Also, the northern and eastern Mediterranean coasts and the coast of Tunisia appear as compound flooding hotspots (Camus et al., 2021; Couasnon et al., 2020; Eilander et al., 2020). Regarding precipitation and surge, higher dependency is concentrated along the Atlantic coast and in the Mediterranean Sea (particularly in the regions of the Gulf of Valencia, Spain, northwest Algeria, the Gulf of Lion, France, the Adriatic coast of Balkan Peninsula, the Aegean coast, southern Turkey and the Mediterranean Levantine region) (Bevacqua et al., 2019; Camus et al., 2021, Figure 4a).

270 Historical trends in compound flooding resulting from high coastal water levels and peak river discharge have been assessed over north-western Europe (near the North and the Baltic Seas) over 1901–2014 using 37 stream gauges (Ganguli & Merz, 2019b). Increasing trends were identified in the region from 47°N to 60°N while decreasing trends were identified along higher latitude coasts (>60°N).

In most regional, continental, or global dependence-based analyses, compound events are identified using a two-sided
275 conditional sampling to bivariate drivers (Camus et al., 2021; Couasnon et al., 2020; Wahl et al., 2015; Ward et al., 2018) which implies that events are either conditioned to one driver or the other. Another approach is to select pairs of high values when both variables exceed individual high percentiles (e.g., 95th percentile, as in Bevacqua et al., 2019). However, extreme water levels might be driven by events not being extreme themselves. Impact-focused approaches, modelling the relationship between extreme water levels and underlying drivers, allow the selection of large impact events whose drivers are not necessary
280 extreme (Bermúdez et al., 2019; Bevacqua et al., 2017; Santos et al., 2021). Most of these studies rely on modelled data products which generally capture compound flooding, but contain biases, false positive or missed extremes (Paprotny et al., 2020).



Climate change can affect flooding dynamics through SLR, changes in each of the flood drivers and in the interaction processes between them. SLR can increase future compound flood hazard (Ganguli et al., 2020; Moftakhari et al., 2017). However, 285 uncertainties due to internal climate variability and climate model differences dominate the large uncertainty in the concurrence of flood extremes in addition to changes of the drivers (e.g. storm surges or river discharge) themselves (Bevacqua et al., 2021). Large-scale sea-level/rainfall-driven compound flooding potential is projected to increase globally by more than 25% by 2100 under RCP8.5 scenario compared to present times (Bevacqua et al., 2020a). The probability of compound flooding due to the co-occurrence of high sea level and precipitation is projected to robustly increase by the end of the 21st century 290 particularly in northern Europe, e.g., along the west coast of Great Britain, northern France, the east and south coast of the North Sea, but also along the coastlines of the eastern half of the Black Sea (Bevacqua et al., 2019, Figure 4b). Increases in compound flood events were observed to be mostly due to the rising mean sea level (Heinrich et al., 2023). These results contrast with a predominant decrease of compound flood hazard projected for north-western Europe at the middle of the 21st century (Ganguli et al., 2020), which may be caused by the fact that the two studies considered different flooding drivers and 295 multi-model ensembles.

In terms of pathways for the impact of compound flooding, elevated sea levels can block or slow down river drainage into the sea, leading to increased upstream water levels which can overflow river channels onto adjacent floodplains thereby increasing inundation extent. Besides, floodwater can also erode riverbanks and cause landslides, leading to further flooding. Hydrodynamic modeling is required to provide a detailed spatial mapping of water levels in estuaries or coastal river deltas. 300 However, high-resolution hydrodynamical modelling, which includes the non-linear interaction between hydraulic processes, topography, and human interventions is only feasible at local scale.

In terms of receptors and consequences, historical information on past damaging floods reveals that compound events have occurred in many locations in Europe. According to the HANZE database (Paprotny et al., 2018), out of 1564 floods that occurred in 37 European countries between 1870 and 2016, 23 (i.e. 1.5%) were identified to be compound floods, recorded in 305 six different countries (the Adriatic Sea—Italian regions of Veneto and Friuli-Venezia Giulia, Mediterranean and western coast of France, Ireland, UK, Belgium and Poland). Specific examples of compound events include: flooding in December 2000 in Brittany, France – 600 persons were affected by a coastal flood that occurred due to the combination of heavy precipitation over several river catchments and a storm surge generated by an extra-tropical storm (Paprotny et al., 2018); and a flood in December 1999 in Lyminster, UK – the town was flooded due to tidal locking of the high run-off down the 310 Lyminster River by a large surge produced by the same storm system, with this flooding overwhelming recently upgraded defences (Ruocco et al., 2011; Hendry et al., 2019).




Figure 4: (a) Present day (1980-2014) probability of potential compound flooding (CF). Return periods of CF (co-occurring sea level and precipitation extremes, i.e., larger than the individual 1-year return levels) based on ERA-Interim data. (b) Future probability of potential CF multi-model mean of projected change (%) of CF return periods, between future (2070-2099) and present (1970-2004) climate (Bevacqua et al., 2019).

4.3 Pathway for coastal flooding

315 Seawater tends to inundate normally dry land via three main pathways: (1) by still water simply overflowing, where the water
level exceeds the elevation of a natural (e.g., barrier beaches) or artificial (e.g., sea wall) barrier; (2) by waves overtopping a
natural or artificial barrier; and (3) by breaching and lowering of a natural or artificial barrier, often as a consequence of
prolonged overwashing or erosion at the front-face of the barrier or by groundwater seepage (Figure 3). Climate change, and
other factors, can influence these pathways, altering local flood risk profiles. For example, SLR in regions with hard structures
320 like barriers or dykes typically lead to a decline in the extent of natural habitats, such as saltmarshes, mudflats, and sand dunes,
which can act as a natural buffer to flooding (Hall et al., 2019). Decline in these natural features (and deterioration of flood
protection infrastructure) can impact flood pathways and can increase flood hazard. In contrast, building new or maintaining
and improving existing flood defences, or application of artificial nourishment and stabilisation of beaches, or providing more
space for water through managed re-alignment, can alter flood pathways and reduce flood risk along coasts (Haigh et al.,
325 2022).

It is increasingly recognised that natural systems that provide important buffering against floods are in decline across parts of Europe. For example, Campbell and Keddy (2022) identified a 136 (confidence interval 39–236) km² loss of salt marsh extent from 2000 to 2019 across Europe. Other green adaptation options include restoration of seagrass meadows (which reduce wave height and sediment erosion) and the creation of buffer zones (Wolff et al., 2023). With mean SLR accelerating over the



330 coming century, and thereby increasing pressure on the narrow coastal zone there is likely to be a continued decline in the
extent of natural systems contributing to natural buffers reducing coastal flood risk, which will lead to defence capital and
maintenance costs increasing dramatically (Haigh et al., 2022). This is corroborated by projections of shoreline retreat along
most of the global shorelines induced by SLR (Vousdoukas et al., 2020; Section 5) and a consequent reduction of ecosystem
services (Paprotny et al., 2021).

335 Obviously, current flood risk around the coastline of Europe would be considerably higher without the decades of investment
into extensive flood risk management infrastructure. Data on flood defences over time are not well documented, but massive
investments in defences have occurred over the 20th and early 21st century in Europe. For example, extensive flood defence
infrastructure has been built in the Netherlands as part of the Delta Works and standards of protection along stretches of the
Dutch coastline now reach 1-in-10,000-year levels (Eijgenraam et al., 2014). Governing policy directives incorporate future
340 sea level rise into the periodic risk assessments and defence strategy updates (Kothuis and Kok, 2017). Nearly a quarter of
England's coast is now defended (Sayers et al., 2015). As for France, more than 16,000 coastal works and developments are
covering more than 2,300 km of coastlines, implying 17% of metropolitan French coasts are shaped by human interventions.
Around 20 movable storm surge barriers have been built around the coast of Europe since 1958 offering flood protection to
millions of people and trillions of Euros of infrastructure (Mooyaart and Jonkman, 2017). This includes six surge barriers in
345 Netherlands (e.g., Eastern Scheldt and Maeslant Barrier, Figure 5), thirteen in the UK (e.g., Thames and Hull Barriers), and
one in Italy (the MOSE system for protecting the historical city of Venice and the lagoon settlements, operating since October
2020), while a new surge barrier is being constructed in Belgium (Nieuwpoort). However, storm surge barriers can be used to
mitigate the flood impact only in semi-enclosed coastal environments (such as bays, estuaries, lagoons) but not along coasts
facing the open sea. Extensive beach nourishment has also taken place in many European countries to counteract coastal
350 erosion and flooding. For example, the Dutch coast is one of the most heavily nourished coasts globally (Brand et al., 2022);
since 1990, more than 300 nourishments programs have taken place (including the notable Sand Engine project; Roest et al.,
2021) adding an average of 12 million m³ annually to the 432 km of Dutch coastline.

However, as sea-levels continue to rise it will become increasingly costly to maintain existing flood defences and surge
barriers, and to carry out coastal nourishments. In the UK, Sayers et al. (2015) showed that length of coastal defences 'highly
355 vulnerable' to failure would almost double (triple) under 0.5 m (2.5 m) mean SLR, with the number of properties affected
rising by around 160% (490%). Furthermore, many of the existing storm surge barriers will need to be strongly upgraded or
replaced over the coming century as sea-level rises. For example, plans are underway to replace the Thames Barrier around
2070, with options including a new barrier built farther downstream of the current one (Environment Agency, 2021). The
Eastern Scheldt Barrier (Figure 5) was designed for only 40 cm SLR, implying that a revision is needed to avoid it to be closed
360 too often with higher sea-level (Haasnoot et al., 2020) with as consequence that the availability of maintenance windows
decreases too much.



Figure 1. Map of the Netherlands showing flood prone zones (blue shadings) and features of the water management system (adapted from PBL [58]). N.A.P. is the Amsterdam Ordnance Level which is the reference plane for sea level height in the Netherlands.

Figure 5: Map of the Netherlands showing flood prone zones (blue shadings) and features of the water management system. NAP is the Amsterdam Ordnance Level which is the reference plane for sea level height in the Netherlands. Extracted from Haasnoot et al., 2020.

365

4.4 Receptor for and Consequences of flooding


The receptors and consequences associated with coastal flood events can be broadly grouped into social (e.g., loss of life, number of people evacuated, damage to residential property, or loss of cultural heritage), economic (e.g., overall monetary cost, disruptions to ports, transport, energy, public services, water systems, agricultural production losses) and environmental (e.g., coastal erosion, degradation or losses of coastal habitats, damage) impacts (Haigh et al., 2017). Importantly these consequences can be long-lasting (e.g., injury or long-term physical and mental health effects; Jackson and Devadason, 2019) and can also extend outside of the coastline area directly impacted by flooding (e.g., disruption to transport or supply chains; Dawson et al., 2016). As SLR and increases in storminess enhance flood risk and its consequences, the number of receptors in flood-prone areas will grow accordingly. Changes in land use and increasing asset values in floodplains can also increase consequences of coastal flooding (Haigh et al., 2022). In contrast, improvements in flood forecasting, early warning, emergency response and planning can greatly reduce the consequences of flooding (Section 4.5). Careful spatial planning and building codes can be effective at reducing risk. Evidence from Haigh et al. (2017) suggests that the number and consequences of coastal floods have declined since 1915 in the UK, reflecting better defences and improvements in flood forecasting,

370

375



380 warning, emergency response and planning (Haigh et al., 2022). As a concrete example, more than 2000 lives were lost around
the coastlines of the North Sea during the flood of 1953; however, similar conditions occurring in December 2013 hardly had
any societal impact, which can be attributed to these improvements and infrastructure investments (Wadey et al., 2015).
As shown in Figure 6, the flooded area from the 100-year event is projected to increase by 10%-15% in 2050, depending on
the emissions scenario, and by 12%-20% by the end of the century. For more rare events, like a 1000-year event, SLR will
385 drive a 48%-67% increase in the flood extent by the year 2100 (Paprotny et al., 2019). The country level flood extents depend
on different factors, such as the exposure to extreme weather conditions, the protection standards in place, as well as the
country's size and the percentage of low-lying coastal areas. For the baseline, the UK and Norway have the largest flood extent
area, exceeding 4,500 km² for the 100-year event for each country. This is a result of their long coastlines exposed to intense
weather conditions. Denmark and Germany follow with values around 3,000 km², and for these cases the main driver is the
390 flat and low-lying configuration of the coastal zones. Other countries with flood extents slightly below 2,000 km² are Greece
and Italy, both characterized by long coastlines.



395 **Figure 6: Indication of the current area vulnerable for flooding with a return period of once in a hundred years and for two different time slices under two different climate scenarios. From Paprotny et al. (2018).**

It is noteworthy that also events with low flood levels (i.e., nuisance flooding, Moftakhari et al., 2018) may have high impacts from disruption of everyday routine activities and property damages, especially in low land settlements. SLR is expected to increase not only the frequency of ECWL but also the frequency of low-level floods, when astronomic tide will become a major driver of flood events (Ferrarin et al., 2022). Towards the end of the century SLR will also result in permanent flooding of certain areas. For example, in the Balearic Islands and by the year 2100, 7.8–27.7 km² and up to 10.9–36.5 km² will be permanently lost to SLR under RCP4.5 and RCP8.5 emission scenarios, respectively (Luque et al., 2021).



Pan-European assessments of future coastal flood risk show general trends and allow for regional comparisons, but also come with large uncertainty due to data scarcity and non-stationary conditions (Hinkel et al., 2021; Vousdoukas et al., 2018a) and cannot replace local studies (Paprotny et al., 2019). For example, flood assessment results are very sensitive to the coastal protection standards assumed which are largely unknown along most of the European coastline (Scussolini et al., 2016). Another crucial factor is digital elevation data. Since several countries lack high-resolution LIDAR data, many assessments are based on less accurate global datasets, which come with vertical biases exceeding the extent of anticipated SLR (Bove et al., 2020; Kulp & Strauss, 2018; Yamazaki et al., 2017). Despite these limitations, all known flood risk assessments highlight an increase in flooded area during the century, which also accelerates as SLR gathers pace.

4.5 Initiatives to develop flood-related climate services in Europe

The EU Adaptation Strategy emphasises the significance of climate services in adapting to climate change. As defined by the Global Framework for Climate Services (Hewitt et al., 2012) of the World Meteorological Organization (WMO), "Climate services provide climate information to help individuals and organizations make climate smart decisions". According to the European Commission's Roadmap for Climate Services (European Commission, 2015) definition, climate services cover "the transformation of climate-related data - together with other relevant information - into customized products such as projections, forecasts, information, trends, economic analysis, assessments (including technology assessment), counselling on best practices development and evaluation of solutions and any other services in relation to climate that may be used for the society at large".

Information on past and future sea level change close to the coastline is being made available, but still suffers from various limitations (e.g., resolution, incomplete physics in models, scarcity of observations). In parallel, core European services have been put in place. Yet, authoritative, consistent and decision-oriented climate services to support policies and decision-making with SLR are still in their early development worldwide.

The European Union's Earth Observation Programme (Copernicus) monitors our planet and its environment, for the ultimate benefit of society. This includes the monitoring of sea level changes and the provision of ancillary fields needed to assess coastal SLR risks, as well as the transformation of the wealth of satellite, in situ and integrated numerical model information into added-value datasets and information usable by scientists, managers, decision-makers and the wider public to guide adaptation and to support related policies and directives (Melet et al., 2021).

New initiatives in the framework of dedicated European research projects in the Horizon Europe program are also underway, for instance the Coastal Climate Core Services (CoCliCo) project. CoCliCo aims to deliver an open web platform that will provide up-to-date information on present and future SLR and its impacts to support decision-making on coastal flood risk management and adaptation. The platform will grant access to the latest and consistent hindcast and projections of sea level, process-based coastal flood maps and shoreline change estimates, flood exposure and vulnerability information, as well as adaptation strategies and options. Users of the platform will be able to visualize, download, and analyse high-quality geospatial



information layers encompassing multiple decision-oriented coastal risk scenarios. For shorter timescales, Early Warning Systems (EWSs) are integrated systems allowing a real time monitoring of potential natural hazards, issuing natural hazard warnings with a few days lead time. Informing the relevant stakeholders (e.g., civil protection agencies, regional and local authorities, environmental agencies) is part of an integrated risk management procedure to mitigate risks. EWSs can play a critical role in classical disaster risk management cycles, supporting the preparedness and response phases, including the deployment of emergency measures for rapid response after a disaster, as well as longer term damage assessment after the occurrence of an event. EWSs are an efficient adaptation measure by providing more than a tenfold return on investment (Global Commission on Adaptation, 2019). The H2020 ECFAS project aims at contributing to the evolution of the Copernicus Emergency Management Service (CEMS) by demonstrating the technical and operational feasibility of a European Coastal Flood Awareness System. Such a system will complement the existing early-warning system of the CEMS for river/pluvial floods, CEMS-EFAS (European Flood Awareness System), by adding a pan-European marine coastal flood awareness system and by tackling coastal resilience to climate risk (marine storminess and exposure). ECFAS provides an integrated risk cycle monitoring and management service, from water level forecasts at the coast with a 5-day lead time (Irazaqui Apecechea et al., 2023), rapid mapping of coastal floods and impacts on population and assets (Le Gal et al., 2022), and risk and recovery Mapping (RRM) for adding coast-targeted products in the aftermath of a marine flood event (e.g., shoreline displacement, maps of flooding and damages). At national scale, the Norwegian mapping authority developed a webtool for inundation mapping showing extreme still water levels and projected sea level, including statistics on the areas, roads, and buildings affected now and in the future (Breili et al., 2020, <https://www.kartverket.no/en/at-sea/se-havniva/se-havniva-i-kart>). The webtool includes statistics on the areas, roads, and buildings affected now and in the future. The German Sea Level Monitor ([https://meeresspiegel-monitor.de/index.php.en](https://meeresspiegel-monitor.de/index.php/en)) provides observed and projected changes at tide gauges along the German North Sea and Baltic Sea coast.

5 Coastal erosion

5.1 Definition and drivers of coastal erosion

Coastal erosion is the permanent loss of land to the sea. Coastal erosion can take place at different time scales and is driven by a wide range of natural and anthropogenic factors. Among the different types of coastal systems are estuaries, lagoons, barrier islands, sandy and gravel beaches, dunes, cliffs, rocky shores, and built areas. Sandy beaches are especially prone to erosion (EUROSION, 2004a, 2004b). They occupy 31% of the ice-free global coastline (Luijendijk et al., 2018) and tend to be the most heavily utilized beach typology (Davenport & Davenport, 2006). Sandy beaches also tend to be geomorphologically complex and dynamic, affected by natural and anthropogenic factors (Mentaschi et al., 2018). A large part of the European coastline is developed by humans, and the built area deprives coastal systems from their natural capacity to accommodate, or recover from erosion (Small & Nicholls, 2003). In addition, human developments and dams tend to prevent terrestrial sediments from reaching the coastline, which favours coastal erosion (Milliman, 1997). As a result of the above, Europe's



beaches have been eroding (EUROSION, 2004a, 2004b; Masselink et al., 2022; Romagnoli et al., 2022), a trend which is projected to accelerate with climate change and SLR (Vousdoukas et al., 2020).

470 One of the consequences of erosion is shoreline change which is the combined result of numerous factors, such as wind and wave climates, terrestrial sediment supply, geological control and human interventions, among others. There is a clear cause and effect relationship between increasing sea levels and shoreline retreat (Bruun, 1962), which justifies the eroding trend reported by either large scale (Hinkel et al., 2013; Vousdoukas et al., 2020), regional (Alvarez-Cuesta et al., 2021; Toimil, Losada, Camus, et al., 2017), or local scale studies (Alvarez-Cuesta et al., 2021; de Santiago et al., 2021; Luque et al., 2021; 475 Romagnoli et al., 2022; Toimil, Losada, Díaz-Simal, et al., 2017). Negative sediment budgets can be another factor driving a robust erosion trend (López-Olmedilla et al., 2022). The former can be the result of several potential natural or anthropogenic factors. In addition, several of the observed changes can relate to quasi-periodical climatic patterns affecting the wave regime (e.g., Barnard et al., 2015).

Apart from major drivers like the ones mentioned above, beach erosion and accretion often depend on a delicate balance of 480 contrasting forces with very similar amplitudes and are therefore very difficult to predict. This applies to all scales; i.e. each wave can transport sand both towards and away from the coast (Vousdoukas et al., 2014), while storms drive erosion that can be followed by complete or partial recovery (e.g., Kroon et al., 2008; Lee et al., 1999). Moreover, the impact of extreme storm events is very much controlled by the initial beach morphological state, and thus by meso-to-macro scale processes (e.g., Qi et al., 2010; Vousdoukas et al., 2012). In this Section, the source, pathway, receptor and consequence are discussed for coastal 485 erosion in Section 5.2. Monitoring methods, including field surveys, video monitoring and Earth observation, are discussed in Section 5.3. Historical shoreline changes in Europe are summarized in Section 5.4 while projected shoreline changes are summarized in Section 5.5. Finally, interactions between coastal erosion and coastal flooding are addressed in Section 5.6.

5.2 Source/Pathway/Receptor/Consequence for coastal erosion

490 *Source for erosion*

Coastal erosion shares the same sources as coastal flooding; e.g., all the components that drive ECWL (tides, storm surges, waves and SLR) can drive coastal erosion, Figure 7. In addition, given that coastal erosion is strongly related to the sediment budget, sediment sources and sinks are very important. As a result, human development and dams play an important role as they tend to prevent terrestrial sediments from reaching the coastline favoring coastal erosion (Anthony et al., 2019; Milliman, 495 1997).



Coastal Erosion




Figure 7 Source, Pathway, Receptor Consequence framework for coastal erosion.

Pathway for erosion

500 Coastal erosion takes place when the sediment budget of a given area becomes negative. Understanding coastal change is a very challenging task since erosion or accretion is a result of multiple factors like: (i) processes that take place in various temporal and spatial scales (Kroon et al., 2008; Larson & Kraus, 1994); (ii) the presence of various features which emerge through complex, self-organization processes often linking different scales and processes (Murray et al., 2009; B. T. Werner, 1999); (iii) the intrinsic uncertainty in predicting the intensity and frequency of extreme events, as well as the related beach
505 morphological response (Vousdoukas, 2012); and (iv) the high complexity of long-term processes like sediment transport and vertical land motion, which are also interconnected, among others, with geological and meteorological phenomena (Gallop et al., 2011; Vousdoukas et al., 2007), as well as human interventions which are increasing in number and extent (Luijendijk et al., 2018; Mentaschi et al., 2018). In principle, coastal erosion can be the result of any process that alters the sediment transport patterns. This can be either hydrodynamic (e.g., changes in wave intensity or direction, sea level, etc) (Sierra & Casas-Prat,
510 2014), related to the presence of obstacles like hard structures (Loureiro et al., 2012; Noble, 1978), or factors that affect the erodibility of the beach (Feagin et al., 2019).

Receptor and consequences for erosion

515 Coastal erosion is the process by which the land is worn away and is submerged in water. Coastal erosion can take many forms, such as the loss of sand dunes, cliffs, or beaches and can have several consequences. Among them are the destruction of buildings, roads, and other infrastructure located near the coast.



Moreover, coastal erosion can have significant impacts on the environment, economy, and human health and safety, as it can lead to habitat loss for coastal species, reduce the recreational value of beaches, and increase the risk of flooding and storm damage for coastal communities.

520

5.3 Monitoring methods for coastal erosion

Coastal monitoring is crucial to gaining a better understanding of the likely impacts of climate change at the coast. While the long-term implications of SLR have received considerable attention, Nicholls et al. (2007) point out that more attention needs to be given to finer temporal and spatial scales, including the localized impacts of potential changes in wave climate and storminess regimes. However, monitoring the coast at higher temporal (daily to decadal) and spatial (three-dimensional) resolutions presents many challenges. Conventional survey techniques for ongoing beach surveys are both costly and labor-intensive, and meaningful trends typically require several years of data to emerge (Short & Trembanis, 2004). Currently, only half a dozen multi-decadal, high-resolution coastal monitoring programs are in operation worldwide, with only 2 in Europe: Truc Vert (France) since 2005, and Noordwijk (Netherlands) since 1964.

525

530

The various monitoring methods for coastal erosion are described briefly in Table 1, highlighting the evolution from traditional to present-day monitoring techniques.

Table 1: Summary of the methods for monitoring coastal erosion

Type of method	Brief description of method
Field surveys	GPS surveys have proven to be a successful method for beach topographical profiling on sandy coasts (Hansen & Barnard, 2010; Harley et al., 2011). This technique is commonly used for monitoring morphological changes in the short term and involves repeating measurements at regular intervals to understand the physical aspects of coastal environments, including daily, monthly, and annual variations of specific parameters (Komar, 1998; Short, 1999). Spatial scales of such surveys are typically from 100m's to several km's.
Video monitoring	Coastal video monitoring systems (e.g. Holman and Stanley, 2007; Stringari and Power, 2022) are ideal for long-term deployments (i.e., years) acquiring data continuously, covering few km of coastline. Images are processed to generate the system's 'basic products': time-averaged, variance, snapshot and timestack images, which are all projected in geographic coordinates using standard photogrammetric techniques. Then, a set of post-processing tools allows extracting quantitative information on various coastal processes at various temporal and spatial scales, such as beach-face/shoreline morphology, inner bar configuration, rip/longshore current systems and wave run-up. Moreover, they have been proved to be useful coastal management tools, supporting sustainable and safe recreational beach use (Jiménez et al., 2007).



Drones	For the past 10 years, drones or Small Unmanned Aerial Vehicles (e.g. Vousdoukas et al., 2011) have been used more frequently for coastal monitoring (Chapapria et al., 2022). These unmanned aerial vehicles are equipped with cameras and other sensors that allow researchers to collect high-resolution images and data over km’s of the coastline. This technology offers several advantages over traditional methods of monitoring. Drones can cover a larger area than ground-based surveys and provide more detailed images than satellite imagery. Additionally, drones can be deployed quickly and easily, making them ideal for collecting data in remote or hard-to-reach areas.
Terrestrial 3D Laser Scanning	Terrestrial 3D Laser Scanning (TLS) or Terrestrial LIDAR has increasingly become the method of choice for beach surveying (e.g., Pietro et al., 2008). Portable scanners allow the completion of beach surveys in excess of hundreds of meters, with sub-centimeter spatial resolution, during the course of a few hours. These advances provide a reliable means of addressing geomorphic relationships, as well as along- and cross-shore changes on the beach; while the accuracy and spatial resolution is practically impossible with traditional GPS surveys. The technique has been very recently introduced in coastal research and has several unexploited possibilities.
Earth Observation	<p>In recent years a new source of geospatial data for studies from regional to planetary scale is provided by earth observation satellites generating an ever-increasing flow of raster image data. An exponential increase in the availability of free geospatial data has recently emerged with the Copernicus Earth observation and monitoring program of the European Union that delivers satellite imagery complemented by in-situ observations (Malenovský et al., 2012).</p> <p>The positional accuracy of satellite derived shorelines (SDS) based on single images has been evaluated to range between 1.6 and 10 m (e.g. Liu et al., 2017). The increasing availability, resolution and spatial coverage of satellite imagery in recent years now provide a powerful alternative to derive reliable, global scale shoreline data.</p>

535

5.4 Historical shoreline change

A comprehensive European study of coastal erosion collected and analysed aerial photographs and local surveys up to 2002 to estimate coastal erosion at the scale of Europe (EUROSION, 2004a). About 20% of the European Union’s coastline suffered serious erosion impacts, with the area lost or seriously impacted estimated at 15 km² per year. More recent studies at continental to global scale confirmed that large stretches of the European coast are suffering from erosion. Using freely available optical satellite images captured since 1984, in conjunction with machine learning and image analysis methods, the shoreline changes have been mapped at global scale (Luijendijk et al., 2018; Mentaschi et al., 2018). The application of an automated shoreline detection method to the sandy shorelines resulted in a global dataset of shoreline change rates for the 33-year period 1984–2016 (Luijendijk et al., 2018). Analysis on satellite-derived sandy beach detection reveals that about 35% of the European



545 coastline is sandy, which agrees largely with the 40% EuroSION estimate (EUROSION, 2004a). Analysis of the satellite derived
shoreline data indicates that 22% of the European sandy beaches are eroding at rates exceeding 0.5 m/yr, while 26% are
accreting and 52% are stable. This means that in Europe a total of more than 8,200 km of sandy beaches have significantly
retreated over the last decades. Areas that experience severe erosion are found at various locations across Europe (see Figure
8). About 4% of the European sandy beaches experience erosion rates classified as severe (> 3m/year). Erosion rates exceed
550 5 m/yr along 2% of the sandy shoreline. The statistics of Europe are rather similar to the global statistics stating that 24%
(28%) of the world's beaches are eroding (accreting) (Luijendijk et al., 2018).




Figure 8: Long-term shoreline changes of the European beaches in m/yr since 1984. The red and green points indicate erosion and accretion respectively (data based on Luijendijk et al. (2018)).

555

5.5 Future shoreline change

Mediterranean beaches are more susceptible to the negative effects of SLR because they are narrower as a consequence of the beach slope. This is highlighted by both large scale (Vousdoukas et al., 2020) and regional scale projections (Monioudi et al., 2017). For example, a recent study in the Balearic Islands projects at least 20% of the islands' beaches losing more than 50%
560 of their surface by the end of the century, even if greenhouse gas emissions are mitigated (Luque et al., 2021). But even projections along the Atlantic coast report shoreline retreat; e.g., in the range of 10-45 m under the middle-of-the-road (RCP4.5) scenario and 14-66 m under the very-high emission (RCP8.5) scenario by the year 2100 for 150 km of Basque coast (de Santiago et al., 2021). A recent pan-European study projects a mean SLR driven median shoreline retreat of 97 m (54 m)



565 under RCP8.5 (respectively under RCP4.5) by the year 2100. This retreat translates to 2,500 km² (1,400 km²) of SLR driven
coastal land loss (Athanasίου et al., 2020), Figure 9. Given the complexity of coastal morphodynamics, projections of shoreline
changes come with high uncertainties. The future variability of wave forcing is more prominent until 2060 with respect to
uncertainties, whereas after that year the uncertainties in predicting sea-level rise become dominant (D'Anna et al., 2021,
2022). Of similar amplitude is also the uncertainty associated with the choice of geophysical datasets in continental scale
assessments (Athanasίου et al., 2020), reaching 45% (26%) of the variance in coastal land loss projections for Europe by 2050
570 (2100).




Figure 9: Projections of shoreline retreat in m (a) and land loss in km² per km of coastline (b) for Europe in the year 2100 under a very high emission scenario (RCP8.5) (from Athanasίου et al., 2020).

575

5.6 Coastal erosion and flooding interactions

Coastal erosion and flooding are extremely interrelated impacts that influence each other (Leaman et al., 2021; Pollard et al., 2019; Sallenger, 2000). Erosion is a physical phenomenon where sand is removed from the shoreface and deposited elsewhere, usually offshore. Erosion and deposition processes can change the shoreface, which affects coastal flooding caused by high
580 water levels. In turn, these high-water levels can cause further erosion or deposition. This feedback manifests itself on different time scales. In the short term, coastal morphology plays a significant role in wave energy dissipation, total water levels reaching the coast, dune breaching, and subsequent flooding. SLR is expected to increase the frequency of episodic erosion and flooding, which in turn could be altered by changes in storminess. At longer time scales, SLR is expected to drive permanent erosion and inundation (R. J. Nicholls & Cazenave, 2010; Cazenave & Le Cozannet, 2014). This effect could be compensated or
585 enhanced in areas with intense alongshore gradients in longshore sediment transport or chronic fluvial sediment supply. Higher



water levels cause wave-driven erosion to occur higher up in the profile resulting in net erosion and deposition on the nearshore bottom (Bruun, 1962). Additionally, deeper water reduces wave refraction and allows waves to get closer to the shore before breaking (Arns et al., 2017; Chaigneau et al., 2023), leading to increased flooding. A coastline subject to sustained erosion over time may lead to the loss of natural flood defenses (Toimil, Losada, et al., 2023; Toimil, Álvarez-Cuesta, et al., 2023).

590 The need for the joint modelling of flooding and erosion has long been recognized in the literature (Bilskie et al., 2014; Lentz et al., 2016; Passeri, Hagen, Bilskie, et al., 2015). However, most studies to date continue to analyse these two impacts separately because of the complex relationship between driving processes and morphological response (Pollard et al., 2019; Toimil et al., 2020; Toimil, Losada, et al., 2023). Amongst the studies that couple flooding and erosion, most of them have typically focused on historical events without considering climate change (Gharagozlou et al., 2020; McCall et al., 2010; Van

595 Ormond et al., 2020). These studies have primarily been conducted at the storm scale and have used pre- and post-storm topobathymetry data to simulate erosion and breaching using a hydro-morphodynamic model. Following the same modelling approach but in the context of climate change, some studies have incorporated the effect of SLR on hydrodynamic forcing conditions (Grases et al., 2020; Passeri et al., 2018). Sanuy and Jimenez (2021) presented a more recent application in the Tordera Delta (Spain) where the baseline topo-bathymetry was modified to consider medium-term erosion.

600 Studies that consider long-term shoreline changes in flooding follow more diverse approaches. Stripling et al. (2017) delivered flood maps considering long-term changes in seawall toe levels along the west coast of Calabria (Italy) and in an idealised coastal stretch around Holderness (UK). As for climate change studies, Dawson et al. (2009) developed a methodology to account for shoreline changes in coastal flood projections along the East Anglian coast (UK). Dawson et al. (2009) linked coastal flooding due to storms and SLR with long-term erosion by adjusting the likelihood of flood defence structures failure

605 based on shoreline changes. Other studies have examined the effect of shoreline changes on coastal flooding considering empiric (Grilli et al., 2017) and surveyed profile translation due to SLR (Barnard et al., 2019) and due to SLR and changes in the sediment budget (Passeri et al., 2016; Passeri, Hagen, Medeiros, et al., 2015). Also using real profiles (Toimil, Álvarez-Cuesta, et al., 2023) proposed a suite of numerical and statistical models to analyse the influence of storm morphodynamics, SLR, erosion and longshore sediment transport on total water levels and coastal flooding along a 40-km coastal stretch in the

610 Spanish Mediterranean.

Current studies highlight the need to consider the interconnections between hydrodynamics and morphodynamics to better understand the functioning of the coastal system. Changes in topographic representations can alter the path and pattern of maximum water levels (Bilskie et al., 2014). (Toimil, Álvarez-Cuesta, et al., 2023) found that total water levels are mainly affected by storm erosion and profile geometry, and that long-term erosion is the main shoreline change contributor to the

615 flooded area.

To date, existing research in the EU focuses on the East Mediterranean and North Sea basins. In the East Mediterranean, studies encompass a wider range of coastal typologies than in the North Sea, including cliffs, deltas, and beaches with varying levels of anthropisation. Most studies model the interaction of medium and long-term coastal changes with episodic flooding,



620 but few additionally consider storm erosion as flood enhancer. This can be particularly important for extreme weather events
of high return periods, regardless of the type of coastline. The largest gap lies in studies that combine the interaction of hydro-
morphological processes under climate change scenarios, as such studies are very limited in both basins. However, knowledge
on the potential effects of these interconnections and the enhancing role of SLR is key information to decision makers to make
informed decisions for the future management of coastal communities.

6 Saltwater intrusion

625 6.1 Processes and monitoring of saltwater intrusion

Saltwater intrusion (SWI) represents the increased extent of the mixing zone between inland freshwater and saltwater (Source),
therefore rising the salt content both in surface waters and in groundwaters (Pathway). SWI can hinder the use of water for
agriculture due to salt damage to crops (e.g., Maas and Hoffman, 1977) and can threaten coastal communities that rely on fresh
water supplies for their livelihood (Receptor) (e.g., too salty drinking water can lead to cardiovascular diseases, (He &
630 MacGregor, 2011)), see Figure 10.





Figure 10 A visualisation of the Source Pathway Receptor framing for saltwater impacts.

Rivers and aquifers (good permeable water-bearing porous media) contaminated by high salinity decrease freshwater storage
and water quality, reduce soil fertility, impact on vegetation and freshwater species, and affect human health. Moreover, in
635 deltaic regions, SWI can also negatively impact ecosystem services and aquaculture activities such as clam or shrimp farming
(e.g., Dierberg and Kiattisimkul, 1996; Hou et al., 2022) (Figure 11). SWI is a slow but increasing hazard affecting European
coasts, especially in deltas and estuaries. In these low-elevation coastal zones, climatic changes (including sea-level rise)



combine with the changes induced by human activities such as reducing river flows causing salt-wedge shifts (Maselli & Trincardi, 2013). Figure 11b shows several processes causing problems induced by SWI such as extracting groundwater, creating controlled low-lying areas (polders) and reducing in urbanized areas fresh water depletion into the groundwater systems (sealing). As a result of climate change (and associated SLR) and induced human activities, impacts of SWI are likely to increase (e.g., (Befus et al., 2020; Oude Essink et al., 2010). In this section we discuss the main impacts of SWI, again using the source (Section 6.2), pathway (Section 6.3), receptor/consequence (Section 6.4) framework.



645 **Figure 11: Salinisation of (ground)water resources and land for different stages of progressive development. Salt water is represented in red / pink, brackish water in purple, fresh water in blue. (a) Situation in the distant past, before human settlements; (b) current situation; (c) situation in the future without adaptation; (d) situation in the future with adaptation. Source: Deltares (2023).**

650 The scarcity of data often poses a challenge for sustainable management of groundwater resources worldwide. Mapping and monitoring the spatial extent of near-coastal fresh groundwater resources usually requires detailed information for large coastal regions, which is often not available whereas in-situ measurements are time- and labor consuming. To address this challenge, remotely sensed data can be a cost-effective way to gather both surface and groundwater information, covering a large area in a short period. For instance, Airborne Electromagnetic geophysical methods are particularly useful for detecting groundwater



655 salinity affecting the conductivity of the groundwater. Such methods have been executed in Denmark (Duque et al., 2022),
Germany (Siemon et al., 2015) and The Netherlands (Delsman et al., 2018).

6.2 Source for surface saltwater intrusion


Surface SWI threatens water resourcing and freshwater availability especially in low-lying coastal areas such as deltas and
660 estuaries (van Engelen et al., 2022), which are characterized by natural complex interactions between fresh and salty waters
(Horner-Devine et al., 2015; Valle-Levinson, 2010). The inland intrusion of salty waters along the river courses is controlled
by the forces acting at both the river and the sea domains: the advective dispersion associated with the river flow, the steady
shear dispersion associated with the estuarine exchange flow, and the tidal pumping (Lerczak et al., 2006). The balance among
these forces is regulated mostly by the combined action of river discharge and sea level oscillations (Bellafiore et al., 2021).
665 Surface heat fluxes (evaporation) and the salt content in corresponding sea water also play a role in determining the extent of
surface SWI. Therefore, while more evident in drought conditions (e.g., the extended 2022 drought in the Po Valley - Italy;
Bonaldo et al., 2023), the processes regulating SWI, and the concurring effects cannot be attributed just to one single driver.
Moreover, these natural drivers can act both on short timescales, as tidal fluctuations, storm surges and hurricanes, and at the
long timescales, as climatic fluctuations, subsidence and, among others, SLR. In general, the variation of surface SWI does
670 not only affect the extension of the affected area, but can lead to the predominance of some hydrodynamic processes, as, for
example, shifting the system from a diffusive to an advective dominated river dynamics (e.g., the Po River). Also, a
modification of the environmental conditions in transitional areas can vary the extent of eu-, poly-, meso- and oligo-haline
areas (Rodrigues et al., 2019). The concurring processes affecting surface SWI are linked to progressive river discharge
decrease, increased surface heat fluxes (evaporation) and increase in the salt content and in the relative sea level. In Europe,
675 several estuarine and deltaic systems are suffering from the progressive increase in surface SWI. Not surprisingly, this process
occurs both in micro and macro tidal environments (e.g., the Po River and Elbe Estuary, respectively), with different sea
salinity values (ranging from less haline ones in the Atlantic or North Sea to the more haline ones in the Mediterranean) and
higher or lower surface heat fluxes. These environmental conditions trigger the predominance of one of the several saltwater
drivers. As an example, the Tagus Estuary is exposed to a tidal excursion of up to 3.8 m, also amplified by resonance. Therefore,
680 even in normal conditions, sea saltier water intrudes affecting 43% of the estuary area (intertidal zone). The combination of
tides and the periodic exposure to droughts, with several low discharge events, seem to massively affect the system (Rodrigues
et al., 2019).

The Atlantic coasts already include cases of increased saltwater intrusion in the groundwater system (Figure 12). Studies on
the quantification of effects of changed drivers show their relative effect. In the Minho and Lima estuaries, in the northern
685 coast of Portugal, the future SLR scenarios identify a progressive increase in saltwater extension, and the effect of most extreme
SLR scenarios is leading to a transgression of the saltier front of several kilometers (Pereira et al., 2022). In this case, SLR is
identified as the dominant driver for increased saltwater intrusion, compared to future river discharge reduction. On the other
hand, the quantification of the relative effect of SLR and reduced river discharge in future climate change scenarios leads to



opposite conclusions in a microtidal Mediterranean system, as the Po River delta. River discharge reduction affects SWI more
690 than SLR (Bellafiore et al., 2021). SLR and climate change induced salinization is predicted to worsen also in several coastal
locations in the North Sea (e.g., The Netherlands - Bonte and Zwolsman, 2010; Belgium - Bertels and Willems, 2022). In all
deltaic and estuarine systems, the evaluation of changes in drivers should be carried out considering possible additional long-
term modifications of the morphologic environment, such as subsidence.

Added to these natural factors and often acting in combination with them, several anthropogenic activities can exacerbate SWI
695 by lowering the surface freshwater supply. Examples are changes in land use and land drainage, irrigation, hydropower
production and over-exploitation of coastal aquifers.



700 **Figure 12:** Case-studies where groundwater salinisation is occurring in the coastal zone around Europe (updated and after Post et al., 2018, and Van Weert et al., 2009), where it may cause multifaceted impacts combined with sea-level rise changes.


6.3 Pathway for saltwater intrusion

For SWI, the pathway reflects how sea water reaches coastal freshwater resources causing their salinization. In estuaries and
deltas, rivers are the preferential way to transport sea water upstream, through salt-wedge intrusion along the riverbed and by
705 lateral inflow of the river into the aquifers. Land salinization and aquifer contamination can also result from coastal inundation
with saline waters, through floods induced by storm surges. The coastal morphology and geological characteristics are
therefore strongly influencing the saltwater pathway determining complex hydrogeological interactions between groundwater,
surface water and marine water. As explained in Section 6.2, climate change may enhance SWI drivers by raising the mean
relative sea level as well as by reducing the net freshwater supply in rivers and aquifers. However, as for coastal flooding, SLR
710 will also shorten the pathway of saltwater to reach land and freshwater resources by squeezing the coastline, thereby facilitating



coastal inundation and intrusion of marine waters. In addition, several anthropogenic activities exacerbate SWI by altering the river mouth and coastal morphology, such as diversions of water bodies, river channel deepening, saltmarsh reduction and human induced subsidence (White and Kaplan, 2017).

715 Several prevention and adaption measures have been undertaken in the last decades to limit coastal inundation and ingression
of salty waters along the river channels and the aquifers in Europe. Anthropogenic interventions can affect SWI--impacted
areas by increasing the downstream flow of freshwater (e.g., river diversion, optimization of freshwater withdrawals and
deliveries) or by preventing the upstream transport of saline water. All flood barriers and measures limiting land inundation
described in Section 4 have the co-benefit of limiting salinization of soil and intrusion of salt waters into surface and
720 groundwate systems. The most adopted engineered strategy to prevent salt wedges from intruding in estuaries and deltas is
the installation of (often submerged) mechanical barriers (gates, dams, dikes, levees) near the river mouth that physically block
the upstream flow of saline water (White and Kaplan, 2017). However, salt barriers are regularly damaged and breached during
floods and are not effective during extreme droughts when the saline layer occupies the largest portion of the water column.
Regarding groundwater, subsurface barrier walls (such as sheet piles, clay trenches and injection of chemicals) are considered
725 as one of the most effective methods for inhibiting SWI (Armanuos et al., 2020). Various coastal Managed Aquifer Recharge
(MAR) schemes can be implemented to mitigate groundwater salinization in the coastal zone (Dillon et al., 2019; Oude Essink,
2001) (see Figure 13). Sprenger et al. (2017) identify successful coastal MAR systems for The Netherlands (Amsterdam, The
Hague areas), Spain (Barcelona Llobregat delta), Italy (Po delta), Belgium (De Panne), and Portugal (Algarve). These include
increasing artificial recharge in upland areas to enlarge the outflow of fresh groundwater through the coastal aquifer, injecting
730 or infiltrating (purified) fresh water near the shoreline to create freshwater injection barriers, land reclamation to create a
foreland where a freshwater body can develop or delay the inflow of saline groundwater. This is for instance a consequence
of the sand suppletion ('The Sand Engine' at Zuid-Holland, The Netherlands) where the foreland better protects the low-lying
hinterland against flooding (Huizer et al., 2016).



735




Figure 13: Concept proposed for managing water resources in coastal areas, ensuring a reliable supply of freshwater for domestic, industrial, and agricultural purposes. To store excess freshwater, large-scale coastal Managed Aquifer Recharge (MAR) and Aquifer Storage and Recovery (ASR) methods can be used (see text). Additionally, brackish groundwater can be intercepted to prevent the salinization of aquifers and low-lying areas, while also providing a source of freshwater after desalination (after
740 <https://www.coastar.nl/> COASTAR, 2018).

6.4 Receptor and consequences of saltwater intrusion

The progressive salinization of water resources has severe and long-lasting consequences on several social (e.g., reduction of drinking water reservoirs), economic (e.g., water systems, agricultural production and land losses), and environmental (e.g.,
745 degradation or losses of freshwater habitats, changes in biodiversity) issues. According to Cooper et al. (1964), groundwater SWI is a natural occurrence in coastal groundwater systems. However, human activities such as excessive freshwater pumping from coastal aquifers (Custodio, 2002; Custodio & Bruggeman, 1987; Mastrocicco & Colombani, 2021; Schmork & Mercado, 1969) often disrupt this natural process and even cause land subsidence (Minderhoud et al., 2017). The interaction between groundwater and surface water can also be significant, with saline groundwater sometimes exfiltrating towards surface water
750 systems (De Louw et al., 2011; Delsman, 2015) and negatively impacting agricultural use and nature (Stofberg et al., 2015). In densely populated and industrialized coastal regions around the world (Neumann et al., 2015b), including areas in Europe, groundwater often serves as the primary source of freshwater. Several studies have examined the future availability of groundwater worldwide under climate change and related SLR (Green et al., 2011; Taylor et al., 2013) and human activities (e.g., Wada et al., 2010, 2014). Projected SLR is expected to exacerbate water stress in these densely populated areas,
755 potentially leading to overexploitation and salinization of groundwater resources due to upcoming saline water (Figure 14). This situation is further compounded by the growing demand for freshwater in the future. Groundwater extraction in deltas leads to accelerated salinization of coastal freshwater aquifers and land subsidence if groundwater infiltration is limited by low-permeable layers (Herrera-García et al., 2021). The combination of land subsidence, SLR and sand mining in riverbeds also increases the risk of SWI in estuaries and floods (Eslami et al., 2021).

760 The influence of SLR on coastal groundwater systems has been studied since the 1990s (Navoy, 1991; Oude Essink, 1996; Sherif & Singh, 1999). More recent contributions concern (global) conceptual analyses based on analytical comparisons of fresh-salt interfaces (Chang et al., 2011; Chesnaux, 2015; Chesnaux et al., 2021; Ferguson and Gleeson, 2012; Mazi et al., 2013; Werner and Simmons, 2009), 2D cross-sectional model studies (Ketabchi & Jahangir, 2021; Michael et al., 2013), and 3D model studies (Befus et al., 2020; Loáiciga, 2009; Mabrouk et al., 2018; Masterson & Garabedian, 2007; Oude Essink et al., 2010; Rasmussen et al., 2013; Vandenbohede et al., 2008; Delsman et al., 2023), whereas the effects of SLR on coastal
765 groundwater systems has been reviewed in Ketabchi et al. (2016).



770 **Figure 14: Saltwater intrusion in a coastal aquifer where too much groundwater is extracted leading to upconing of saline groundwater into the extraction well.**

7 Ecosystems and estuaries

SLR can negatively impact coastal ecosystems and estuaries (e.g., degradation of marshes, wetlands, salt water intrusions), either through SLR itself or indirectly through measures implemented to protect the land from flooding or other adverse effects. On the other one hand, several coastal ecosystems can only be adapted or be maintained under narrow ranges of SLR rates
775 (e.g., (Timmerman et al., 2021)). On the other hand, coastal defenses and measures implemented to protect coastal zones from SLR, such as flood barriers (e.g., Orton et al., 2023) also influence coastal ecosystems. These impacts can include increased water levels in estuaries, which may lead to the inundation of adjacent low-lying lands, shoreline erosion, and the potential failure of coastal and stormwater infrastructures (Hanslow et al., 2018; Pachauri et al., 2014; Sweet and Park, 2014). As such, the sustainable management of estuaries, adjacent low-lying areas, and their associated ecosystems requires a thorough
780 understanding of the complex impacts of (accelerating) SLR on estuarine processes (Saintilan et al., 2018; Schuerch et al., 2018). This knowledge will inform effective strategies for adaptation and mitigation in the face of SLR challenges.

Estuarine habitats, located at the interface between land and sea, face significant pressures from anthropogenic stressors originating within their catchment areas, including sedimentation, eutrophication, and pollution (Smith et al., 2013). These stressors often result from excessive inputs of terrestrial sediment, nutrients, and contaminants. Additionally, the estuaries'
785 ecosystems are vulnerable to the effects of global-scale climate change, including SLR and associated impacts. However, the ecological consequences of SLR and related factors have received relatively little attention (Beukerna, 2002; Rullens et al., 2022). SLR will have a significant impact on coastal and ecological risks, including the reduction and degradation of coastal biodiversity. While further data and analysis are needed to fully understand the combined evolution of these effects, it is evident that the decline in ecosystem services will exacerbate the vulnerability of coastal systems. Once these systems reach a



790 threshold or tipping point, their restoration becomes challenging, and their ability to adapt to new climate or management conditions is severely limited.

Intensified flooding and erosion resulting from SLR pose increasing risks to coastal activities and infrastructure (Genua-Olmedo et al., 2016; Sections 4 and 5). These risks are often projected to amplify under future climate scenarios. A reduction in riverine fluxes contributes to a sedimentary deficit and accelerates relative SLR (Ericson et al., 2006). To restore such
795 systems effectively, progressively larger volumes of sediment are required to compensate for SLR and subsidence (see Sea Level Rise in Europe: observations and projections), maintaining the relative sea levels and ensuring deltaic sustainability.

8 Conclusions

In this chapter, the main impacts of SLR, namely coastal flooding, erosion and saltwater intrusion, have been reviewed using the concept of the Source-Pathway-Receptor-Consequence. Regarding coastal flooding, SLR, along with changes in storm
800 surge and wave height, and in some place's increases in tidal range, has driven an increase in the frequency with which extreme coastal water levels are exceeding high thresholds (Source). This in turn, along with an ongoing decline in the extent of natural habitats that act as a buffer to flooding (Pathways) and rapid population growth and urban encroachment in flood-prone areas (Receptors), has driven an increase in coastal flooding and its impact (Consequences) around the coast of Europe. However, current flood risk around the coastline of Europe would be considerably higher without the decades of investment into
805 extensive flood risk management infrastructure and advances in flood forecasting and emergency response. At the same time, losses in major events that exceed defence design standards are growing and are expected to increase massively in the future with higher rates of SLR, unless further adaptation is taken. Furthermore, events with low flood levels (i.e., nuisance flooding) are likely to increase causing widespread disruption of everyday routine activities and property damages, especially in low lying areas.

810 Coastal erosion shares the same source as flooding; i.e. extreme waves and storm surge, even though in many areas additional long term anthropogenic, geological and climatic factors are also important, as they affect sediment budgets (Pathways). A total of more than 8,200 km of Europe's sandy beaches have significantly retreated over the last decades, found at various locations across the continent (Receptors). Climate change and rising seas are expected to accelerate the current trend and retreating shorelines combined with built areas backshore will result in coastal squeeze and threaten sandy beaches. The above
815 will not only have social, economic, and ecological consequences, but is expected to also exacerbate coastal flooding.

In many regions in Europe, saltwater intrusion into surface and groundwater systems is emerging as a problem. Sea-level rise is an important driver (Source). SLR facilitates coastal inundation and intrusion of marine waters into fresh water resources (Pathway). But other (human) processes also play an important role such as groundwater extractions and low-lying areas
820 attracting salt water and reduced fresh river flows. These processes diminish fresh water availability leading to health risks caused by drinking too salty water, less fresh water for economic purposes (e.g. salt damage to crops) and nature (stresses on



existing biodiversity). A wide variety of adaptation measurements has been developed over the last decades, reducing the impact of saltwater intrusion. At the same time projected SLR is expected to increase the impact of saltwater intrusion in the densely populated coastal areas of Europe, particular in combination with the growing demand for freshwater in the future.

825 In summary, we can conclude that the impacts of SLR are emerging at many places across Europe and it might be expected that these impacts on freshwater availability will increase over time, thereby providing an incentive for further mitigation measures whereas at the same time smart adaptation measures are needed to reduce the impacts itself.

Author Contributions

830 RvdW and AM coordinated the paper. RvdW let the writing of the abstract. Section 2 was led by PL. IH and RvdW drafted section 3. IH and AT drafted section 4. MV, PC and AL drafted section 5. GUO, CF, DB drafted section 6 and JS section 7. All author contributed to the different versions of the entire manuscript. Kate Davis from Southampton University drafted the figures 2,3,7 and 10.

835 Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgements

A. T. acknowledges financial support from the Ministerio de Ciencia e Innovación (MCIN/AEI and
840 NextGenerationEU/PRTR) through the Ramon y Cajal Programme (RYC2021-030873-I). I.D.H time was supported by the UK Natural Environment Research Council grant CHANCE (NE/S010262/1). AM and AT were supported by CoCliCo, which has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 101003598. D.B. and C.F. contribution is embedded in the scientific work of the ESFRI DANUBIUS Research Infrastructure – The International Centre for Advanced Studies on River-Sea Systems (<http://www.danubius-ri.eu/>). RvdW received funding from the NPP.
845



850 References

- Aarnes, O., Reistad, M., Breivik, O., Bitner-Gregersen, E., Eide, L. I., Gramstad, O., Magnusson, A. K., Natvig, B., & Vanem, E. (2017). Projected changes in significant wave height towards the end of the 21st century- Northeast Atlantic. *Journal of Geophysical Research - Oceans*, *122*, 3394–3403.
- Abadie, L. M., Jackson, L. P., Sainz de Murieta, E., Jevrejeva, S., & Galarraga, I. (2020). Comparing urban coastal flood risk
855 in 136 cities under two alternative sea-level projections: RCP 8.5 and an expert opinion-based high-end scenario. *Ocean & Coastal Management*, *193*, 105249. <https://doi.org/10.1016/j.ocecoaman.2020.105249>
- Ali, E., Cramer, W., Carnicer, J., Georgopoulou, E., Hilmi, N., Le Cozannet, G., & Lionello, P. (2022). Cross-Chapter Paper
4: Mediterranean Region. In H.-O. Pörtner, C. D. Roberts, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria,
M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, & B. Rama (Eds.), *Climate Change 2022: Impacts,
860 Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change* (Cambridge University Press, pp. 2233–2272).
[10.1017/9781009325844.021](https://doi.org/10.1017/9781009325844.021)
- Alvarez-Cuesta, M., Toimil, A., & Losada, I. J. (2021). Modelling long-term shoreline evolution in highly anthropized coastal
areas. Part 2: Assessing the response to climate change. *Coastal Engineering*, *168*, 103961.
865 <https://doi.org/10.1016/j.coastaleng.2021.103961>
- Andreadis, K. M., Wing, O. E. J., Colven, E., Gleason, C. J., Bates, P. D., & Brown, C. M. (2022). Urbanizing the floodplain:
Global changes of imperviousness in flood-prone areas. *Environmental Research Letters*, *17*(10), 104024.
<https://doi.org/10.1088/1748-9326/ac9197>
- Anthony, E. J., Almar, R., Besset, M., Reyns, J., Laibi, R., Ranasinghe, R., Abessolo Ondo, G., & Vacchi, M. (2019).
870 Response of the Bight of Benin (Gulf of Guinea, West Africa) coastline to anthropogenic and natural forcing, Part 2:
Sources and patterns of sediment supply, sediment cells, and recent shoreline change. *Continental Shelf Research*,
173, 93–103. <https://doi.org/10.1016/j.csr.2018.12.006>



- Armanuos, A. M., Al-Ansari, N., & Yaseen, Z. M. (2020). Underground Barrier Wall Evaluation for Controlling Saltwater Intrusion in Sloping Unconfined Coastal Aquifers. *Water*, *12*(9), 2403. <https://doi.org/10.3390/w12092403>
- 875 Arns, A., Dangendorf, S., Jensen, J., Talke, S., Bender, J., & Pattiaratchi, C. (2017). Sea-level rise induced amplification of coastal protection design heights. *Scientific Reports*, *7*(1), 40171. <https://doi.org/10.1038/srep40171>
- Arns, A., Wahl, T., Wolff, C., Vafeidis, A. T., Haigh, I. D., Woodworth, P., Niehüser, S., & Jensen, J. (2020). Non-linear interaction modulates global extreme sea levels, coastal flood exposure, and impacts. *Nature Communications*, *11*(1), 1918. <https://doi.org/10.1038/s41467-020-15752-5>
- 880 Athanasiou, P., van Dongeren, A., Giardino, A., Vousdoukas, M. I., Ranasinghe, R., & Kwadijk, J. (2020). Uncertainties in projections of sandy beach erosion due to sea level rise: An analysis at the European scale. *Scientific Reports*, *10*(1), 11895. <https://doi.org/10.1038/s41598-020-68576-0>
- Barnard, P. L., Erikson, L. H., Foxgrover, A. C., Hart, J. A. F., Limber, P., O'Neill, A. C., van Ormondt, M., Vitousek, S., Wood, N., Hayden, M. K., & Jones, J. M. (2019). Dynamic flood modeling essential to assess the coastal impacts of
885 climate change. *Scientific Reports*, *9*(1), 4309. <https://doi.org/10.1038/s41598-019-40742-z>
- Barnard, P. L., Short, A. D., Harley, M. D., Splinter, K. D., Vitousek, S., Turner, I. L., Allan, J., Banno, M., Bryan, K. R., Doria, A., Hansen, J. E., Kato, S., Kuriyama, Y., Randall-Goodwin, E., Ruggiero, P., Walker, I. J., & Heathfield, D. K. (2015). Coastal vulnerability across the Pacific dominated by El Niño/Southern Oscillation. *Nature Geoscience*, *8*(10), 801–807. <https://doi.org/10.1038/ngeo2539>
- 890 Bednar-Friedl, Biesbroek, R., Schmidt, D. N., Alexander, P., Børsheim, K. Y., Carnicer, J., Georgopoulou, E., Haasnoot, M., Le Cozannet, G., Lionello, P., Lipka, O., Möllmann, C., Muccione, V., Mustonen, T., Piepenburg, D., & Whitmarsh, L. (2022). Chapter 13—Europe. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. https://www.ipcc.ch/report/ar6/wg2/downloads/report/IPCC_AR6_WGII_Chapter13.pdf
- 895 Befus, K. M., Barnard, P. L., Hoover, D. J., Finzi Hart, J. A., & Voss, C. I. (2020). Increasing threat of coastal groundwater hazards from sea-level rise in California. *Nature Climate Change*, *10*(10), 946–952. <https://doi.org/10.1038/s41558-020-0874-1>



- Bellafiore, D., Ferrarin, C., Maicu, F., Manfè, G., Lorenzetti, G., Umgiesser, G., Zaggia, L., & Levinson, A. V. (2021). Saltwater Intrusion in a Mediterranean Delta Under a Changing Climate. *Journal of Geophysical Research: Oceans*, 126(2). <https://doi.org/10.1029/2020JC016437>
- 900
- Benetazzo, A., Davison, S., Barbariol, F., Mercogliano, P., Favaretto, C., & Scavo, M. (2022). Correction of ERA5 Wind for Regional Climate Projections of Sea Waves. *Water*, 14(10), 1590. <https://doi.org/10.3390/w14101590>
- Bermúdez, M., Cea, L., & Sopelana, J. (2019). Quantifying the role of individual flood drivers and their correlations in flooding of coastal river reaches. *Stochastic Environmental Research and Risk Assessment*, 33(10), 1851–1861. <https://doi.org/10.1007/s00477-019-01733-8>
- 905
- Bermúdez, M., Farfán, J. F., Willems, P., & Cea, L. (2021). Assessing the Effects of Climate Change on Compound Flooding in Coastal River Areas. *Water Resources Research*, 57(10). <https://doi.org/10.1029/2020WR029321>
- Bertels, D., & Willems, P. (2022). Climate change impact on salinization of drinking water inlets along the Campine Canals, Belgium. *Journal of Hydrology: Regional Studies*, 42, 101129. <https://doi.org/10.1016/j.ejrh.2022.101129>
- 910
- Bevacqua, E., De Michele, C., Manning, C., Couasnon, A., Ribeiro, A. F. S., Ramos, A. M., Vignotto, E., Bastos, A., Blesić, S., Durante, F., Hillier, J., Oliveira, S. C., Pinto, J. G., Ragno, E., Rivoire, P., Saunders, K., Wiel, K., Wu, W., Zhang, T., & Zscheischler, J. (2021). Guidelines for Studying Diverse Types of Compound Weather and Climate Events. *Earth's Future*, 9(11). <https://doi.org/10.1029/2021EF002340>
- Bevacqua, E., Maraun, D., Hobæk Haff, I., Widmann, M., & Vrac, M. (2017). Multivariate statistical modelling of compound events via pair-copula constructions: Analysis of floods in Ravenna (Italy). *Hydrology and Earth System Sciences*, 21(6), 2701–2723. <https://doi.org/10.5194/hess-21-2701-2017>
- 915
- Bevacqua, E., Maraun, D., Vousdoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., & Widmann, M. (2019). Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change. *Science Advances*, 5(9), eaaw5531. <https://doi.org/10.1126/sciadv.aaw5531>
- 920
- Bevacqua, E., Vousdoukas, M. I., Shepherd, T. G., & Vrac, M. (2020b). Brief communication: The role of using precipitation or river discharge data when assessing global coastal compound flooding. *Natural Hazards and Earth System Sciences*, 20(6), 1765–1782. <https://doi.org/10.5194/nhess-20-1765-2020>



- Bevacqua, E., Vousdoukas, M. I., Zappa, G., Hodges, K., Shepherd, T. G., Maraun, D., Mentaschi, L., & Feyen, L. (2020a). More meteorological events that drive compound coastal flooding are projected under climate change. *Communications Earth & Environment*, *1*(1), 47. <https://doi.org/10.1038/s43247-020-00044-z>
- 925
- Bilskie, M. V., Hagen, S. C., Medeiros, S. C., & Passeri, D. L. (2014). Dynamics of sea level rise and coastal flooding on a changing landscape: BILSKIE ET AL. *Geophysical Research Letters*, *41*(3), 927–934. <https://doi.org/10.1002/2013GL058759>
- Bonaldo, D., Bellafore, D., Ferrarin, C., Ferretti, R., Ricchi, A., Sangelantoni, L., & Vitelletti, M. L. (2023). The summer 2022 drought: A taste of future climate for the Po valley (Italy)? *Regional Environmental Change*, *23*(1), 1. <https://doi.org/10.1007/s10113-022-02004-z>
- 930
- Bonte, M., & Zwolsman, J. J. G. (2010). Climate change induced salinisation of artificial lakes in the Netherlands and consequences for drinking water production. *Water Research*, *44*(15), 4411–4424. <https://doi.org/10.1016/j.watres.2010.06.004>
- 935
- Boucharel, J., David, M., Almar, R., & Melet, A. (2023). Contrasted influence of climate modes teleconnections to the interannual variability of coastal sea level components—implications for statistical forecasts. *Climate Dynamics*. <https://doi.org/10.1007/s00382-023-06771-1>
- Bove, G., Becker, A., Sweeney, B., Vousdoukas, M., & Kulp, S. (2020). A method for regional estimation of climate change exposure of coastal infrastructure: Case of USVI and the influence of digital elevation models on assessments. *Science of The Total Environment*, *710*, 136162. <https://doi.org/10.1016/j.scitotenv.2019.136162>
- 940
- Brand, E., Ramaekers, G., & Lodder, Q. (2022). Dutch experience with sand nourishments for dynamic coastline conservation – An operational overview. *Ocean & Coastal Management*, *217*, 106008. <https://doi.org/10.1016/j.ocecoaman.2021.106008>
- Breili, K., Simpson, M. J. R., Klokkervold, E., & Roaldsdotter Ravndal, O. (2020). High-accuracy coastal flood mapping for Norway using lidar data. *Natural Hazards and Earth System Sciences*, *20*(2), 673–694. <https://doi.org/10.5194/nhess-20-673-2020>
- 945
- Bruun, P. (1962). Sea level rise as a cause of shore erosion. *Journal of Waterway, Harbors Division, ASCE* *88*(117).



- Calafat, F. M., & Marcos, M. (2020). Probabilistic reanalysis of storm surge extremes in Europe. *Proceedings of the National Academy of Sciences*, 117(4), 1877–1883. <https://doi.org/10.1073/pnas.1913049117>
- 950 Calafat, F. M., Wahl, T., Tadesse, M. G., & Sparrow, S. N. (2022). Trends in Europe storm surge extremes match the rate of sea-level rise. *Nature*, 603(7903), 841–845. <https://doi.org/10.1038/s41586-022-04426-5>
- Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., Romero, J., Aldunce, P., Barrett, K., Blanco, G., Cheung, W. W. L., Connors, S., Denton, F., Diongue-Niang, A., Dodman, D., Garschagen, M., Geden, O., Hayward, B., Jones, C., ... Péan, C. (2023). *IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. (First)*. Intergovernmental Panel on Climate Change (IPCC). <https://doi.org/10.59327/IPCC/AR6-9789291691647>
- 955
- Campbell, D., & Keddy, P. (2022). The Roles of Competition and Facilitation in Producing Zonation Along an Experimental Flooding Gradient: A Tale of Two Tails with Ten Freshwater Marsh Plants. *Wetlands*, 42(1), 5. <https://doi.org/10.1007/s13157-021-01524-4>
- 960
- Camus, P., Haigh, I. D., Nasr, A. A., Wahl, T., Darby, S. E., & Nicholls, R. J. (2021). Regional analysis of multivariate compound coastal flooding potential around Europe and environs: Sensitivity analysis and spatial patterns. *Natural Hazards and Earth System Sciences*, 21(7), 2021–2040. <https://doi.org/10.5194/nhess-21-2021-2021>
- Camus, P., Haigh, I. D., Wahl, T., Nasr, A. A., Méndez, F. J., Darby, S. E., & Nicholls, R. J. (2022). Daily synoptic conditions associated with occurrences of compound events in estuaries along North Atlantic coastlines. *International Journal of Climatology*, 42(11), 5694–5713. <https://doi.org/10.1002/joc.7556>
- 965
- Carbognin, L., & Tosi, L. (2002). Interaction between Climate Changes, Eustacy and Land Subsidence in the North Adriatic Region, Italy. *Marine Ecology*, 23(s1), 38–50. <https://doi.org/10.1111/j.1439-0485.2002.tb00006.x>
- Cardona, O. D., van Aalst, M. K., Birkmann, J., Fordham, M., McGregor, G., Perez, R., Pulwarty, R. S., Schipper, E. L. F., & Sinh, B. T. (2012). Determinants of risk: Exposure and vulnerability. In C. B. Field, V. R. Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G.-K. Plattner, S. K. Allen, M. Tignor, & P. M. Midgley (Eds.), *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special*
- 970



Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC) (Cambridge University Press, pp. 65–108).

- 975 Cazenave, A., & Le Cozannet, G. (2014). Sea level rise and its coastal impacts. *Earth's Future*, 2, 15–34.
- Chaigneau, A. A., Law-Chune, S., Melet, A., Voldoire, A., Reffray, G., & Aouf, L. (2023). Impact of sea level changes on future wave conditions along the coasts of western Europe. *Ocean Science*, 19(4), 1123–1143. <https://doi.org/10.5194/os-19-1123-2023>
- Chang, S. W., Clement, T. P., Simpson, M. J., & Lee, K.-K. (2011). Does sea-level rise have an impact on saltwater intrusion? *Advances in Water Resources*, 34(10), 1283–1291. <https://doi.org/10.1016/j.advwatres.2011.06.006>
- 980 Chaparría, V. E., Peris, J. S., & González-Escrivá, J. A. (2022). Coastal Monitoring Using Unmanned Aerial Vehicles (UAVs) for the Management of the Spanish Mediterranean Coast: The Case of Almenara-Sagunto. *International Journal of Environmental Research and Public Health*, 19(9), 5457. <https://doi.org/10.3390/ijerph19095457>
- Chesnaux, R. (2015). Closed-form analytical solutions for assessing the consequences of sea-level rise on groundwater resources in sloping coastal aquifers. *Hydrogeology Journal*, 23(7), 1399–1413. <https://doi.org/10.1007/s10040-015-1276-8>
- 985 Chesnaux, R., Marion, D., Boumaiza, L., Richard, S., & Walter, J. (2021). An analytical methodology to estimate the changes in fresh groundwater resources with sea-level rise and coastal erosion in strip-island unconfined aquifers: Illustration with Savary Island, Canada. *Hydrogeology Journal*, 29(3), 1355–1364. <https://doi.org/10.1007/s10040-020-02300-0>
- 990 Cooley, S. W., Schoeman, D., Bopp, L., Boyd, P., Donner, S., Ito, S.-I., Kiessling, W., Martinetto, P., Ojea, E., Racault, M.-F., Rost, B., Skern-Mauritzen, M., & Ghebrehiwet, D. Y. (2022). Chapter 3: Oceans and Coastal Ecosystems and their Services. In *Climate Change 2022: Impacts, Adaptation and Vulnerability. Working group II contribution to the sixth Assessment Report of the Intergovernmental Panel on Climate Change*.
- 995 Cooper, H. H. J., Kohout, F. A., Henry, H. R., & Glover, R. E. (1964). *Sea Water in Coastal Aquifers* (Geological Survey Water-Supply Paper 1613–C). United States Department of the Interior. <https://pubs.usgs.gov/wsp/1613c/report.pdf>



- Couasnon, A., Eilander, D., Muis, S., Veldkamp, T. I. E., Haigh, I. D., Wahl, T., Winsemius, H. C., & Ward, P. J. (2020). Measuring compound flood potential from river discharge and storm surge extremes at the global scale. *Natural Hazards and Earth System Sciences*, 20(2), 489–504. <https://doi.org/10.5194/nhess-20-489-2020>
- Custodio, E. (2002). Aquifer overexploitation: What does it mean? *Hydrogeology Journal*, 10(2), 254–277. <https://doi.org/10.1007/s10040-002-0188-6>
- Custodio, E., & Bruggeman, G. A. (1987). *Groundwater problems in coastal areas: A contribution to the International Hydrological Programme*. Unesco.
- D’Anna, M., Castelle, B., Idier, D., Rohmer, J., Le Cozannet, G., Thieblemont, R., & Bricheno, L. (2021). Uncertainties in Shoreline Projections to 2100 at Truc Vert Beach (France): Role of Sea-Level Rise and Equilibrium Model Assumptions. *Journal of Geophysical Research: Earth Surface*, 126(8). <https://doi.org/10.1029/2021JF006160>
- D’Anna, M., Idier, D., Castelle, B., Rohmer, J., Cagigal, L., & Mendez, F. J. (2022). Effects of stochastic wave forcing on probabilistic equilibrium shoreline response across the 21st century including sea-level rise. *Coastal Engineering*, 175, 104149. <https://doi.org/10.1016/j.coastaleng.2022.104149>
- Davenport, J., & Davenport, J. L. (2006). The impact of tourism and personal leisure transport on coastal environments: A review. *Estuarine, Coastal and Shelf Science*, 67(1–2), 280–292. <https://doi.org/10.1016/j.ecss.2005.11.026>
- Dawson, D., Shaw, J., & Roland Gehrels, W. (2016). Sea-level rise impacts on transport infrastructure: The notorious case of the coastal railway line at Dawlish, England. *Journal of Transport Geography*, 51, 97–109. <https://doi.org/10.1016/j.jtrangeo.2015.11.009>
- Dawson, R. J., Dickson, M. E., Nicholls, R. J., Hall, J. W., Walkden, M. J. A., Stansby, P. K., Mokrech, M., Richards, J., Zhou, J., Milligan, J., Jordan, A., Pearson, S., Rees, J., Bates, P. D., Koukoulas, S., & Watkinson, A. R. (2009). Integrated analysis of risks of coastal flooding and cliff erosion under scenarios of long term change. *Climatic Change*, 95(1–2), 249–288. <https://doi.org/10.1007/s10584-008-9532-8>
- De Louw, P. G. B., Eeman, S., Siemon, B., Voortman, B. R., Gunnink, J., Van Baaren, E. S., & Oude Essink, G. H. P. (2011). Shallow rainwater lenses in deltaic areas with saline seepage. *Hydrology and Earth System Sciences*, 15(12), 3659–3678. <https://doi.org/10.5194/hess-15-3659-2011>



- de Santiago, I., Camus, P., González, M., Liria, P., Epelde, I., Chust, G., del Campo, A., & Uriarte, A. (2021). Impact of climate change on beach erosion in the Basque Coast (NE Spain). *Coastal Engineering*, 167, 103916. <https://doi.org/10.1016/j.coastaleng.2021.103916>
- Delsman, J. R. (2015). *Saline groundwater—Surface water interaction in coastal lowlands*. IOS Press BV.
- 1025 Delsman, J. R., Mulder, T., Romero Verastegui, B., Bootsma, H., Zitman, P., Huizer, S., & Oude Essink, G. H. P. (2023). Reproducible construction of a high-resolution national variable-density groundwater salinity model for the Netherlands. *Environmental Modelling & Software*, 164, 105683. <https://doi.org/10.1016/j.envsoft.2023.105683>
- Delsman, J. R., Van Baaren, E. S., Siemon, B., Dabekaussen, W., Karaoulis, M. C., Pauw, P. S., Vermaas, T., Bootsma, H., De Louw, P. G. B., Gunnink, J. L., Dubelaar, C. W., Menkovic, A., Steuer, A., Meyer, U., Revil, A., & Oude Essink, 1030 G. H. P. (2018). Large-scale, probabilistic salinity mapping using airborne electromagnetics for groundwater management in Zeeland, the Netherlands. *Environmental Research Letters*, 13(8), 084011. <https://doi.org/10.1088/1748-9326/aad19e>
- Dierberg, F. E., & Kiattisimkul, W. (1996). Issues, impacts, and implications of shrimp aquaculture in Thailand. *Environmental Management*, 20(5), 649–666. <https://doi.org/10.1007/BF01204137>
- 1035 Dillon, P., Stuyfzand, P., Grischek, T., Lluria, M., Pyne, R. D. G., Jain, R. C., Bear, J., Schwarz, J., Wang, W., Fernandez, E., Stefan, C., Pettenati, M., Van Der Gun, J., Sprenger, C., Massmann, G., Scanlon, B. R., Xanke, J., Jokela, P., Zheng, Y., ... Sapiano, M. (2019). Sixty years of global progress in managed aquifer recharge. *Hydrogeology Journal*, 27(1), 1–30. <https://doi.org/10.1007/s10040-018-1841-z>
- Dodet, G., Melet, A., Arduin, F., Bertin, X., Idier, D., & Almar, R. (2019). The Contribution of Wind-Generated Waves to 1040 Coastal Sea-Level Changes. *Surveys in Geophysics*, 40(6), 1563–1601. <https://doi.org/10.1007/s10712-019-09557-5>
- Dodman, D., Hayward, B., Pelling, M., Castan Broto, V., Chow, W., Chu, E., Dawson, R., Khirfan, L., McPhearson, T., Prakash, A., Zheng, Y., & Ziervogel, G. (2022). Cities, Settlements, and Key Infrastructure. In H.-O. Pörtner, C. D. Roberts, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, & B. Rama (Eds.), *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working*



- 1045 *Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press, pp. 2233–2272). 10.1017/9781009325844.021
- Duque, C., Meyer, R., University of Copenhagen (Denmark), Sonnenborg, T. O., & Geological Survey of Denmark and Greenland (GEUS). (2022). Saltwater intrusion in Denmark. *BOLETÍN GEOLÓGICO Y MINERO*, 133(1).
<https://doi.org/10.21701/bolgeomin/133.1/002>
- 1050 Eijgenraam, C., Kind, J., Bak, C., Brekelmans, R., den Hertog, D., Duits, M., Roos, K., Vermeer, P., & Kuijken, W. (2014). Economically Efficient Standards to Protect the Netherlands Against Flooding. *Interfaces*, 44(1), 7–21.
<https://doi.org/10.1287/inte.2013.0721>
- Eilander, D., Couason, A., Ikeuchi, H., Muis, S., Yamazaki, D., Winsemius, H. C., & Ward, P. J. (2020). The effect of surge on riverine flood hazard and impact in deltas globally. *Environmental Research Letters*, 15(10), 104007.
1055 <https://doi.org/10.1088/1748-9326/ab8ca6>
- Environment Agency. (2021). *Thames Estuary 2100: 10-Year Review—Technical Monitoring Report*. Environment Agency, London, UK.
- Ericson, J., Vorosmarty, C., Dingman, S., Ward, L., & Meybeck, M. (2006). Effective sea-level rise and deltas: Causes of change and human dimension implications. *Global and Planetary Change*, 50(1–2), 63–82.
1060 <https://doi.org/10.1016/j.gloplacha.2005.07.004>
- Eslami, S., Hoekstra, P., Minderhoud, P. S. J., Trung, N. N., Hoch, J. M., Sutanudjaja, E. H., Dung, D. D., Tho, T. Q., Voepel, H. E., Woillez, M.-N., & Van Der Vegt, M. (2021). Projections of salt intrusion in a mega-delta under climatic and anthropogenic stressors. *Communications Earth & Environment*, 2(1), 142. <https://doi.org/10.1038/s43247-021-00208-5>
- 1065 European Commission. (2015). *Towards an EU research and innovation policy agenda for nature-based solutions & re-naturing cities – Final report of the Horizon 2020 expert group on “Nature-based solutions and re-naturing cities.”*
<https://data.europa.eu/doi/10.2777/479582>
- EUROSION. (2004a). *Living with coastal erosion in Europe: Sediment and Space for Sustainability. PART I - Major findings and Policy Recommendations of the EUROSION project*. Directorate General Environment. European Commission.



- 1070 EUROSION. (2004b). *Living with coastal erosion in Europe: Sediment and Space for Sustainability. PART II: Maps and Statistics. Final Report of the Project 'Coastal erosion—Evaluation of the need for action'. Directorate General Environment. European Commission.*
- Feagin, R. A., Furman, M., Salgado, K., Martinez, M. L., Innocenti, R. A., Eubanks, K., Figlus, J., Huff, T. P., Sigren, J., & Silva, R. (2019). The role of beach and sand dune vegetation in mediating wave run up erosion. *Estuarine, Coastal and Shelf Science*, 219, 97–106. <https://doi.org/10.1016/j.ecss.2019.01.018>
- 1075 Ferguson, G., & Gleeson, T. (2012). Vulnerability of coastal aquifers to groundwater use and climate change. *Nature Climate Change*, 2(5), 342–345. <https://doi.org/10.1038/nclimate1413>
- Ferrarin, C., Lionello, P., Orlić, M., Raicich, F., & Salvadori, G. (2022). Venice as a paradigm of coastal flooding under multiple compound drivers. *Scientific Reports*, 12(1), 5754. <https://doi.org/10.1038/s41598-022-09652-5>
- 1080 Ferrarin, C., Tomasin, A., Bajo, M., Petrizzo, A., & Umgiesser, G. (2015). Tidal changes in a heavily modified coastal wetland. *Continental Shelf Research*, 101, 22–33. <https://doi.org/10.1016/j.csr.2015.04.002>
- Ferreira, Ó., Kupfer, S., & Costas, S. (2021). Implications of sea-level rise for overwash enhancement at South Portugal. *Natural Hazards*, 109(3), 2221–2239. <https://doi.org/10.1007/s11069-021-04917-0>
- Gallop, S. L., Bosserelle, C., Pattiaratchi, C., & Eliot, I. (2011). Rock topography causes spatial variation in the wave, current and beach response to sea breeze activity. *Marine Geology*, 290(1–4), 29–40. <https://doi.org/10.1016/j.margeo.2011.10.002>
- 1085
- Galluccio, G., Bisaro, A., Beckhauser, E. F., Aleu, R. B., Hinkel, J., Casas, M. F., Espin, O., & Vafeidis, N. (2023). Sea Level Rise in Europe: Adaptation measures and decision-making principles. *State of the Planet, submitted.*
- Ganguli, P., & Merz, B. (2019a). Extreme Coastal Water Levels Exacerbate Fluvial Flood Hazards in Northwestern Europe. *Scientific Reports*, 9(1), 13165. <https://doi.org/10.1038/s41598-019-49822-6>
- 1090
- Ganguli, P., & Merz, B. (2019b). Trends in Compound Flooding in Northwestern Europe During 1901–2014. *Geophysical Research Letters*, 46(19), 10810–10820. <https://doi.org/10.1029/2019GL084220>
- Ganguli, P., Paprotny, D., Hasan, M., Güntner, A., & Merz, B. (2020). Projected Changes in Compound Flood Hazard From Riverine and Coastal Floods in Northwestern Europe. *Earth's Future*, 8(11). <https://doi.org/10.1029/2020EF001752>



- 1095 Genua-Olmedo, A., Alcaraz, C., Caiola, N., & Ibáñez, C. (2016). Sea level rise impacts on rice production: The Ebro Delta as an example. *Science of The Total Environment*, 571, 1200–1210. <https://doi.org/10.1016/j.scitotenv.2016.07.136>
- Gharagozlou, A., Dietrich, J. C., Karanci, A., Luettich, R. A., & Overton, M. F. (2020). Storm-driven erosion and inundation of barrier islands from dune-to region-scales. *Coastal Engineering*, 158, 103674. <https://doi.org/10.1016/j.coastaleng.2020.103674>
- 1100 Glavovic, B., Dawson, R., Chow, W., Garschagen, M., Haasnoot, M., Singh, C., & Thomas, A. (2022). Cross-Chapter Paper 2: Cities and Settlements by the Sea. In *Climate Change 2022: Impacts, Adaptation and Vulnerability. Working group II contribution to the sixth Assessment Report of the Intergovernmental Panel on Climate Change*.
- Global Commission on Adaptation. (2019). *Adapt Now: A Global Call for Leadership on Climate Resilience*. Washington, DC: World Resources Institute. <https://openknowledge.worldbank.org/handle/10986/32362> License: CC BY 4.0
- 1105 *International*. Washington, DC: World Resources Institute.
- Granger, C. W. J. (2001). Estimating the probability of flooding on a tidal river. In *Essays in econometrics: Collected Papers of Clive W. J. Granger Volume 1, Spectral Analysis, Seasonality, Nonlinearity, Methodology, and Forecasting* (pp. 355–365). Cambridge University Press.
- Grases, A., Gracia, V., García-León, M., Lin-Ye, J., & Sierra, J. P. (2020). Coastal Flooding and Erosion under a Changing
- 1110 Climate: Implications at a Low-Lying Coast (Ebro Delta). *Water*, 12(2), 346. <https://doi.org/10.3390/w12020346>
- Green, T. R., Taniguchi, M., Kooi, H., Gurdak, J. J., Allen, D. M., Hiscock, K. M., Treidel, H., & Aureli, A. (2011). Beneath the surface of global change: Impacts of climate change on groundwater. *Journal of Hydrology*, 405(3–4), 532–560. <https://doi.org/10.1016/j.jhydrol.2011.05.002>
- Grilli, A., Spaulding, M. L., Oakley, B. A., & Damon, C. (2017). Mapping the coastal risk for the next century, including sea
- 1115 level rise and changes in the coastline: Application to Charlestown RI, USA. *Natural Hazards*, 88(1), 389–414. <https://doi.org/10.1007/s11069-017-2871-x>
- Haasnoot, M., Kwadijk, J., Alphen, J. van, Bars, D. L., Diermanse, F., Spek, A. van der, Essink, G. O., Delsman, J., & Mens, M. (2020). *Adaptation to uncertain sea-level rise; how uncertainty in Antarctic mass-loss impacts the coastal adaptation strategy of the Netherlands*. <https://doi.org/10.1088/1748-9326/ab666c>



- 1120 Haigh, I. D., Eliot, M., & Pattiaratchi, C. (2011). Global influences of the 18.61 year nodal cycle and 8.85 year cycle of lunar perigee on high tidal levels. *Journal of Geophysical Research*, *116*(C6), C06025. <https://doi.org/10.1029/2010JC006645>
- Haigh, I. D., Nicholls, R. J., Penning-Rowsell, E. C., & Sayers, P. (2022). Climate change impacts on coastal flooding relevant to the UK and Ireland. *MCCIP Rolling Evidence Updates*, 18 pages. <https://doi.org/10.14465/2022.REU02.CFL>
- 1125 Haigh, I. D., Ozsoy, O., Wadey, M. P., Nicholls, R. J., Gallop, S. L., Wahl, T., & Brown, J. M. (2017). An improved database of coastal flooding in the United Kingdom from 1915 to 2016. *Scientific Data*, *4*(1), 170100. <https://doi.org/10.1038/sdata.2017.100>
- Haigh, I. D., Pickering, M. D., Green, J. A. M., Arbic, B. K., Arns, A., Dangendorf, S., Hill, D. F., Horsburgh, K., Howard, T., Idier, D., Jay, D. A., Jänicke, L., Lee, S. B., Müller, M., Schindelegger, M., Talke, S. A., Wilmes, S., & Woodworth, P. L. (2020). The Tides They Are A-Changin’: A Comprehensive Review of Past and Future Nonastronomical Changes in Tides, Their Driving Mechanisms, and Future Implications. *Reviews of Geophysics*, *58*(1). <https://doi.org/10.1029/2018RG000636>
- 1130 Haigh, I. D., Wadey, M. P., Gallop, S. L., Loehr, H., Nicholls, R. J., Horsburgh, K., Brown, J. M., & Bradshaw, E. (2015). A user-friendly database of coastal flooding in the United Kingdom from 1915–2014. *Scientific Data*, *2*(1), 150021. <https://doi.org/10.1038/sdata.2015.21>
- 1135 Hall, A. E., Herbert, R. J. H., Britton, J. R., Boyd, I. M., & George, N. C. (2019). Shelving the Coast With Vertipools: Retrofitting Artificial Rock Pools on Coastal Structures as Mitigation for Coastal Squeeze. *Frontiers in Marine Science*, *6*, 456. <https://doi.org/10.3389/fmars.2019.00456>
- Hallegatte, S., Green, C., Nicholls, R. J., & Corfee-Morlot, J. (2013). Future flood losses in major coastal cities. *Nature Climate Change*, *3*, 802–806.
- 1140 Hansen, J. E., & Barnard, P. L. (2010). Sub-weekly to interannual variability of a high-energy shoreline. *Coastal Engineering*, *57*(11–12), 959–972. <https://doi.org/10.1016/j.coastaleng.2010.05.011>



- Harley, M. D., Turner, I. L., Short, A. D., & Ranasinghe, R. (2011). Assessment and integration of conventional, RTK-GPS and image-derived beach survey methods for daily to decadal coastal monitoring. *Coastal Engineering*, 58(2), 194–
1145 205. <https://doi.org/10.1016/j.coastaleng.2010.09.006>
- He, F. J., & MacGregor, G. A. (2011). Salt reduction lowers cardiovascular risk: Meta-analysis of outcome trials. *The Lancet*, 378(9789), 380–382. [https://doi.org/10.1016/S0140-6736\(11\)61174-4](https://doi.org/10.1016/S0140-6736(11)61174-4)
- Heinrich, P., Hagemann, S., Weisse, R., Schrum, C., Daewel, U., & Gaslikova, L. (2023). Compound flood events: Analysing the joint occurrence of extreme river discharge events and storm surges in northern and central Europe. *Natural*
1150 *Hazards and Earth System Sciences*, 23(5), 1967–1985. <https://doi.org/10.5194/nhess-23-1967-2023>
- Hendry, A., Haigh, I. D., Nicholls, R. J., Winter, H., Neal, R., Wahl, T., Joly-Laugel, A., & Darby, S. E. (2019). Assessing the characteristics and drivers of compound flooding events around the UK coast. *Hydrology and Earth System Sciences*, 23(7), 3117–3139. <https://doi.org/10.5194/hess-23-3117-2019>
- Herrera-García, G., Ezquerro, P., Tomás, R., Béjar-Pizarro, M., López-Vinielles, J., Rossi, M., Mateos, R. M., Carreón-Freyre,
1155 D., Lambert, J., Teatini, P., Cabral-Cano, E., Erkens, G., Galloway, D., Hung, W.-C., Kakar, N., Sneed, M., Tosi, L., Wang, H., & Ye, S. (2021). Mapping the global threat of land subsidence. *Science*, 371(6524), 34–36. <https://doi.org/10.1126/science.abb8549>
- Hewitt, C., Mason, S., & Walland, D. (2012). The Global Framework for Climate Services. *Nature Climate Change*, 2(12), 831–832. <https://doi.org/10.1038/nclimate1745>
- 1160 Hinkel, J., Feyen, L., Hemer, M., Le Cozannet, G., Lincke, D., Marcos, M., Mentaschi, L., Merkens, J. L., de Moel, H., Muis, S., Nicholls, R. J., Vafeidis, A. T., van de Wal, R. S. W., Vousdoukas, M. I., Wahl, T., Ward, P. J., & Wolff, C. (2021). Uncertainty and Bias in Global to Regional Scale Assessments of Current and Future Coastal Flood Risk. *Earth's Future*, 9(7). <https://doi.org/10.1029/2020EF001882>
- Hinkel, J., Nicholls, R. J., Tol, R. S. J., Wang, Z. B., Hamilton, J. M., Boot, G., Vafeidis, A. T., McFadden, L., Ganopolski,
1165 A., & Klein, R. J. T. (2013). A global analysis of erosion of sandy beaches and sea-level rise: An application of DIVA. *Global and Planetary Change*, 111, 150–158. <https://doi.org/10.1016/j.gloplacha.2013.09.002>



- Holman, R. A., & Stanley, J. (2007). The history and technical capabilities of Argus. *Coastal Engineering*, 54(6–7), 477–491.
<https://doi.org/10.1016/j.coastaleng.2007.01.003>
- 1170 Horner-Devine, A. R., Hetland, R. D., & MacDonald, D. G. (2015). Mixing and Transport in Coastal River Plumes. *Annual Review of Fluid Mechanics*, 47(1), 569–594. <https://doi.org/10.1146/annurev-fluid-010313-141408>
- Horsburgh, K. J., & Wilson, C. (2007). Tide-surge interaction and its role in the distribution of surge residuals in the North Sea. *Journal of Geophysical Research: Oceans*, 112(C8), 2006JC004033. <https://doi.org/10.1029/2006JC004033>
- Hou, Y., Yang, J., Russoniello, C. J., Zheng, T., Wu, M., & Yu, X. (2022). Impacts of Coastal Shrimp Ponds on Saltwater Intrusion and Submarine Groundwater Discharge. *Water Resources Research*, 58(7).
1175 <https://doi.org/10.1029/2021WR031866>
- Hugo, G. (2011). Future demographic change and its interactions with migration and climate change. *Global Environmental Change*, 21, S21–S33. <https://doi.org/10.1016/j.gloenvcha.2011.09.008>
- Huizer, S., Oude Essink, G. H. P., & Bierkens, M. F. P. (2016). Fresh groundwater resources in a large sand replenishment. *Hydrology and Earth System Sciences*, 20(8), 3149–3166. <https://doi.org/10.5194/hess-20-3149-2016>
- 1180 Hurrell, J. W. (1995). Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation. *Science*, 269(5224), 676–679. <https://doi.org/10.1126/science.269.5224.676>
- Idier, D., Bertin, X., Thompson, P., & Pickering, M. D. (2019). Interactions Between Mean Sea Level, Tide, Surge, Waves and Flooding: Mechanisms and Contributions to Sea Level Variations at the Coast. *Surveys in Geophysics*, 40(6), 1603–1630. <https://doi.org/10.1007/s10712-019-09549-5>
- 1185 Idier, D., Paris, F., Cozannet, G. L., Boulahya, F., & Dumas, F. (2017). Sea-level rise impacts on the tides of the European Shelf. *Continental Shelf Research*, 137, 56–71. <https://doi.org/10.1016/j.csr.2017.01.007>
- Irazoqui Apecechea, M., Melet, A., & Armaroli, C. (2023). Towards a pan-European coastal flood awareness system: Skill of extreme sea-level forecasts from the Copernicus Marine Service. *Frontiers in Marine Science*, 9, 1091844. <https://doi.org/10.3389/fmars.2022.1091844>



- 1190 Jackson, L., & Devadason, C. A. (2019). *Climate Change, Flooding and Mental Health. Report from the Secretariat of the Rockefeller Foundation Economic Council on Planetary Health*. <https://www.planetaryhealth.ox.ac.uk/wp-content/uploads/sites/7/2019/04/Climate-Change-Flooding-and-Mental-Health-2019.pdf>
- Jiménez, J. A., Osorio, A., Marino-Tapia, I., Davidson, M., Medina, R., Kroon, A., Archetti, R., Ciavola, P., & Aarnikhof, S. G. J. (2007). Beach recreation planning using video-derived coastal state indicators. *Coastal Engineering*, 54(6–7), 507–521. <https://doi.org/10.1016/j.coastaleng.2007.01.012>
- 1195 Ketabchi, H., & Jahangir, M. S. (2021). Influence of aquifer heterogeneity on sea level rise-induced seawater intrusion: A probabilistic approach. *Journal of Contaminant Hydrology*, 236, 103753. <https://doi.org/10.1016/j.jconhyd.2020.103753>
- Ketabchi, H., Mahmoodzadeh, D., Ataie-Ashtiani, B., & Simmons, C. T. (2016). Sea-level rise impacts on seawater intrusion in coastal aquifers: Review and integration. *Journal of Hydrology*, 535, 235–255. <https://doi.org/10.1016/j.jhydrol.2016.01.083>
- 1200 Kew, S. F., Selten, F. M., Lenderink, G., & Hazeleger, W. (2013). The simultaneous occurrence of surge and discharge extremes for the Rhine delta. *Natural Hazards and Earth System Sciences*, 13(8), 2017–2029. <https://doi.org/10.5194/nhess-13-2017-2013>
- 1205 Khanal, S., Ridder, N., De Vries, H., Terink, W., & Van Den Hurk, B. (2018). *Storm surge and extreme river discharge: A compound event analysis using ensemble impact modelling* [Preprint]. Catchment hydrology/Modelling approaches. <https://doi.org/10.5194/hess-2018-103>
- Komar, P. D. (1998). *Beach Processes and Sedimentation* (2nd ed., Vol. 54). Prentice Hall. <http://id.erudit.org/iderudit/004810ar>
- 1210 Kothuis, B., & Kok, M. (Eds.). (2017). *Integral design of multifunctional flood defenses: Multidisciplinary approaches & examples*. Delft University Publishers.
- Kroon, A., Larson, M., Möller, I., Yokoki, H., Rozynski, G., Cox, J., & Larroude, P. (2008). Statistical analysis of coastal morphological data sets over seasonal to decadal time scales. *Coastal Engineering*, 55(7–8), 581–600. <https://doi.org/10.1016/j.coastaleng.2007.11.006>



- 1215 Kulp, S. A., & Strauss, B. H. (2018). CoastalDEM: A global coastal digital elevation model improved from SRTM using a neural network. *Remote Sensing of Environment*, 206, 231–239. <https://doi.org/10.1016/j.rse.2017.12.026>
- Larson, M., & Kraus, N. C. (1994). Temporal and spatial scales of beach profile change, Duck, North Carolina. *Marine Geology*, 117(1–4), 75–94. [https://doi.org/10.1016/0025-3227\(94\)90007-8](https://doi.org/10.1016/0025-3227(94)90007-8)
- Le Cozannet, G., Lawrence, J., Schoeman, D., Adelekan, I., Cooley, S., Glavovic, B., Haasnoot, M., Harris, R., Kiessling, W.,
1220 Kopp, R. E., Mukherji, A., Nunn, P., Piepenburg, D., Schmidt, D., Simmons, C. T., Singh, C., Slangen, A. B. A., & Supratid, S. (2022). Cross-Chapter Box SLR: Sea level rise. In *Climate Change 2022: Impacts, Adaptation and Vulnerability. Working group II contribution to the sixth Assessment Report of the Intergovernmental Panel on Climate Change*.
- Le Gal, M., Fernández-Montblanc, T., Montes Perez, J., Souto Cecon, P. E., Duo, E., Gastal, V., Delbour, S., & Ciavola, P.
1225 (2022). *A flood map catalogue for integration into a European flood awareness system (ECFAS)* [Other]. display. <https://doi.org/10.5194/egusphere-egu22-9832>
- Leaman, C. K., Harley, M. D., Splinter, K. D., Thran, M. C., Kinsela, M. A., & Turner, I. L. (2021). A storm hazard matrix combining coastal flooding and beach erosion. *Coastal Engineering*, 170, 104001. <https://doi.org/10.1016/j.coastaleng.2021.104001>
- 1230 Lee, J. J., Phillips, D. L., & Benson, V. W. (1999). Soil erosion and climate change: Assessing potential impacts and adaptation practices. *Journal of Soil and Water Conservation*, 54(3), 529–536.
- Lentz, E. E., Thieler, E. R., Plant, N. G., Stippa, S. R., Horton, R. M., & Gesch, D. B. (2016). Evaluation of dynamic coastal response to sea-level rise modifies inundation likelihood. *Nature Climate Change*, 6(7), 696–700. <https://doi.org/10.1038/nclimate2957>
- 1235 Lerczak, J. A., Geyer, W. R., & Chant, R. J. (2006). Mechanisms Driving the Time-Dependent Salt Flux in a Partially Stratified Estuary*. *Journal of Physical Oceanography*, 36(12), 2296–2311. <https://doi.org/10.1175/JPO2959.1>
- Lionello, P., Barriopedro, D., Ferrarin, C., Nicholls, R. J., Orlić, M., Raicich, F., Reale, M., Umgiesser, G., Vousdoukas, M., & Zanchettin, D. (2021). Extreme floods of Venice: Characteristics, dynamics, past and future evolution (review article). *Natural Hazards and Earth System Sciences*, 21(8), 2705–2731. <https://doi.org/10.5194/nhess-21-2705-2021>



- 1240 Liu, Q., Trinder, J., & Turner, I. L. (2017). Automatic super-resolution shoreline change monitoring using Landsat archival data: A case study at Narrabeen–Collaroy Beach, Australia. *Journal of Applied Remote Sensing*, *11*(1), 016036. <https://doi.org/10.1117/1.JRS.11.016036>
- Loáiciga, H. A. (2009). Long-term climatic change and sustainable ground water resources management. *Environmental Research Letters*, *4*(3), 035004. <https://doi.org/10.1088/1748-9326/4/3/035004>
- 1245 López-Olmedilla, L., Almeida, L. P., de Figueiredo, S. A., Fontán-Bouzas, Á., Silva, P. A., & Alcántara-Carrió, J. (2022). Effect of alongshore sediment supply gradients on projected shoreline position under sea-level rise (northwestern Portuguese coast). *Estuarine, Coastal and Shelf Science*, *271*, 107876. <https://doi.org/10.1016/j.ecss.2022.107876>
- Loureiro, C., Ferreira, Ó., & Cooper, J. A. G. (2012). Geologically constrained morphological variability and boundary effects on embayed beaches. *Marine Geology*, *329–331*, 1–15. <https://doi.org/10.1016/j.margeo.2012.09.010>
- 1250 Luijendijk, A., Hagenaars, G., Ranasinghe, R., Baart, F., Donchyts, G., & Aarninkhof, S. (2018). The State of the World’s Beaches. *Scientific Reports*, *8*(1), 6641. <https://doi.org/10.1038/s41598-018-24630-6>
- Luque, P., Gómez-Pujol, L., Marcos, M., & Orfila, A. (2021). Coastal Flooding in the Balearic Islands During the Twenty-First Century Caused by Sea-Level Rise and Extreme Events. *Frontiers in Marine Science*, *8*, 676452. <https://doi.org/10.3389/fmars.2021.676452>
- 1255 Maas, E. V., & Hoffman, G. J. (1977). Crop Salt Tolerance—Current Assessment. *Journal of the Irrigation and Drainage Division*, *103*(2), 115–134. <https://doi.org/10.1061/JRCEA4.0001137>
- Mabrouk, M., Jonoski, A., H. P. Oude Essink, G., & Uhlenbrook, S. (2018). Impacts of Sea Level Rise and Groundwater Extraction Scenarios on Fresh Groundwater Resources in the Nile Delta Governorates, Egypt. *Water*, *10*(11), 1690. <https://doi.org/10.3390/w10111690>
- 1260 Malenovský, Z., Rott, H., Cihlar, J., Schaepman, M. E., García-Santos, G., Fernandes, R., & Berger, M. (2012). Sentinels for science: Potential of Sentinel-1, -2, and -3 missions for scientific observations of ocean, cryosphere, and land. *Remote Sensing of Environment*, *120*, 91–101. <https://doi.org/10.1016/j.rse.2011.09.026>



- Mantz, P., & Wakeling, H. (1979). FORECASTING FLOOD LEVELS FOR JOINT EVENTS OF RAINFALL AND TIDAL SURGE FLOODING USING EXTREME VALUE STATISTICS. *Proceedings of the Institution of Civil Engineers*, 67(1), 31–50. <https://doi.org/10.1680/iicep.1979.2315>
- Marcos, M., Calafat, F. M., Berihuete, Á., & Dangendorf, S. (2015). Long-term variations in global sea level extremes. *Journal of Geophysical Research: Oceans*, 120(12), 8115–8134. <https://doi.org/10.1002/2015JC011173>
- Marcos, M., Wöppelmann, G., Matthews, A., Ponte, R. M., Birol, F., Arduin, F., Coco, G., Santamaría-Gómez, A., Ballu, V., Testut, L., Chambers, D., & Stopa, J. E. (2019). Coastal Sea Level and Related Fields from Existing Observing Systems. *Surveys in Geophysics*, 40(6), 1293–1317. <https://doi.org/10.1007/s10712-019-09513-3>
- Maselli, V., & Trincardi, F. (2013). Man made deltas. *Scientific Reports*, 3(1), 1926. <https://doi.org/10.1038/srep01926>
- Masselink, G., Brooks, S., Poate, T., Stokes, C., & Scott, T. (2022). Coastal dune dynamics in embayed settings with sea-level rise – Examples from the exposed and macrotidal north coast of SW England. *Marine Geology*, 450, 106853. <https://doi.org/10.1016/j.margeo.2022.106853>
- Masterson, J. P., & Garabedian, S. P. (2007). Effects of Sea-Level Rise on Ground Water Flow in a Coastal Aquifer System. *Ground Water*, 45(2), 209–217. <https://doi.org/10.1111/j.1745-6584.2006.00279.x>
- Mastrocicco, M., & Colombani, N. (2021). The Issue of Groundwater Salinization in Coastal Areas of the Mediterranean Region: A Review. *Water*, 13(1), 90. <https://doi.org/10.3390/w13010090>
- Mazas, F., & Hamm, L. (2017). An event-based approach for extreme joint probabilities of waves and sea levels. *Coastal Engineering*, 122, 44–59. <https://doi.org/10.1016/j.coastaleng.2017.02.003>
- Mazi, K., Koussis, A. D., & Destouni, G. (2013). *Intensively exploited Mediterranean aquifers: Resilience and proximity to critical points of seawater intrusion* [Preprint]. Groundwater hydrology/Modelling approaches. <https://doi.org/10.5194/hessd-10-13817-2013>
- McCall, R. T., Van Thiel De Vries, J. S. M., Plant, N. G., Van Dongeren, A. R., Roelvink, J. A., Thompson, D. M., & Reniers, A. J. H. M. (2010). Two-dimensional time dependent hurricane overwash and erosion modeling at Santa Rosa Island. *Coastal Engineering*, 57(7), 668–683. <https://doi.org/10.1016/j.coastaleng.2010.02.006>



- McGranahan, G., Balk, D., & Anderson, B. (2007). The rising tide: Assessing the risks of climate change and human settlements in low elevation coastal zones. *Environment and Urbanization*, 19(1), 17–37.
<https://doi.org/10.1177/0956247807076960>
- 1290 McMichael, C., Dasgupta, S., Ayebe-Karlsson, S., & Kelman, I. (2020). A review of estimating population exposure to sea-level rise and the relevance for migration. *Environmental Research Letters*, 15(12), 123005.
<https://doi.org/10.1088/1748-9326/abb398>
- Mel, R. A., Carniello, L., & D’Alpaos, L. (2021). How long the Mo.S.E. barriers will be effective in protecting all urban settlements within the Venice Lagoon? The wind setup constraint. *Coastal Engineering*, 168, 103923.
1295 <https://doi.org/10.1016/j.coastaleng.2021.103923>
- Melet, A., Almar, R., Hemer, M., Le Cozannet, G., Meyssignac, B., & Ruggiero, P. (2020). Contribution of Wave Setup to Projected Coastal Sea Level Changes. *Journal of Geophysical Research: Oceans*, 125(8).
<https://doi.org/10.1029/2020JC016078>
- Melet, A., Buontempo, C., Mattiuzzi, M., Salamon, P., Baharel, P., Breyiannis, G., Burgess, S., Crosnier, L., Le Traon, P.-Y.,
1300 Mentaschi, L., Nicolas, J., Solari, L., Vamborg, F., & Voukouvalas, E. (2021). European Copernicus Services to Inform on Sea-Level Rise Adaptation: Current Status and Perspectives. *Frontiers in Marine Science*, 8, 703425.
<https://doi.org/10.3389/fmars.2021.703425>
- Melet, A., Van De Wal, R., Bellafiore, D., Vousdoulas, M., Camus, P., Ferrarin, C., Oude Essink, G., Haigh, I. D., Lionello, P., Luijendijk, A., Toimil, A., & Staneva, J. (2023). Sea level rise in Europe: Observations and projections. *State of the Planet*, submitted.
1305
- Menendez, M., & Woodworth, P. L. (2010). Changes in extreme high water levels based on a quasi-global tide-gauge data set. *Journal of Geophysical Research*, 115, C10011.
- Mentaschi, L., Vousdoulas, M. I., Pekel, J.-F., Voukouvalas, E., & Feyen, L. (2018). Global long-term observations of coastal erosion and accretion. *Scientific Reports*, 8(1), 12876. <https://doi.org/10.1038/s41598-018-30904-w>



- 1310 Mentaschi, L., Vousdoukas, M. I., Voukouvalas, E., Dosio, A., & Feyen, L. (2017). Global changes of extreme coastal wave energy fluxes triggered by intensified teleconnection patterns. *Geophysical Research Letters*, *44*(5), 2416–2426. <https://doi.org/10.1002/2016GL072488>
- Michael, H. A., Russoniello, C. J., & Byron, L. A. (2013). Global assessment of vulnerability to sea-level rise in topography-limited and recharge-limited coastal groundwater systems: GLOBAL ASSESSMENT OF VULNERABILITY TO
1315 SEA-LEVEL RISE. *Water Resources Research*, *49*(4), 2228–2240. <https://doi.org/10.1002/wrcr.20213>
- Milliman, J. D. (1997). Blessed dams or damned dams? *Nature*, *386*(6623), 325–327. <https://doi.org/10.1038/386325a0>
- Minderhoud, P. S. J., Erkens, G., Pham, V. H., Bui, V. T., Erban, L., Kooi, H., & Stouthamer, E. (2017). Impacts of 25 years of groundwater extraction on subsidence in the Mekong delta, Vietnam. *Environmental Research Letters*, *12*(6), 064006. <https://doi.org/10.1088/1748-9326/aa7146>
- 1320 Moftakhari, H. R., AghaKouchak, A., Sanders, B. F., Allaire, M., & Matthew, R. A. (2018). What Is Nuisance Flooding? Defining and Monitoring an Emerging Challenge. *Water Resources Research*, *54*(7), 4218–4227. <https://doi.org/10.1029/2018WR022828>
- Moftakhari, H. R., Salvadori, G., AghaKouchak, A., Sanders, B. F., & Matthew, R. A. (2017). Compounding effects of sea level rise and fluvial flooding. *Proceedings of the National Academy of Sciences*, *114*(37), 9785–9790.
1325 <https://doi.org/10.1073/pnas.1620325114>
- Monioudi, I. N., Velegrakis, A. F., Chatzipavlis, A. E., Rigos, A., Karambas, T., Vousdoukas, M. I., Hasiotis, T., Koukourouvli, N., Peduzzi, P., Manoutsoglou, E., Poulos, S. E., & Collins, M. B. (2017). Assessment of island beach erosion due to sea level rise: The case of the Aegean archipelago (Eastern Mediterranean). *Natural Hazards and Earth System Sciences*, *17*(3), 449–466. <https://doi.org/10.5194/nhess-17-449-2017>
- 1330 Mooyaart, L. F., & Jonkman, S. N. (2017). Overview and Design Considerations of Storm Surge Barriers. *Journal of Waterway, Port, Coastal, and Ocean Engineering*, *143*(4), 06017001. [https://doi.org/10.1061/\(ASCE\)WW.1943-5460.0000383](https://doi.org/10.1061/(ASCE)WW.1943-5460.0000383)
- Morim, J., Hemer, M., Wang, X. L., Cartwright, N., Trenham, C., Semedo, A., Young, I., Bricheno, L., Camus, P., Casas-Prat, M., Erikson, L., Mentaschi, L., Mori, N., Shimura, T., Timmermans, B., Aarnes, O., Breivik, O., Behrens, A.,



- 1335 Dobrynin, M., ... Andutta, F. (2019). Robustness and uncertainties in global multivariate wind-wave climate projections. *Nature Climate Change*, 9(9), 711–718. <https://doi.org/10.1038/s41558-019-0542-5>
- Muis, S., Apecechea, M. I., Dullaart, J., de Lima Rego, J., Madsen, K. S., Su, J., Yan, K., & Verlaan, M. (2020). A High-Resolution Global Dataset of Extreme Sea Levels, Tides, and Storm Surges, Including Future Projections. *Frontiers in Marine Science*, 7, 263. <https://doi.org/10.3389/fmars.2020.00263>
- 1340 Murray, A. B., Lazarus, E., Ashton, A., Baas, A., Coco, G., Coulthard, T., Fonstad, M., Haff, P., McNamara, D., Paola, C., Pelletier, J., & Reinhardt, L. (2009). Geomorphology, complexity, and the emerging science of the Earth's surface. *Geomorphology*, 103(3), 496–505. <https://doi.org/10.1016/j.geomorph.2008.08.013>
- Navoy, A. S. (1991). *Aquifer-estuary interaction and vulnerability of groundwater supplies to sea level rise-driven saltwater intrusion*. Pennsylvania State University.
- 1345 Neumann, B., Vafeidis, A. T., Zimmermann, J., & Nicholls, R. J. (2015a). Future Coastal Population Growth and Exposure to Sea-Level Rise and Coastal Flooding—A Global Assessment. *PLOS ONE*, 10(3), e0118571. <https://doi.org/10.1371/journal.pone.0118571>
- Neumann, B., Vafeidis, A. T., Zimmermann, J., & Nicholls, R. J. (2015b). Future Coastal Population Growth and Exposure to Sea-Level Rise and Coastal Flooding—A Global Assessment. *PLOS ONE*, 10(3), e0118571. <https://doi.org/10.1371/journal.pone.0118571>
- 1350 Nicholls, R. J., & Cazenave, A. (2010). Sea-Level Rise and Its Impact on Coastal Zones. *Science*, 328(5985), 1517–1520. <https://doi.org/10.1126/science.1185782>
- Nicholls, R., Wong, P., Burkett, V., Codignotto, J., Hay, J., McLean, R., & Woodroffe, S. R. C. (2007). Coastal systems and low-lying areas. In *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 315–356). Cambridge University Press.
- 1355 Noble, R. M. (1978). Coastal Structures' Effects on Shorelines. *Coastal Engineering 1978*, 2068–2085. <https://doi.org/10.1061/9780872621909.127>



- 1360 Oppenheimer, M., Glavovic, B., Hinkel, J., van de Wal, R. S. W., Magnan, A., Abd-Elgawad, A., Cai, R., Cifuentes-Jara, M.,
DeConto, R. M., Ghosh, T., Hay, J., Isla, F. I., Marzeion, B., Meyssignac, B., & Sebesvari, Z. (2019). Sea Level Rise
and Implications for Low Lying Islands, Coasts and Communities. In *IPCC Special Report on the Ocean and
Cryosphere in a Changing Climate* (Pörtner H.-O. et al.). Cambridge University Press.
- Orton, P., Ralston, D., van Prooijen, B., Secor, D., Ganju, N., Chen, Z., Fernald, S., Brooks, B., & Marcell, K. (2023). Increased
Utilization of Storm Surge Barriers: A Research Agenda on Estuary Impacts. *Earth's Future*, *11*(3), e2022EF002991.
1365 <https://doi.org/10.1029/2022EF002991>
- Oude Essink, G. H. P. (1996). *Impact of sea level rise on groundwater flow regimes: A sensitivity analysis for the Netherlands*.
Delft University Press.
- Oude Essink, G. H. P. (2001). Salt Water Intrusion in a Three-dimensional Groundwater System in The Netherlands: A
Numerical Study. *Transport in Porous Media*, *43*(1), 137–158. <https://doi.org/10.1023/A:1010625913251>
- 1370 Oude Essink, G. H. P., Van Baaren, E. S., & De Louw, P. G. B. (2010). Effects of climate change on coastal groundwater
systems: A modeling study in the Netherlands. *Water Resources Research*, *46*(10), 2009WR008719.
<https://doi.org/10.1029/2009WR008719>
- Paprotny, D., Morales-Nápoles, O., & Jonkman, S. N. (2018). HANZE: A pan-European database of exposure to natural
hazards and damaging historical floods since 1870. *Earth System Science Data*, *10*(1), 565–581.
1375 <https://doi.org/10.5194/essd-10-565-2018>
- Paprotny, D., Morales-Nápoles, O., Vousdoukas, M. I., Jonkman, S. N., & Nikulin, G. (2019). Accuracy of pan-European
coastal flood mapping. *Journal of Flood Risk Management*, *12*(2), e12459. <https://doi.org/10.1111/jfr3.12459>
- Paprotny, D., Terefenko, P., Giza, A., Czapliński, P., & Vousdoukas, M. I. (2021). Future losses of ecosystem services due to
coastal erosion in Europe. *Science of The Total Environment*, *760*, 144310.
1380 <https://doi.org/10.1016/j.scitotenv.2020.144310>
- Paprotny, D., Vousdoukas, M. I., Morales-Nápoles, O., Jonkman, S. N., & Feyen, L. (2020). Pan-European hydrodynamic
models and their ability to identify compound floods. *Natural Hazards*, *101*(3), 933–957.
<https://doi.org/10.1007/s11069-020-03902-3>



- Passeri, D. L., Bilskie, M. V., Plant, N. G., Long, J. W., & Hagen, S. C. (2018). Dynamic modeling of barrier island response to hurricane storm surge under future sea level rise. *Climatic Change*, 149(3), 413–425. <https://doi.org/10.1007/s10584-018-2245-8>
- Passeri, D. L., Hagen, S. C., Bilskie, M. V., & Medeiros, S. C. (2015). On the significance of incorporating shoreline changes for evaluating coastal hydrodynamics under sea level rise scenarios. *Natural Hazards*, 75(2), 1599–1617. <https://doi.org/10.1007/s11069-014-1386-y>
- 1385
- Passeri, D. L., Hagen, S. C., Medeiros, S. C., Bilskie, M. V., Alizad, K., & Wang, D. (2015). The dynamic effects of sea level rise on low-gradient coastal landscapes: A review: THE DYNAMIC EFFECTS OF SEA LEVEL RISE ON LOW-GRADIENT COASTAL LANDSCAPES. *Earth's Future*, 3(6), 159–181. <https://doi.org/10.1002/2015EF000298>
- 1390
- Passeri, D. L., Hagen, S. C., Plant, N. G., Bilskie, M. V., Medeiros, S. C., & Alizad, K. (2016). Tidal hydrodynamics under future sea level rise and coastal morphology in the Northern Gulf of Mexico. *Earth's Future*, 4(5), 159–176. <https://doi.org/10.1002/2015EF000332>
- 1395
- Pereira, H., Sousa, M. C., Vieira, L. R., Morgado, F., & Dias, J. M. (2022). Modelling Salt Intrusion and Estuarine Plumes under Climate Change Scenarios in Two Transitional Ecosystems from the NW Atlantic Coast. *Journal of Marine Science and Engineering*, 10(2), 262. <https://doi.org/10.3390/jmse10020262>
- Petroliagkis, T. I., Joint Research Centre (European Commission), Bidlot, J., Voukouvalas, E., & Disperati, J. (2016). *Joint probabilities of storm surge, significant wave height and river discharge components of coastal flooding events: Utilising statistical dependence methodologies & techniques*. Publications Office of the European Union. <https://data.europa.eu/doi/10.2788/677778>
- 1400
- Pickering, M. D., Wells, N. C., Horsburgh, K. J., & Green, J. A. M. (2012). The impact of future sea-level rise on the European Shelf tides. *Continental Shelf Research*, 35, 1–15. <https://doi.org/10.1016/j.csr.2011.11.011>
- 1405
- Pietro, L. S., O'Neal, M. A., & Puleo, J. A. (2008). Developing Terrestrial-LIDAR-Based Digital Elevation Models for Monitoring Beach Nourishment Performance. *Journal of Coastal Research*, 246, 1555–1564. <https://doi.org/10.2112/07-0904.1>



- Pollard, J., Spencer, T., & Brooks, S. (2019). The interactive relationship between coastal erosion and flood risk. *Progress in Physical Geography: Earth and Environment*, 43(4), 574–585. <https://doi.org/10.1177/0309133318794498>
- 1410 Pörtner, H.-O., Roberts, C. D., Poloczanska, E., Mintenbeck, K., Tignor, M., Alegria, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., & Okem, A. (Eds.). (2022). IPCC Summary for Policymakers. In *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 3–33). Cambridge University Press. <https://dx.doi.org/10.1017/9781009325844.001>
- 1415 Post, V. E. A., Oude Essink, G., Szymkiewicz, A., Bakker, M., Houben, G., Custodio, E., & Voss, C. (2018). Celebrating 50 years of SWIMs (Salt Water Intrusion Meetings). *Hydrogeology Journal*, 26(6), 1767–1770. <https://doi.org/10.1007/s10040-018-1800-8>
- Pugh, D., & Woodworth, P. (2014). *Sea-Level Science: Understanding Tides, Surges, Tsunamis and Mean Sea-Level Changes* (1st ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9781139235778>
- 1420 Qi, H., Cai, F., Lei, G., Cao, H., & Shi, F. (2010). The response of three main beach types to tropical storms in South China. *Marine Geology*, 275(1–4), 244–254. <https://doi.org/10.1016/j.margeo.2010.06.005>
- Rasmussen, P., Sonnenborg, T. O., Goncear, G., & Hinsby, K. (2013). Assessing impacts of climate change, sea level rise, and drainage canals on saltwater intrusion to coastal aquifer. *Hydrology and Earth System Sciences*, 17(1), 421–443. <https://doi.org/10.5194/hess-17-421-2013>
- 1425 Rodrigues, M., Fortunato, A. B., & Freire, P. (2019). Saltwater Intrusion in the Upper Tagus Estuary during Droughts. *Geosciences*, 9(9), 400. <https://doi.org/10.3390/geosciences9090400>
- Roest, B., de Vries, S., de Schipper, M., & Aarninkhof, S. (2021). Observed Changes of a Mega Feeder Nourishment in a Coastal Cell: Five Years of Sand Engine Morphodynamics. *Journal of Marine Science and Engineering*, 9(1), 37. <https://doi.org/10.3390/jmse9010037>
- 1430 Romagnoli, C., Bosman, A., Casalbore, D., Anzidei, M., Doumaz, F., Bonaventura, F., Meli, M., & Verdirame, C. (2022). Coastal Erosion and Flooding Threaten Low-Lying Coastal Tracts at Lipari (Aeolian Islands, Italy). *Remote Sensing*, 14(13), 2960. <https://doi.org/10.3390/rs14132960>



- Rueda, A., Camus, P., Tomás, A., Vitousek, S., & Méndez, F. J. (2016). A multivariate extreme wave and storm surge climate emulator based on weather patterns. *Ocean Modelling*, *104*, 242–251. <https://doi.org/10.1016/j.ocemod.2016.06.008>
- 1435 Ruocco, A. C., Nicholls, R. J., Haigh, I. D., & Wadey, M. P. (2011). Reconstructing coastal flood occurrence combining sea level and media sources: A case study of the Solent, UK since 1935. *Natural Hazards*, *59*(3), 1773–1796. <https://doi.org/10.1007/s11069-011-9868-7>
- Sallenger, A. H. Jr. (2000). Storm Impact Scale for Barrier Islands. *Journal of Coastal Research*, *16*(3), 890–895.
- Samuels, P. G., & Burt, N. (2002). A new joint probability appraisal of flood risk. *Proceedings of the Institution of Civil Engineers - Water and Maritime Engineering*, *154*(2), 109–115. <https://doi.org/10.1680/wame.2002.154.2.109>
- 1440 Sanò, M., Jiménez, J. A., Medina, R., Stanica, A., Sanchez-Arcilla, A., & Trumbic, I. (2011). The role of coastal setbacks in the context of coastal erosion and climate change. *Ocean & Coastal Management*, *54*(12), 943–950. <https://doi.org/10.1016/j.ocecoaman.2011.06.008>
- Santos, V. M., Casas-Prat, M., Poschlod, B., Ragno, E., Van Den Hurk, B., Hao, Z., Kalmár, T., Zhu, L., & Najafí, H. (2021). Statistical modelling and climate variability of compound surge and precipitation events in a managed water system: A case study in the Netherlands. *Hydrology and Earth System Sciences*, *25*(6), 3595–3615. <https://doi.org/10.5194/hess-25-3595-2021>
- 1445 Sanuy, M., & Jiménez, J. A. (2021). Probabilistic characterisation of coastal storm-induced risks using Bayesian networks. *Natural Hazards and Earth System Sciences*, *21*(1), 219–238. <https://doi.org/10.5194/nhess-21-219-2021>
- 1450 Sayers, P. B., Hall, J. W., & Meadowcroft, I. C. (2002). Towards risk-based flood hazard management in the UK. *Proceedings of the Institution of Civil Engineers - Civil Engineering*, *150*(5), 36–42. <https://doi.org/10.1680/cien.2002.150.5.36>
- Sayers, P. B., Horritt, M. S., Penning-Rowsell, E., & McKenzie, A. (2015). *Climate Change Risk Assessment 2017: Projections of future flood risk in the UK. Research undertaken by Sayers and Partners on behalf of the Committee on Climate Change*. Committee on Climate Change, London. <https://www.theccc.org.uk/wp-content/uploads/2015/10/CCRA-Future-Flooding-Main-Report-Final-06Oct2015.pdf>
- 1455 Schmork, S., & Mercado, A. (1969). Upconing of Fresh Water-Sea Water Interface Below Pumping Wells, Field Study. *Water Resources Research*, *5*(6), 1290–1311. <https://doi.org/10.1029/WR005i006p01290>



- Scussolini, P., Aerts, J. C. J. H., Jongman, B., Bouwer, L. M., Winsemius, H. C., de Moel, H., & Ward, P. J. (2016). FLOPROS: An evolving global database of flood protection standards. *Natural Hazards and Earth System Sciences*, 16(5), 1049–1061. <https://doi.org/10.5194/nhess-16-1049-2016>
- 1460
- Sherif, M. M., & Singh, V. P. (1999). Effect of climate change on sea water intrusion in coastal aquifers. *Hydrological Processes*, 13(8), 1277–1287. [https://doi.org/10.1002/\(SICI\)1099-1085\(19990615\)13:8<1277::AID-HYP765>3.0.CO;2-W](https://doi.org/10.1002/(SICI)1099-1085(19990615)13:8<1277::AID-HYP765>3.0.CO;2-W)
- Short, A. D. (Ed.). (1999). *Handbook of beach and shoreface morphodynamics*. John Wiley.
- 1465 Short, A. D., & Trembanis, A. C. (2004). Decadal Scale Patterns in Beach Oscillation and Rotation Narrabeen Beach, Australia—Time Series, PCA and Wavelet Analysis. *Journal of Coastal Research*, 20(2), 523–532. [https://doi.org/10.2112/1551-5036\(2004\)020\[0523:DSPIBO\]2.0.CO;2](https://doi.org/10.2112/1551-5036(2004)020[0523:DSPIBO]2.0.CO;2)
- Simon, B., Costabel, S., Voß, W., Meyer, U., Deus, N., Elbracht, J., Günther, T., & Wiederhold, H. (2015). Airborne and ground geophysical mapping of coastal clays in Eastern Friesland, Germany. *GEOPHYSICS*, 80(3), WB21–WB34.
- 1470 <https://doi.org/10.1190/geo2014-0102.1>
- Sierra, J. P., & Casas-Prat, M. (2014). Analysis of potential impacts on coastal areas due to changes in wave conditions. *Climatic Change*, 124(4), 861–876. <https://doi.org/10.1007/s10584-014-1120-5>
- Small, C., & Nicholls, R. J. (2003). A global analysis of human settlement in coastal zones. *Journal of Coastal Research*, 19(3), 584–599.
- 1475 Sprenger, C., Hartog, N., Hernández, M., Vilanova, E., Grützmacher, G., Scheibler, F., & Hannappel, S. (2017). Inventory of managed aquifer recharge sites in Europe: Historical development, current situation and perspectives. *Hydrogeology Journal*, 25(6), 1909–1922. <https://doi.org/10.1007/s10040-017-1554-8>
- Stevens, A. J., Clarke, D., & Nicholls, R. J. (2016). Trends in reported flooding in the UK: 1884–2013. *Hydrological Sciences Journal*, 61(1), 50–63. <https://doi.org/10.1080/02626667.2014.950581>
- 1480 Stofberg, S. F., Klimkowska, A., Paulissen, M. P. C. P., Witte, J.-P. M., & van der Zee, S. E. A. T. M. (2015). Effects of salinity on growth of plant species from terrestrializing fens. *Aquatic Botany*, 121, 83–90. <https://doi.org/10.1016/j.aquabot.2014.12.004>



- Stringari, C. E., & Power, H. (2022). Picoastal: A low-cost coastal video monitoring system. *SoftwareX*, 18, 101073.
<https://doi.org/10.1016/j.softx.2022.101073>
- 1485 Stripling, S., Panzeri, M., Blanco, B., Rossington, K., Sayers, P., & Borthwick, A. (2017). Regional-scale probabilistic shoreline evolution modelling for flood-risk assessment. *Coastal Engineering*, 121, 129–144.
<https://doi.org/10.1016/j.coastaleng.2016.12.002>
- Svensson, C., & Jones, D. A. (2002). Dependence between extreme sea surge, river flow and precipitation in eastern Britain. *International Journal of Climatology*, 22(10), 1149–1168. <https://doi.org/10.1002/joc.794>
- 1490 Svensson, C., & Jones, D. A. (2004). Dependence between sea surge, river flow and precipitation in south and west Britain. *Hydrology and Earth System Sciences*, 8(5), 973–992. <https://doi.org/10.5194/hess-8-973-2004>
- Taylor, R. G., Scanlon, B., Döll, P., Rodell, M., van Beek, R., Wada, Y., Longuevergne, L., Leblanc, M., Famiglietti, J. S., Edmunds, M., Konikow, L., Green, T. R., Chen, J., Taniguchi, M., Bierkens, M. F. P., MacDonald, A., Fan, Y., Maxwell, R. M., Yechieli, Y., ... Treidel, H. (2013). Ground water and climate change. *Nature Climate Change*, 3(4),
1495 322–329. <https://doi.org/10.1038/nclimate1744>
- Thompson, G., & Law, F. (1983). An assessment of the fluvial tidal flooding problem of the River Ancholme, UK. *Proceedings from the IUGG Interdisciplinary Symposium on Assessment of Natural Hazards*. IUGG Interdisciplinary Symposium on Assessment of Natural Hazards, Hamburg.
- Timmerman, A., Haasnoot, M., Middelkoop, H., Bouma, T., & McEvoy, S. (2021). Ecological consequences of sea level rise and flood protection strategies in shallow coastal systems: A quick-scan barcoding approach. *Ocean & Coastal Management*, 210, 105674. <https://doi.org/10.1016/j.ocecoaman.2021.105674>
- 1500 Toimil, A., Álvarez-Cuesta, M., & Losada, I. J. (2023). Neglecting the effect of long- and short-term erosion can lead to spurious coastal flood risk projections and maladaptation. *Coastal Engineering*, 179, 104248.
<https://doi.org/10.1016/j.coastaleng.2022.104248>
- 1505 Toimil, A., Losada, I. J., Álvarez-Cuesta, M., & Le Cozannet, G. (2023). Demonstrating the value of beaches for adaptation to future coastal flood risk. *Nature Communications*, 14(1), 3474. <https://doi.org/10.1038/s41467-023-39168-z>



- Toimil, A., Losada, I. J., Camus, P., & Díaz-Simal, P. (2017). Managing coastal erosion under climate change at the regional scale. *Coastal Engineering*, *128*, 106–122. <https://doi.org/10.1016/j.coastaleng.2017.08.004>
- Toimil, A., Losada, I. J., Díaz-Simal, P., Izaguirre, C., & Camus, P. (2017). Multi-sectoral, high-resolution assessment of climate change consequences of coastal flooding. *Climatic Change*, *145*(3–4), 431–444. <https://doi.org/10.1007/s10584-017-2104-z>
- Toimil, A., Losada, I. J., Nicholls, R. J., Dalrymple, R. A., & Stive, M. J. F. (2020). Addressing the challenges of climate change risks and adaptation in coastal areas: A review. *Coastal Engineering*, *156*, 103611. <https://doi.org/10.1016/j.coastaleng.2019.103611>
- 1515 Valle-Levinson, A. (2010). Definition and classification of estuaries. In A. Valle-Levinson (Ed.), *Contemporary Issues in Estuarine Physics* (1st ed., pp. 1–11). Cambridge University Press. <https://doi.org/10.1017/CBO9780511676567.002>
- Van Den Hurk, B., Van Meijgaard, E., De Valk, P., Van Heeringen, K.-J., & Gooijer, J. (2015a). Analysis of a compounding surge and precipitation event in the Netherlands. *Environmental Research Letters*, *10*(3), 035001. <https://doi.org/10.1088/1748-9326/10/3/035001>
- 1520 Van Den Hurk, B., Van Meijgaard, E., De Valk, P., Van Heeringen, K.-J., & Gooijer, J. (2015b). Analysis of a compounding surge and precipitation event in the Netherlands. *Environmental Research Letters*, *10*(3), 035001. <https://doi.org/10.1088/1748-9326/10/3/035001>
- van Engelen, J., Essink, G. H. P. O., & Bierkens, M. F. P. (2022). Sustainability of fresh groundwater resources in fifteen major deltas around the world. *Environmental Research Letters*, *17*(12), 125001. <https://doi.org/10.1088/1748-9326/aca16c>
- 1525
- Van Ormondt, M., Nelson, T. R., Hapke, C. J., & Roelvink, D. (2020). Morphodynamic modelling of the wilderness breach, Fire Island, New York. Part I: Model set-up and validation. *Coastal Engineering*, *157*, 103621. <https://doi.org/10.1016/j.coastaleng.2019.103621>
- van Weert, F., van der Gun, J., & Reckman, J. (2009). *Global Overview of Saline Groundwater Occurrence and Genesis* (GP 2009-1; p. 107). International Groundwater Resources Assessment Centre. <https://www.un->
- 1530



igrac.org/sites/default/files/resources/files/Global%20Overview%20of%20Saline%20Groundwater%20Occurrences
%20and%20Genesis.pdf

- Vandenbohede, A., Luyten, K., & Lebbe, L. (2008). Effects of Global Change on Heterogeneous Coastal Aquifers: A Case Study in Belgium. *Journal of Coastal Research*, 2, 160–170. <https://doi.org/10.2112/05-0447.1>
- 1535 Vousdoukas, M. I. (2012). Erosion/accretion patterns and multiple beach cusp systems on a meso-tidal, steeply-sloping beach. *Geomorphology*, 141–142, 34–46. <https://doi.org/10.1016/j.geomorph.2011.12.003>
- Vousdoukas, M. I., Almeida, L. P. M., & Ferreira, Ó. (2012). Beach erosion and recovery during consecutive storms at a steep-sloping, meso-tidal beach: BEACH EROSION/RECOVERY DURING CONSECUTIVE STORMS. *Earth Surface Processes and Landforms*, 37(6), 583–593. <https://doi.org/10.1002/esp.2264>
- 1540 Vousdoukas, M. I., Kirupakaramoorthy, T., Oumeraci, H., de la Torre, M., Wübbold, F., Wagner, B., & Schimmels, S. (2014). The role of combined laser scanning and video techniques in monitoring wave-by-wave swash zone processes. *Coastal Engineering*, 83, 150–165. <https://doi.org/10.1016/j.coastaleng.2013.10.013>
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Bianchi, A., Dottori, F., & Feyen, L. (2018a). Climatic and socioeconomic controls of future coastal flood risk in Europe. *Nature Climate Change*, 8(9), 776–780. <https://doi.org/10.1038/s41558-018-0260-4>
- 1545
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., & Feyen, L. (2017). Extreme sea levels on the rise along Europe’s coasts: EXTREME SEA LEVELS ALONG EUROPE’S COASTS. *Earth’s Future*, 5(3), 304–323. <https://doi.org/10.1002/2016EF000505>
- Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., Jevrejeva, S., Jackson, L. P., & Feyen, L. (2018b). Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard. *Nature Communications*, 9(1), 2360. <https://doi.org/10.1038/s41467-018-04692-w>
- 1550
- Vousdoukas, M. I., Pennucci, G., Holman, R. A., & Conley, D. C. (2011). A semi automatic technique for Rapid Environmental Assessment in the coastal zone using Small Unmanned Aerial Vehicles (SUAV). *Journal of Coastal Research, Special Issue 64: PROCEEDINGS OF THE 11th INTERNATIONAL COASTAL SYMPOSIUM ICS2011*(64), 1755–
1555 1759.



- Vousdoukas, M. I., Ranasinghe, R., Mentaschi, L., Plomaritis, T. A., Athanasiou, P., Luijendijk, A., & Feyen, L. (2020). Sandy coastlines under threat of erosion. *Nature Climate Change*, *10*(3), 260–263. <https://doi.org/10.1038/s41558-020-0697-0>
- Vousdoukas, M. I., Velegrakis, A. F., & Plomaritis, T. A. (2007). Beachrock occurrence, characteristics, formation mechanisms and impacts. *Earth-Science Reviews*, *85*(1–2), 23–46. <https://doi.org/10.1016/j.earscirev.2007.07.002>
- Vousdoukas, M. I., Voukouvalas, E., Annunziato, A., Giardino, A., & Feyen, L. (2016). Projections of extreme storm surge levels along Europe. *Climate Dynamics*, *47*(9–10), 3171–3190. <https://doi.org/10.1007/s00382-016-3019-5>
- Wada, Y., Van Beek, L. P. H., Van Kempen, C. M., Reckman, J. W. T. M., Vasak, S., & Bierkens, M. F. P. (2010). Global depletion of groundwater resources. *Geophysical Research Letters*, *37*(20), n/a-n/a. <https://doi.org/10.1029/2010GL044571>
- Wada, Y., Wisser, D., & Bierkens, M. F. P. (2014). Global modeling of withdrawal, allocation and consumptive use of surface water and groundwater resources. *Earth System Dynamics*, *5*(1), 15–40. <https://doi.org/10.5194/esd-5-15-2014>
- Wadey, M. P., Haigh, Ivan. D., Nicholls, R. J., Brown, J. M., Horsburgh, K., Carroll, B., Gallop, S. L., Mason, T., & Bradshaw, E. (2015). A comparison of the 31 January–1 February 1953 and 5–6 December 2013 coastal flood events around the UK. *Frontiers in Marine Science*, *2*. <https://doi.org/10.3389/fmars.2015.00084>
- Wahl, T., Jain, S., Bender, J., Meyers, S. D., & Luther, M. E. (2015). Increasing risk of compound flooding from storm surge and rainfall for major US cities. *Nature Climate Change*, *5*(12), 1093–1097. <https://doi.org/10.1038/nclimate2736>
- Ward, P. J., Couasnon, A., Eilander, D., Haigh, I. D., Hendry, A., Muis, S., Veldkamp, T. I. E., Winsemius, H. C., & Wahl, T. (2018). Dependence between high sea-level and high river discharge increases flood hazard in global deltas and estuaries. *Environmental Research Letters*, *13*(8), 084012. <https://doi.org/10.1088/1748-9326/aad400>
- Werner, A. D., & Simmons, C. T. (2009). Impact of Sea-Level Rise on Sea Water Intrusion in Coastal Aquifers. *Ground Water*, *47*(2), 197–204. <https://doi.org/10.1111/j.1745-6584.2008.00535.x>
- Werner, B. T. (1999). Complexity in Natural Landform Patterns. *Science*, *284*(5411), 102–104. <https://doi.org/10.1126/science.284.5411.102>



- 1580 White, C. J. (2007). *The use of joint probability analysis to predict flood frequency in estuaries and tidal rivers* [Phd, University of Southampton]. <https://eprints.soton.ac.uk/63847/>
- White, E., & Kaplan, D. (2017). Restore or retreat? Saltwater intrusion and water management in coastal wetlands. *Ecosystem Health and Sustainability*, 3(1), e01258. <https://doi.org/10.1002/ehs2.1258>
- Wolff, C., Bonatz, H., & Vafeidis, A. T. (2023). Setback zones can effectively reduce exposure to sea-level rise in Europe. 1585 *Scientific Reports*, 13(1), 5515. <https://doi.org/10.1038/s41598-023-32059-9>
- Wu, W., McInnes, K., O’Grady, J., Hoeke, R., Leonard, M., & Westra, S. (2018). Mapping Dependence Between Extreme Rainfall and Storm Surge. *Journal of Geophysical Research: Oceans*, 123(4), 2461–2474. <https://doi.org/10.1002/2017JC013472>
- Yamazaki, D., Ikeshima, D., Tawatari, R., Yamaguchi, T., O’Loughlin, F., Neal, J. C., Sampson, C. C., Kanae, S., & Bates, P. 1590 D. (2017). A high-accuracy map of global terrain elevations: Accurate Global Terrain Elevation map. *Geophysical Research Letters*, 44(11), 5844–5853. <https://doi.org/10.1002/2017GL072874>
- Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., Van Den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., & Vignotto, E. (2020). A 1595 typology of compound weather and climate events. *Nature Reviews Earth & Environment*, 1(7), 333–347. <https://doi.org/10.1038/s43017-020-0060-z>
- Zscheischler, J., Westra, S., Van Den Hurk, B. J. J. M., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., Wahl, T., & Zhang, X. (2018). Future climate risk from compound events. *Nature Climate Change*, 8(6), 469–477. <https://doi.org/10.1038/s41558-018-0156-3>

1600

1605