

# Laboratory experiments in ocean alkalinity enhancement research

M. Débora Iglesias-Rodríguez<sup>1,2</sup>, Rosalind E.M. Rickaby<sup>3</sup>, Arvind Singh<sup>4</sup>, James Gatley<sup>1,2</sup>

<sup>1</sup> Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara; Santa Barbara, CA 93106, U.S.A.

<sup>2</sup> Marine Science Institute, University of California, Santa Barbara, CA 93106, U.S.A.

<sup>3</sup> Department of Earth Sciences, University of Oxford, Oxford, U.K.

<sup>4</sup> Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, India.

*Correspondence to:* M. Débora Iglesias-Rodríguez, [iglesias@ucsb.edu](mailto:iglesias@ucsb.edu)

**Abstract.** Recent concern about the consequences of continuing increases in atmospheric CO<sub>2</sub> as a key heat-trapping agent (USGCRP, 2017; IPCC, 2021) have prompted ocean experts to come together to discuss how to provide science-based solutions. Ocean alkalinity enhancement (OAE) is being considered not only as an ocean carbon dioxide removal (CDR) approach, but also as a potential way to mitigate ocean acidification. Over the last two decades, inter-laboratory comparisons have proven valuable in evaluating the reliability of methodologies associated with sampling and analysis of carbonate chemistry parameters, which have been routinely used in ocean acidification research. Given the complexity of processes and mechanisms related to ecosystem responses to OAE, consolidating protocols to ensure compatibility across studies is fundamental for synthesis and upscaling analysis. This chapter provides an overview of best practice in OAE laboratory experimentation and facilitates awareness of the importance of applying standardized methods to promote data re-use, inter-lab comparisons, meta-analysis and transparency. This chapter provides the reader with the tools to (1) identify the criteria to achieve the best laboratory practice and experimental design; (2) provide guidance on the selection of response variables for various purposes (physiological, biogeochemical, ecological, evolutionary) for inter-lab comparisons; (3) offer recommendation for a minimum set of variables that should be sampled and propose additional variables critical for different types of synthesis and upscaling; and (4) identify protocols for standardized measurements of response variables. Key recommendations include ensuring reproducibility through appropriate experimental design and replication, assessing alkalinity thresholds for secondary precipitates for each experimental approach and condition, using recommended targets of alkalinity (3000-4000 μmol kg<sup>-1</sup>) and levels exceeding these concentrations to mimic responses at the site of deployment/non equilibrium and use intermediate alkalinity levels to identify potential nonlinear responses, and establish the appropriate experimental design to address questions at specific levels of organization (chemical, physiological, molecular) and assuming different scenarios (e.g., mimicking impacts at the site of deployment in a non-equilibrated system versus steady state scenarios in an equilibrated system).

## 1. Introduction

Laboratory studies need to be reproducible, consistent and transparent (Box 1) to provide the scientific community and regulators with useful information to move the field forward and facilitate the development of safe guidelines. Based on numerous modeling studies, ocean alkalinity enhancement (OAE) appears to be a promising ocean carbon dioxide removal (CDR) approach, with the likely beneficial side effect of mitigating ocean acidification (Burt et al., 2021; Hartmann et al., 2023; NASEM, 2022; Wang et al., 2023). Laboratory experiments are urgently needed to determine the CDR potential of various OAE methods as well as OAE impacts at various levels of biological organization (ecological, physiological, biochemical, molecular). The emerging empirical studies are offering insight while revealing gaps in our knowledge of the mechanisms governing OAE and its effect on marine biota (e.g., Ferderer et al., 2022; Gatley et al., 2023; Yang et al., 2023). For example, the conditions preventing or limiting the formation of secondary precipitates and the pros and cons of various alkali are still under debate. Given that empirical work on OAE is still in its infancy and that some of the assumptions based on modeling studies remain untested, this chapter is an evolving document that will be updated as the OAE community continues to release results.

54  
55 Laboratory manipulations allow making observations in a highly controlled environment using model  
56 species or subsets of populations (selected species or populations). Results are generally considered highly  
57 reproducible (Box 1) and therefore laboratory manipulations are viewed as a necessary step to either generate  
58 hypotheses to test in the field or vice versa, when field experimentation is an option. Under the latter, field  
59 observations guide the laboratory experiments to validate field results in well-known systems and under tightly  
60 controlled conditions.

61  
62 A number of approaches – batch, semi-continuous and continuous cultures – have been used to address  
63 diverse OAE settings (e.g., at the point of deployment, under steady state conditions, air- versus non air-equilibrated  
64 seawater) and various biological scenarios (specific stages of growth, life cycle, and to explore abrupt/short term  
65 versus long term responses to manipulations). In some cases, specific stages during the life cycle of organisms can  
66 be selected (for example, larval versus adult stage; sexual versus asexual phase). Time series laboratory experiments  
67 are less restricted than mesocosm experiments with regards to the duration of experiments because they tend to be  
68 ‘cleaner’, with relatively low bacteria numbers and generally without biological confounding factors (viruses,  
69 predation, competition for resources, etc.). Therefore, the cause–effect relationships are easier to elucidate as  
70 conditions and organisms can be tested in relative isolation, and there is the possibility of extensive replication.

71  
72 The main limitation of laboratory experiments is that the dynamic phenomena occurring in the natural  
73 environment cannot be captured in the laboratory and, therefore, results may not be applicable to real life scenarios.  
74 For example, in laboratory experiments the influence of mixing processes, conditions governing particle flocculation  
75 or the linkage to higher levels of biological organization (e.g., predation) are difficult to discern (see Forbes and  
76 Calow, 2002; Martin et al., 2014). Portable lab experiments, such as deck incubations aboard research vessels or  
77 outdoor incubations, with some influence from the local environment (e.g., diurnal alterations of light, water flow  
78 through from the coast to maintain in-situ temperature) as well as community-level mesocosm experiments are the  
79 conduit to field manipulations. These large-scale community experimental tanks address the importance of the  
80 physico-chemical conditions, space, density-dependent effects, biotic interactions and complexity of natural  
81 environments in their response to OAE manipulations\buffering, or boosting, the direct effects of environmental  
82 stress on organisms (Paiva et al., 2021).

83  
84 This chapter provides best practice guidelines in OAE laboratory experimentation and offers  
85 recommendations to enable data re-use, inter-lab comparisons, and transparency. We offer recommendations  
86 regarding (1) the criteria to achieve the best laboratory practice and experimental design; (2) the selection of  
87 response variables for various purposes (physiological, biogeochemical, ecological, evolutionary) for inter-lab  
88 comparisons; (3) a minimum set of variables that should be sampled and additional variables critical for different  
89 types of synthesis and upscaling; and (4) protocols for standardized measurements of response variables.

## 90 91 92 **2. Lessons learned from ocean acidification research**

93 The rich insights obtained in ocean acidification research are key to supporting OAE studies. However, as  
94 crucial as it is to follow guidelines when designing laboratory experiments, it is equally important to acknowledge  
95 that there may be potential confounders and challenges that may not be accounted for in the guidelines. Being able  
96 to conduct quantitative laboratory intercomparisons, including interspecies comparisons, will be critically dependent  
97 on identifying recommendations regarding experimental design, sample collection and data analysis. Important  
98 considerations include the source of alkalinity, rate of alkalinity addition, testing air-CO<sub>2</sub>-equilibrated *versus* non-  
99 equilibrated seawater, and the effect of ancillary variables (e.g., temperature) in multifactorial experiments which  
100 are known to yield complex and variable results (e.g., see the interactive effects of ocean acidification and warming  
101 - Harvey et al., 2013). The guidelines provided in this chapter should significantly improve the quality and impact of  
102 the OAE research, which is required to meet the identified societal need for research on OAE and other types of  
103 ocean CDR (NASEM, 2021).

104  
105 An exploration of procedures, patterns and challenges associated with ocean acidification research has  
106 offered ideas on how to design rigorous and reproducible laboratory experiments that enable measuring and  
107 monitoring carbonate chemistry shifts and biological responses to ocean acidification (Cornwall and Hurd, 2016).  
108 Cornwall and Hurd (2016) reported that 95% of the experimental work between 1993 and 2014 had interdependent  
109 or lacked replication in clearly defined treatments, or did not report sufficient methodological detail. More broadly,

110 results from Wernberg et al. (2012) from marine climate change experiments between 2000 and 2009, reported that  
111 ~49% of the experiments had identifiable issues with their experimental procedures, and 91% of the experiments  
112 reported showed a lack of treatment replication or pseudo-replication. Amongst the studies, 9% included  
113 extreme/unrealistic treatments of temperature or pH far beyond worst case scenario projections (Wernberg et al.,  
114 2012) although ‘extreme’ pH/alkalinity conditions may prove useful to define thresholds of tolerance and upper  
115 limits of alkalinity enhancement, and to understand underlying physiological mechanisms of acclimation to  
116 alkalinization. While the urgent need for field trials requires careful consideration of treatment levels, in order to  
117 maximize the insight gained from OAE experiments, testing conditions outside the year 2100 IPCC CO<sub>2</sub> emission  
118 scenarios are encouraged. These conditions outside worst case scenario projections will further our knowledge on  
119 the mechanisms governing biological (e.g., shell production) and abiotic (e.g., particle aggregation, secondary  
120 precipitation) responses to applied chemical CDR.  
121

122 Like in ocean acidification research, careful attention should be given to the advantages and disadvantages  
123 that concern the choices of dissolved inorganic carbon species to measure, and how error propagation will affect the  
124 calculated parameters (Martz et al., 2015). Moreover, dissolved organic matter (DOM) is known to contribute to  
125 alkalinity (Kim and Lee, 2009; Koeve et al., 2010) although the presence of strong acidic groups in organic matter  
126 can decrease net alkalinity (Hu, 2020; Middelburg et al., 2020). Depending on the type of system under  
127 investigation, attention should be paid to whether to apply titration alkalinity (typically used in ocean studies) versus  
128 the charge balance approach (often used in freshwater systems, with high concentrations of dissolved organic  
129 matter) (see Middlebufig et al., 2020). Results from ocean acidification mesocosm experiments focused on  
130 phytoplankton revealed that nutrient-limited communities appeared to be more responsive to changing carbonate  
131 chemistry than those having access to high inorganic nutrient concentrations (see Paul et al., 2015; Sala et al., 2015;  
132 Bach et al., 2016). These observations indicate that trophic state might play a role in the susceptibility of organisms  
133 to the changes in carbonate chemistry driven by alkalinization. Also, competition between species has been found to  
134 be altered under various carbonate chemistry conditions (see Kroeker et al., 2013a), which merits a focus on  
135 experiments that address preferential selection of taxonomic groups under different alkalinity conditions. Although  
136 applying nutrient-limiting conditions is experimentally challenging, understanding how species succession and  
137 community composition might respond to alkalinization could in part be addressed in a laboratory context.  
138

139 While it is fairly straightforward to determine how individual changes in parameters influence chemical  
140 and biological responses, understanding impacts of multiple parameters [e.g., increased alkalinity and warming,  
141 increased alkalinity and resource availability (nutrients, light, prey)] can be challenging as they can interact in  
142 complex ways. Indeed, ocean acidification research revealed antagonistic, synergistic, and additive responses when  
143 studying ocean acidification and warming (Byrne and Przeslawski, 2013; Kroeker et al., 2013b; Harvey et al., 2015;  
144 Pistevo et al., 2016). Identifying tipping points and interactive effects when other parameters (e.g., temperature) are  
145 altered in seawater, in addition to alkalinity, is critical given the capacity of these parameters to drive (otherwise  
146 unpredictable) shifts in species abundances, biodiversity and community composition, physiological outputs,  
147 survival, and reproduction (Crain et al., 2008; Darling and Côté, 2008; Galic et al., 2018).

#### 148 **Box 1. Criteria for best laboratory practice**

149 · **Reproducibility.** From the emerging OAE research (e.g., regarding the formation of secondary  
150 precipitates - see Montserrat et al., 2017 versus Fuhr et al., 2022; and Moras et al., 2022) and the ocean  
151 acidification literature (e.g., see Ridgwell et al., 2009), we have learned that similar approaches can  
152 lead to conflicting and unresolved outcomes. Without appropriate reporting of sample collection,  
153 methodology and data processing, it is challenging to re-analyze the data and reconcile the  
154 discrepancies. As the field emerges and evolves, it will be required to reevaluate early experiments and  
155 possibly re-analyze results with updated protocols.

156 · **Defining inclusion and exclusion criteria.** In order to reduce confounding covariates, attention  
157 must be paid to factors affecting flocculation, aggregation of particles (e.g., possibly impacted by  
158 dissolved organic matter increases after phytoplankton blooms), fluctuations in temperature, which  
159 affect mineral dissolution and precipitation rates, and biological and physiological properties,  
160 including stage during the life cycle, trophic state, and seasonality, that affect the susceptibility of  
161 organisms to OAE (e.g., see Vandamme et al., 2015; Subhas et al., 2022).

162 · **Establishing experimental controls.** In OAE experimental designs, controls must be appropriately  
163 selected. These could include seawater without added alkalinity, seawater ± nutrients/food, treatments

164 with and without the organisms tested. When mineral dissolution is too slow, an alternative analog that  
165 reproduces the basic chemistry is encouraged (for example, the use of salts and alkali; e.g.,  $\text{CaCl}_2$  and  
166  $\text{Na}_2\text{CO}_3$  to mimic the effect of limestone-based mineral dissolution). Controls could also contain an  
167 alternative form of alkalinity that alters the seawater carbonate chemistry solely, without adding carbon  
168 or trace metals (e.g.,  $\text{NaOH}$ ).

169 · **Basic biological responses.** Studies on organisms' physiological responses (e.g., growth,  
170 respiration, size, reproduction, photosynthesis and calcification) are recommended. These responses  
171 can be measured directly; for example, as uptake rates of solutes using traditional assays, mass  
172 spectrometric methods for indirect assessment of changes in elements, or molecular responses using  
173 markers of functional processes. Rates of growth and calcification can also be measured by changes in  
174 dry mass or buoyant mass in many types of organisms, especially in macroinvertebrates and  
175 macroalgae (see Dodge et al., 1984; Davis, 1989; Sanders et al., 2018). For organisms that undergo  
176 development one must determine which stage of development (e.g., larval vs adult; vegetative vs  
177 gamete stage) to target. Also, when altering more than one parameter, particular attention must be paid  
178 to potential confounding effects. Multi-factorial experiments can be used to explore the weight of each  
179 parameter.

### 181 3. Seawater media preparation and manipulation of carbonate chemistry

182 The different steps in experimental design are outlined in Table 1. The process starts with natural or  
183 artificially made seawater with or without nutrient additions. One must consider whether adding nutrients/food/prey  
184 is required; for example, whether exploring OAE impacts is intended in conjunction with specific scenarios, e.g.,  
185 nutrient fertilization, specific stages of growth or population development, and the extent to which nutrient additions  
186 or any other basic manipulation of the environmental conditions might impact the interpretation of results. For OAE  
187 manipulations where sterilization is required for the experimental set up, autoclaving is discouraged given the  
188 alterations in carbonate chemistry, including loss of  $\text{CO}_2$ , leading to a decrease in dissolved inorganic carbon and  
189 alterations in alkalinity (increase with increasing salinity/decrease with precipitation of carbonate) triggered by  
190 autoclaving. Instead, filter-sterilization of seawater through small pore size filters (e.g.,  $0.22\ \mu\text{m}$  filters) is required  
191 to remove particles and most bacteria, and produce the stock media where different manipulations are applied to  
192 create different alkalinity treatments.

194 There are several approaches to simulating the addition of alkalinity that capture different components of  
195 any manipulation experiment. The first approach could be testing the impact of instantaneous addition of alkalinity  
196 to seawater to mimic the impact on seawater chemistry and ecosystems at the point of deployment. The second  
197 involves aeration and equilibration with the atmosphere to explore the physico-chemical response to a steady state/  
198 equilibrated scenario. In the latter instance, the medium is aliquoted out to the experimental vessels/tanks where  
199 aeration is applied to promote air equilibration. Monitoring carbonate chemistry through time enables determining  
200 when equilibration of seawater with air occurs.

202 **Table 1. Experimental considerations for OAE experimentation. Medium preparation:** the seawater can be  
203 obtained from coastal or open ocean sites. Filtered seawater or, when appropriate (e.g., when growing autotrophic  
204 organisms), seawater supplemented with nutrients; for example, using f/2 or variations of f/2 media (see Guillard  
205 and Ryder, 1962) will be used for growing organisms. Seawater media can also be prepared from artificial recipes  
206 (e.g., Aquil medium; Morel et al., 1979) when specific compounds or elements need to be altered in seawater. Media  
207 must be sterilized by filtration rather than through autoclaving and nutrients can be added, typically from stock  
208 solutions. When possible, moderate aeration should be applied. Types of alkali include adding pulverized mineral  
209 directly to the media and promote dissolution physically (e.g., by stirring); dissolving the mineral separately and  
210 filter out any particles remaining in the media before experimentation; dissolving salts to mimic the chemistry of the  
211 dissolved alkali (e.g., to mimic limestone dissolution, dissolve  $\text{CaCl}_2$  and  $\text{NaCO}_3$ , which result in higher dissolution  
212 rates); and adding liquid alkali such as  $\text{NaOH}$ . Establishing time series prior to the experiment to determine time  
213 frames regarding length of experiment, frequency of sampling, etc. is recommended. **Experimental design:** in  
214 addition to optimizing reproducibility by designing enough replication and test the reproducibility of the method,  
215 researchers should remain engaged with respect to protocols and experimental design to avoid artifacts and  
216 undesirable side effects of methodology. When possible, ensure equilibration of seawater gasses with air and define  
217 experimental time frames to test impacts under conditions representative of the site of deployment (where limited  
218 gas exchange occurs) and those representative of steady state/equilibrated conditions. Although most laboratory  
219 experiments address short term impacts, chronic effects can be tested in long term incubations. **Sampling and**

220 **analysis:** the parameters to be considered should allow inter-lab comparisons, address functional properties of  
 221 organisms (e.g., calcification, silicification, particulate organic carbon) and fulfill needs to improve model  
 222 parameterizations. It is important to establish well defined time windows for sampling as well as frequency of  
 223 sampling to capture physical, chemical and biological properties of the studied system. It is advisable to limit the  
 224 time of sample storage to minimize observations that might confound interpretation of results (e.g., reverse  
 225 weathering during storage). Stock solutions (e.g., nutrient and alkalinity solutions) must be stored in the appropriate  
 226 vessels to avoid contamination from leachates coming out of the vessel itself (e.g., silicate contamination from  
 227 solutions stored in borosilicate containers). Detection limits and accuracy and precision should be offered for each  
 228 protocol.  
 229

Medium preparation	Experiment design	Sampling and analysis
<p><b>Natural/artificial seawater</b></p> <p><b>Filter sterilization (e.g., 0.22 um)</b></p> <p>+/- nutrient addition</p> <p>+/- aeration</p> <p><b>Type of alkalinity treatments</b></p> <ul style="list-style-type: none"> <li>• Pulverized mineral</li> <li>• Pre-dissolved mineral</li> <li>• Dissolved salts</li> <li>• Liquid alkali</li> </ul> <p><b>Pre-equilibrated vs non-equilibrated seawater with air phase</b></p> <ul style="list-style-type: none"> <li>• Carbonate chemistry</li> <li>• Flocculation/aggregation</li> <li>• Biology</li> </ul>	<p><b>Best actions to maximize confidence</b></p> <ul style="list-style-type: none"> <li>• Within study replication and pseudo-replication</li> <li>• Coordinated networks (teams sharing progress to decide on best protocols)</li> </ul> <p><b>Preliminary time series of TA and carbonate chemistry</b></p> <ul style="list-style-type: none"> <li>• Define experimental time frames</li> <li>• Assess TA upper limits</li> <li>• Expand the upper limits to address impacts at site of deployment</li> </ul> <p><b>Abrupt vs chronic biology impacts</b></p> <ul style="list-style-type: none"> <li>• Short-term tests (acclimation)</li> <li>• Long-term experiments (adaptation)</li> </ul>	<p><b>Criteria for key parameters</b></p> <ul style="list-style-type: none"> <li>• Inter-lab comparisons</li> <li>• Functional properties</li> <li>• Model parameterization</li> </ul> <p><b>Sampling frequency and timing</b></p> <ul style="list-style-type: none"> <li>• Select time window for sampling</li> <li>• Identify sampling frequency that captures key chemical, physical or biological features</li> </ul> <p><b>Limit storage to minimize artifacts</b></p> <p><b>Identify and report key analytical parameters affecting error</b></p> <ul style="list-style-type: none"> <li>• Detection limits</li> <li>• Measurement accuracy/precision</li> <li>• Identify any impact of experimental design on uncertainties</li> </ul>

230  
 231  
 232 **3.a Sources of alkalinity**

233 As yet, it is unclear what the optimal method or source of alkalinity enhancement may be in order to  
 234 simulate the desired chemistry in seawater media. Proposed sources of alkalinity include silicate minerals (olivine,  
 235 basalt), brucite, limestone and its derivatives (quicklime and portlandite), NaOH and mine tailings (NASEM, 2021;  
 236 Nawaz et al., 2023). Given the slow dissolution kinetics of the minerals, generating alkaline solutions artificially is  
 237 acceptable. For example, Gately et al. (2023) simulated alkalinity enhancement via a limestone-inspired solution by  
 238 adding Na<sub>2</sub>CO<sub>3</sub> and CaCl<sub>2</sub> or its hydrated form (CaCl<sub>2</sub>H<sub>4</sub>O<sub>2</sub>) to seawater. Adding Na<sub>2</sub>CO<sub>3</sub> raises TA and DIC in a  
 239 2:1 ratio, with 2 moles of TA added by 2 conservative Na<sup>+</sup> ions in Na<sub>2</sub>CO<sub>3</sub>, and 1 mole DIC added by CO<sub>3</sub><sup>2-</sup>. CaCl<sub>2</sub>  
 240 does not raise alkalinity because it adds equal amounts of positive and negative conservative charge to the solution  
 241 from Ca<sup>2+</sup> and 2 x Cl<sup>-</sup>. However, it does raise the calcium in solution and therefore the saturation state of the  
 242 seawater with respect to CaCO<sub>3</sub>.  
 243

244 Many possibilities for solid or liquid alkalinity additions are being considered (see chapter 3). While adding  
 245 minerals as precursors of alkalinity can provide a source of potentially beneficial nutrients (e.g., silicate, iron,  
 246 magnesium) (Hartmann et al. 2013), the possible toxic effect of metals leached out of minerals, an example being  
 247 nickel (Ni) leached from olivine (Montserrat et al., 2017) is of concern. The use of NaOH is currently gaining  
 248 attention given that its environmental footprint is perceived as smaller than the mining of alkaline minerals, which  
 249 necessitate an expansion of mining operations, transportation, and industrial processing, which are energetically  
 250 costly and can lead to air pollution. Additionally, the amount of Na added to seawater is very small relative to the  
 251 large background of NaCl in seawater.  
 252

253 The addition of NaOH and other forms of alkalinity to seawater cause initial spikes in pH and a drop in  
 254 aqueous CO<sub>2</sub> that can be balanced to a steady state via bubbling with air (Table 1). Determining abiotic and biotic

255 responses to the initial spikes in pH and drops in CO<sub>2</sub> is an important step in addition to understanding responses  
256 under steady-state conditions. It may be that large manipulations of alkalinity are needed to elicit a measurable and  
257 reproducible response, and the required alkalinity concentrations will be refined with more detailed modeling but,  
258 based on current information, proposed targets for alkalinity manipulations are 3000-4000 μmol kg<sup>-1</sup> (Renforth and  
259 Henderson, 2017). ~4000 μmol/kg is the concentration of alkalinity expected at locations in the ocean where  
260 alkalinity is initially added, and ~3000 μmol/kg is the concentration of alkalinity expected once ocean circulation  
261 has dispersed the alkalinity over a larger area (Renforth and Henderson, 2017). Alkalinity thresholds for the  
262 formation of precipitates will need to be determined for each experimental approach and condition. It is however  
263 recommended that researchers consider using alkalinities exceeding the recommended targets, and utilize  
264 intermediate treatments (e.g., 2000, 4000, 7000 μatm/kg seawater) rather than just low/high treatments, in order to  
265 identify potential nonlinear and even parabolic responses. This approach led to important and unexpected outcomes  
266 in ocean acidification research (e.g., Ries et al., 2009).

### 267 **3.b Impacts of impurities/metal leachates**

268 An important consideration in OAE studies is the impact of metals leached from dissolving minerals and  
269 their ecotoxicological potential on marine organisms. For example, although some elements (e.g., Fe and Mg)  
270 leached out of minerals could be beneficial micronutrients, the potentially toxic effect of metals such as nickel (Ni)  
271 (Montserrat et al., 2017), leached from olivine, is of concern. Diverse responses have however been reported with  
272 respect to Ni and it appears that some cyanobacteria rely on Ni more than other photosynthetic organisms (see  
273 Dupont et al., 2008, 2010; Ho, 2013). A recent laboratory study testing olivine leachates (containing Si, Ni, Mg, Fe,  
274 Cr and Co) in phytoplankton revealed either positive or neutral physiological short term responses in all treatments  
275 (Hutchins et al., 2023). However, one should consider the role of long-term experiments to examine organismal and  
276 population adaptation of metal exposure as well as potential bioaccumulation and biomagnification impacts in  
277 consumers.  
278

279  
280 Another important consideration is the effect of pH on metal speciation as pH and a change in the  
281 concentration of OH<sup>-</sup> and CO<sub>3</sub><sup>2-</sup> ions can affect the solubility, adsorption, toxicity, and rates of redox processes of  
282 metals in seawater thus altering the interactions of metals with marine organisms (Millero et al., 2009). When  
283 dissolving minerals in seawater one must consider nonstoichiometry and incomplete dissolution perhaps as a result  
284 of dissolution of impurities, precipitation of secondary minerals, or preferential leaching of elements from the  
285 mineral surface (Brantley, 2008, NASEM, 2021). The formation of secondary precipitates has been observed in  
286 several studies exploring the dissolution of olivine (Fuhr et al., 2022), and limestone derivatives (Moras et al., 2022;  
287 Gately et al., 2023; Hartmann et al., 2023). Using an alkaline solution rather than reactive alkaline particles has been  
288 recommended to reduce carbonate precipitation unless seawater critical supersaturation levels are exceeded  
289 (Hartmann et al., 2023). In addition, runaway CaCO<sub>3</sub> precipitation, a condition where more alkalinity is removed  
290 than initially added, reduces the OAE CO<sub>2</sub> uptake efficiency. More complex precipitates containing Fe, Si, and P  
291 were observed in a study using a limestone-inspired OAE approach revealing that mineral precipitation caused by  
292 seawater alkalization can also remove inorganic nutrients from solution (Gately et al., 2023).  
293


294 Maintaining alkalinity following OAE is critically dependent on the carbonate saturation state, its temporal  
295 evolution, and particle surface processes (Hartmann et al., 2023). To minimize the loss of alkalinity and maximize  
296 alkalinity enhancement, Hartmann et al (2023) propose the application of an alkaline solution in CO<sub>2</sub> equilibrium  
297 with the atmosphere and/or solutions with tested saturation levels to prevent a further increase in supersaturation,  
298 and the precipitation of carbonate to avoid loss of alkalinity. A separate reservoir where alkaline solutions have  
299 been prepared is desirable for testing upper limits of alkalinity addition and identifying saturation thresholds to  
300 minimize precipitation.

## 301 **4. Experimental design**

### 302 **4.a Experimental replication**

303  
304 Replication is important to determine if results are reproducible although one must consider that when results are so  
305 dependent on precise experimental conditions that replicability is needed for reproducibility, the result may be  
306 unique and potentially less relevant than a phenomenon that can be reproduced by a variety of independent, non-  
307 identical approaches (see Casadevall and Fang, 2010). A number of experimental designs can be used to achieve  
308 adequate statistical replication (Fig. 1). For example, simple replication involves experimental units (each of the  
309 replicates) per treatment where all the conditions are manipulated independently but in the same way for that  
310

311 treatment and where responses to the treatment are measured [defined by Hurlbert (2009) as the “evaluation unit”]  
 312 and each experimental unit can be considered as independent. In temporal replication, multiple measurements are  
 313 made through time (temporal trends) on the same experimental unit. Sacrificial replication involves the use of  
 314 multiple sampling times per treatment (for example, a time series) and multiple experimental units at the time of  
 315 samplin. Each approach has distinct strengths and limitations, and the choice of the approach depends on the  
 316 scientific questions and the extent of the risk of error propagation. For example, one might choose sacrificial  
 317 replication for certain chemical manipulations that require sampling from vessels with comparable volumes but  
 318 choose instead temporal replication for monitoring the evolution of a microbial culture or the physiology of fish over  
 319 time under certain alkalinity conditions.  
 320  
 321  
 322



323  
 324  
 325 **Figure 1.** Examples of experimental laboratory design with regards to replication. Each treatment, represented by a  
 326 colour, contains experimental units ~~contains~~ (replicates). Each experimental unit is treated as an independent  
 327 experiment except in the sacrificial replication approach, where each replicate is treated statistically as an  
 328 experimental unit.  
 329

#### 330 4.b Preliminary experiments

331 In addition to testing the biological responses to abrupt enhanced alkalinity, marine organisms can be  
 332 exposed to enhanced alkalinity conditions after equilibration of seawater pCO<sub>2</sub> with that in the air-phase following  
 333 alkalinity addition. Ideally, aeration should be maintained to ensure O<sub>2</sub> levels required by marine animals and also  
 334 maintain stable pCO<sub>2</sub> levels in the alkalinity perturbation experiments. Depending on the organism tested (a few  
 335 organisms do not tolerate aeration in tanks), aeration might or might not remain for the duration of the experiment  
 336 (Table 1). The vessels used in OAE experiments might not be traditional tanks used in aquaria, but rather any type  
 337 of container adequate for different type of organisms (e.g., culture flasks for bacteria, conical flasks, carboys for  
 338 phytoplankton, open tanks for echinoderms and fish) with air lines to introduce aeration in the media. When running  
 339 multifactorial experiments (e.g., temperature and alkalinity), designing an analysis plan and concrete experimental  
 340 questions to interrogate can help determine the sample size and minimum number of treatments.  
 341

342 An analog to OAE is the use of lime soda and other alkali to combat acid rain, which has caused deleterious  
 343 changes in freshwater ecosystems for more than half a century in northern Europe and North America. To reverse  
 344 some of these' changes a number of governmental and nongovernmental teams have applied lime and other  
 345 neutralizing compounds to streams, rivers, lakes, and catchments in the most affected or most ecologically valuable  
 346 regions (see Clair and Hindar, 2005). Another example is the effects of seawater buffering mainly by addition of  
 347 Na<sub>2</sub>CO<sub>3</sub> addition) utilized by the commercial shellfish industry (e.g., Ragg et al., 2019), which showed a broad  
 348 improvement in larval health compared to undersaturated waters.  
 349  
 350

351 Standardizing technical details in protocols, sampling, sample processing and analyses are crucial to control  
352 for variation introduced by reagents, sample storage and other factors. The collection and curation of metadata  
353 associated with each sample are critical for data interpretation, inter-lab comparison and drawing conclusions to  
354 move forward with planning field deployments for research purposes. For studies involving more than one level of  
355 biological organization; i.e., grazing experiments, competition experiments, particular attention should be paid to  
356 designing adequate controls.

357  
358 The effects of OAE and its interactions with other parameters might differ depending on the duration of the  
359 experiments. Indeed, in ocean acidification studies, compensatory metabolic pathways appear to take longer to  
360 become established, depending on factors such as the exposure history (Calosi et al., 2013) and phase of the life  
361 cycle (Hettinger et al., 2012). In a study testing ocean acidification and warming, biological effects were not  
362 detectable in the short term, but were rather manifested over time (Godbold and Solan, 2013). It was suggested that  
363 species responses to seasonal variations in environmental conditions might explain these differences that, depending  
364 upon timing, can either exacerbate or buffer the long-term directional effects of climatic forcing (Godbold and  
365 Solan, 2013).

#### 366 **4.c Recommended minimum set of variables to report**

367 To improve comparability between future work, we recommend a minimum set of variables with the  
368 understanding that more variables might be added as new results emerge (Table 2). We recommend to measure and  
369 report at least the following variables (shown in bold in Table 2).

- 371 • At least alkalinity and one more parameter of the carbonate system must be measured to calculate key carbonate  
372 chemistry parameters including bicarbonate and carbonate ions, CO<sub>2</sub>, pH and the saturation state of CaCO<sub>3</sub>  
373 polymorphs. This information is critical to determine chemical alterations in the dissolved inorganic carbon  
374 system as a result of alkalization.
- 375 • Resource availability (e.g., prey, dissolved inorganic nutrients, light) are needed to monitor the growth  
376 conditions.
- 377 • Particulate organic carbon (POC), nitrogen (PON), phosphorous (POP) are required to learn about trends in  
378 biomass production and stoichiometry.
- 379 • Basic physiological properties (respiration, photosynthesis) should be measured to inform biogeochemical  
380 models and learn about biologically-mediated fluxes of elements.
- 381 • Some functional group-specific properties, particularly those involving mineral precipitation (calcification,  
382 silicification) and those with environmental effects (e.g., toxin production) and with climate-relevant impacts  
383 (nitrogen fixation/denitrification) in context specific cases.
- 384 • Size of offspring and fecundity rates can be used as indicators of transgenerational plasticity and adaptation to  
385 alkalization.

386  
387 Other variables are important in the exploration of specific questions such as how does seawater alkalization  
388 affect biodiversity?; how does metal bioavailability change under increased pH?; what is the role of organic  
389 alkalinity in coastal systems? The variables and protocols listed in this chapter is not exhaustive and only provides a  
390 proxy sample largely based on the literature on climate impacts on marine systems and ocean acidification.

391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404



405  
406  
407  
408  
409  
410

**Table 2. Examples of responses to ocean alkalinity enhancement to be measured in experimental manipulation studies.** Knowledge need (M=medium, H=high; measurement mode (MM=manual mode; S=sensor; SD=sensor in development). A minimum variable set is highlighted in bold. Selected references are provided as examples of protocols.

Type of response	Variable	Knowledge need	Measurement mode	Protocol reference
Chemical and environmental	<b>Carbonate chemistry parameters</b> {[HCO <sub>3</sub> <sup>-</sup> ], [CO <sub>3</sub> <sup>2-</sup> ], [CO <sub>2</sub> ], pCO <sub>2</sub> , Ω}	H	MM, S, SD	Dickson (2010); Bockmon and Dickson (2015)
	Dissolved organic matter	M	MM	Marañón et al. (2004); Sharp et al. (1995)
	<b>Dissolved inorganic nutrients</b>	H	MM	Worsfold et al. (2013)
	Resource availability (prey, light)	H	MM, S	Lawrence et al. (2017)
	<b>Particulate organic matter (C, N, P)</b>	H	MM	Verardo et al. (1990); Hilton et al. (1996); Pujo-Pay and Raimbault (1994); Fu et al. (2008)
	Trace metals (in solution and in aggregates)	M	MM	Guo et al. (2022); Hutchins et al. (2023)
	Biologically and biogeochemically relevant elements (e.g., Si, Mg:Ca)	M	MM	Brzezinski (1985); de Nooijer et al. (2017)
Physiological	<b>Basic physiology (respiration, photosynthetic, growth rates; morphometric measurements)</b>	H	MM, S	Iglesias-Rodriguez et al. (2008); Kelly et al. (2013); Farrell et al. (2009)
	<b>Some functional group-specific physiology (e.g., calcification, silicification, nitrification/denitri</b>	H	MM, S	Cohen et al. (2017); DeCarlo et al. (2019)

	<b>fication, toxin production)</b>			
	Physiological stress [e.g., heat shock proteins, oxidative stress-related proteins, photosynthetic stress (shifts in quantum yield), morphological alterations (e.g., cyst formation)	M	MM	O'Donnell et al. (2009); Moya et al. (2015); Trimborn et al. (2017)
	Incidence of pathogens and disease	H	MM	Asplund et al. (2014)
Reproduction	Spawning success	M	MM	Liu et al. (2011)
	<b>Size of offspring</b>	M	MM	Cao et al. (2018); Johnson (2022); Albright et al. (2010)
	Sperm motility	M	MM	Esposito et al. (2020); Havenhand et al. (2008)
	Epigenetic analysis	M	MM	Li et al. (2018); Lee et al. (2022)
	<b>Fecundity</b>	M	MM	Maranhão and Marques (2003); Thor and Dupont (2015)
	Hatching success	M	MM	Saigusa (1992)
Species interactions	Competition for resources	M	MM	Connell et al. (2013); Guo et al. (2022)
	Predation and species interactions	M	MM	Greatorex and Knights (2023); Bacus and Kelley (2023); Mitchell et al. (2023)

	Synergistic/antagonistic effects of other environmental parameters	M	MM, S	Gerhard et al. (2023); Khalil et al. (2023)
--	--	---	-------	---

411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461

#### 4.d Type of experiments

Laboratory experiments can be designed to both address short term responses and to explore the longer-term adaptation to chronic exposure to enhanced alkalinity conditions. Filtered natural seawater should be used, when possible, in incubations unless artificial seawater is required (for example, when studying the effect of metal concentrations). Short term manipulations involve the use of batch, semi-continuous and continuous incubation experiments. In *batch incubation experiments*, all resources are provided at the beginning of the incubation, without further addition and sampling takes place during a short time period (hours, days, weeks). Only gases and alkali can be added during the course of the experiment. When biological processes are measured, a phase during the life cycle (e.g., larva/adult; vegetative cells/gametes) or growth (healthy, exponentially-growing/resource-limited, stationary growing organisms /senescent organisms) is typically targeted. Sampling is conducted until the nutrients are consumed and beyond if decaying populations are the focus of the investigation.

Given that resources (light, nutrients) are the limiting factor in batch incubation experiments, the organisms are in the exponential growth phase for a limited time period. To expand sampling and replication during the exponential growth phase, resupply of nutrients using a *semi-continuous culturing* approach can prevent food/nutrients from becoming a limiting factor. When the studied organism is phototrophic, one must ensure subculturing (microbial cultures) or appropriate arrangement of organisms to prevent light limitation. The advantage of semi-continuous culturing is that it allows investigating trends over extended time periods, increase replication and higher yield. Generally, the resource is added manually or pumped from the nutrient supply vessel into the culture vessel during exponential growth or when specific conditions are met (e.g., when a certain biomass concentration is reached).

In *continuous cultures*, the rate of addition of nutrient is controlled to maintain steady state cell growth. This system is known as chemostat, where typically, a volume of culture medium is added and the same volume is removed from the growing culture. A challenge with this type of ‘bioreactors’ is that, over long time periods, they can be more susceptible to microbial contamination and long-term phenotypic and genotypic variance in the cultures (Reusch, 2013).

*Portable incubation experiments* that simulate regional in situ alkalinity deployments are an important step in understanding seawater alkalization and its impact on marine organisms prior to field testing. This type of incubation experiments, which simulate alkalinity additions under diverse local in situ parameters (e.g., temperature, irradiance, nutrients), can be accomplished using portable incubators onboard research vessels (i.e., deck incubations) or outdoors, at coastal research facilities (Fig. 2).

When studying photosynthetic organisms high-quality light filters should be attached to the acrylic tank to adjust photosynthetically active radiation (PAR) within the incubator (e.g., Fig. 2). To maintain in situ seawater temperatures, an inflow port can supply seawater to the incubator. Effort should be taken to ensure movement of seawater quickly through the incubator to maintain a uniform temperature.

When collecting natural seawater, one must consider how biological interactions (e.g., grazing) could confound results and filter accordingly. Unlike laboratory experiment, that allow for seawater-air phase CO<sub>2</sub> equilibration, portable incubation experiments require instantaneous alkalinity additions; thus, careful consideration should be given to the method of alkalinity addition used. When adding liquid alkalinity, e.g., solutions (e.g., 1 M) of NaOH one must consider that flocculation commonly occurs upon alkalinity addition (Subhas et al., 2022). Adding pulverized minerals directly to the treatment vessels is another option although this method may yield incomplete dissolution or slow dissolution (e.g., Fuhr et al., 2022) with undesirable effects including secondary precipitation, particle aggregation and detrimental biological impacts (NASEM, 2022). Some researchers have opted for mimicking mineral dissolution instead (see Gately et al., 2023). As in the traditional laboratory experiments described above, vessels within the incubator should ideally be aerated during experimentation. In addition to

462 chemical and biological parameters, PAR and temperature data should be collected throughout the experimental  
463 timeframe through discrete sampling or semi-continuously using sensors and data loggers. The best practices  
464 outlined in Box 1 should be adhered to when planning portable incubation experiments. Effort should be taken to  
465 position the incubator in a way that avoids confounding factors such as light contamination (e.g., from the ship).

## 466 467 **5. Sampling and analysis**

468  
469 Technical variability amongst experimental methods ranging from sampling and sample processing can  
470 propagate through the various steps before analysis; for example, chemical analysis and molecular work/sequencing  
471 can be error-prone (e.g., Catlett et al., 2020). The use of blanks every time sampling is conducted is essential for  
472 detecting contamination originating from the experiment itself or from the adjacent environment (e.g., exogenous  
473 sources such as surface contamination, flagellates in droplets through aeration, etc.). When possible, several barriers  
474 to contamination are recommended (e.g., filters at various points of aeration). Additionally, for samples (other than  
475 those preserved for analysis of alkalinity, dissolved inorganic carbon analysis or pH) that are kept for further  
476 analyses, contaminants that grow during shipping or while samples are being stored can sometimes be reduced by  
477 freezing at -80 °C, when possible, or by using the appropriate preservatives when storing at ambient temperature is  
478 required (e.g., ethanol, paraformaldehyde, glutaraldehyde). Attention should be paid to the material of vessels where  
479 samples and solutions are stored; for example, avoid borosilicate bottles to store nutrients or alkalinity solutions as  
480 silicate can be leached into solution.

481  
482 Establishing time series prior to the experiment to determine time frames regarding the appropriate length  
483 of the experiment and frequency of sampling is recommended. It is important to establish well defined time  
484 windows for sampling as well as frequency of sampling to capture physical, chemical and biological properties of  
485 the studied system. It is advisable to limit the time of sample storage to minimize observations that might confound  
486 interpretation of results (e.g., reverse weathering during storage) (Subhas et al., 2022).

### 487 488 **5.a Criteria for key parameters**

489 For the most part, laboratory experiments are aimed at elucidating the physiological performance and  
490 biogeochemical responses of organisms (rather than communities) to physical or chemical alterations in the  
491 environment although responses in ecological fitness could be drawn from laboratory experiments (Table 2).  
492 Importantly, environmental change can affect species differently and interactions between species that are sensitive  
493 to environmental change can function as ecological leverage points through which modest changes in abiotic  
494 conditions are amplified into large changes in marine ecosystems (see Kroeker and Sanford, 2022). These  
495 interactions can be measured as competition, predation, and symbiotic relationships (mutualism, commensalism and  
496 parasitism) that can vary along environmental gradients that cause stress (Stachowick, 2001; Bruno et al., 2003; Ma  
497 et al., 2023).

498  
499 Criteria for selection of species should include whether the organism is amenable to laboratory  
500 experimentation, the amount of background knowledge on the organism's physiology and biogeochemistry,  
501 ecological importance of the organism, and local and global impacts. Considerations when selecting organisms  
502 should also include geographic origin (e.g., temperate/tropical/polar) and ecosystem type (e.g., benthic vs pelagic).  
503 Special attention should be paid to those species that (1) significantly impact or respond biogeochemically to  
504 chemical changes caused by alkalinity addition (e.g., possibly calcifiers, photosynthetic organisms); (2) keystone  
505 organisms (e.g., corals, salmon, sea stars, toxin-producing phytoplankton); and (3) organisms/functional groups of  
506 known vulnerability to climate change (corals, urchins).

507  
508 Calcium carbonate producing organisms are particularly interesting because of their known sensitivity to  
509 changes in carbonate chemistry and because any alteration in their abundance or calcification rates could have  
510 implications in the CDR potential of alkalization. Mineralogical composition of carbonate containing organisms  
511 might possibly be affected by alkalization. For example, recent meta-analysis of studies exploring the effects of  
512 the carbonate chemistry shifts caused by ocean acidification revealed effects on shell state, development and growth  
513 rate (Figuerola et al., 2021). Biomineralization studies should explore species-specific responses driven by  
514 mineralogical composition (calcite, aragonitic, high/low Mg calcite) of their tests, shells and skeletons.  
515 Environmental and biological control on calcification particularly any changes in the Mg content in calcite driven by  
516 the use of brucite and other minerals potentially adding Mg to calcite must be reported as calcite with a high Mg  
517 content is less stable in aqueous solutions (Ries et al., 2016). Empirical studies have shown that the Mg/Ca ratio of

518 Mg-calcite producing organisms generally varies proportionally with seawater Mg/Ca (e.g., Ries, 2004; Ries, 2006)  
519 and therefore particular attention should be paid to the Mg content (and solubility) of biomineralized calcite. The  
520 addition to proposed Ca and Mg containing minerals - Ca(OH)<sub>2</sub> (slaked lime), Mg(OH)<sub>2</sub> (brucite), CaCO<sub>3</sub>  
521 (limestone) or (Mg,Ca)CO<sub>3</sub> (dolomite) - will alter the Mg/Ca ratio of the seawater. An extensive body of literature  
522 reports biogenic and abiotic precipitation of low-Mg calcite when seawater Mg/Ca falls within the calcite stability  
523 field (seawater molar Mg/Ca < 2) and the biogenic and abiogenic precipitation of aragonite and high-Mg calcite  
524 when seawater Mg/Ca falls within the aragonite stability field (seawater molar Mg/Ca > 2) (Ries, 2010). Thus,  
525 modification of local seawater Mg/Ca ratios by OAE has the potential to favor aragonite and high-Mg calcite  
526 organisms if seawater Mg/Ca is increased, and low-Mg calcite organisms if seawater Mg/Ca is decreased. This is an  
527 important area of future OAE research.

528  
529 Central to OAE laboratory experimentation is our ability to measure any possible stress induced by  
530 alkalization and learn about underlying mechanisms behind acclimation to the chemical alterations of seawater  
531 caused by OAE. This can be achieved by measuring basic functions (growth rates, size, reproductive success),  
532 sensitivities to alkalization might be organism-specific and possibly trophic level-specific (e.g., Voigt et al. 2003,  
533 Gilman et al. 2010) although most laboratory experiments do not address the complexity of trophic interactions.  
534 Similarly, measuring adaptation and diversity in acclimation between and within related organisms is a challenge  
535 and the ocean acidification literature revealed how important it is to pay attention to diversity of responses (see  
536 Kroeker et al., 2010).

537  
538 Stress is often measured as a reduction in organismal performance or fitness caused by environmental  
539 change (Schulte, 2014). In addition to these general physiological or behavioral responses, markers of stress such as  
540 oxidative stress are often used. For example, it is well established that the production of reactive oxygen species  
541 (ROS) can increase due to environmental stress including ocean acidification (Lesser, 2006; Lushchak, 2011). Many  
542 biomarkers are commonly used for studying oxidative stress in marine organisms (Cailleaud et al., 2007; Vehmaa et  
543 al., 2013) and an increase in ROS and superoxide dismutase and catalase activities have been reported in marine  
544 animals under stress (von Weissenberg et al., 2022). Heat shock proteins (HSPs) are also used as molecular markers  
545 of stress because of their abundance, high sensitivity to stress and being ubiquitously expressed (Gross, 2004).  
546 Among all HSPs, HSP70s are the most studied as a strong up-regulation of HSP70 production has been  
547 demonstrated broadly with the exception of *Hydra oligactis* (Bosch et al., 1998), and some Antarctic animals (La  
548 Terza et al. 2001; Place and Hofmann, 2005).

### 549 **5.b Measurements of nutrient uptake rates**

550  
551 The uptake rate of carbon and other nutrients that results in the observed standing stocks of particulate  
552 matter involve many physiological processes that are sensitive to changes in inorganic carbon chemistry and pH  
553 (Matsumoto et al., 2020). Chemical changes following the addition of alkalinity might alter physiological processes  
554 that represent sources (calcification, respiration) and sinks (photosynthesis) of CO<sub>2</sub>. One should also pay attention to  
555 the reciprocal interactions between these physiological processes and the chemically altered environment as even  
556 minor changes in biological processes, or in the balance between them, can have implications for the CDR potential  
557 and biodiversity.

558  
559 One of the most unknown effects of OAE is the fate of biological fixation rates of different elements (e.g.,  
560 carbon and N<sub>2</sub> fixation rates). Such rates are measured in batch cultures and bioassay (mixed natural community)  
561 incubation experiments (LaRoche et al., 2010). While the objective of culture experiments is to understand the effect  
562 of environmental parameters on the elemental uptake by particular species in a lab, bioassay experiments have to  
563 deal with a rather complex species interaction in the field or after subsampling of mesocosms in a lab (Hutchins et  
564 al., 2007; Paul et al., 2016). Labelled/enriched (~99%) stable isotope tracers is the most used method for rate  
estimation these days. The rate calculation is based on isotopic mass balance equation (Montoya et al., 1996):

$$565 \quad \text{C or N}_2 \text{ fixation rate} = \frac{[POM]}{t} \left( \frac{A_f - A_0}{A_e - A_0} \right) \quad (1)$$

566  
567 where, [POM] is the concentration of element of interest (C or N) at the end of the incubation. Likewise,  $A_f$   
568 = atom% in POM at the end of incubation,  $A_0$  = atom% in POM at the start of the incubation,  $t$  is time of incubation,  
and  $A_e$  = isotopic enrichment in the dissolved form after the tracer addition at the start of the incubation

569 This equation/method is sensitive to analytical protocols in routine incubations (White et al., 2020), and  
570 might be even more sensitive in OAE incubations due to the issue of gas equilibration in tightly capped bottles. While  
571 the C substrate-based incubations are supposedly straightforward in incubations, N<sub>2</sub> gas incubations face the challenge  
572 of under-equilibration leading to underestimation of rates. But OAE incubations can produce larger errors in the C  
573 fixation estimates as well. This is because NaHCO<sub>3</sub> is generally used as a C substrate. To estimate <sup>13</sup>C isotopic  
574 enrichment after tracer addition (term in equation 1), a DIC value is normally assumed (as it does not change much at  
575 a given region). But OAE is expected to increase (or fluctuate) DIC during the experimental period, and thus a  
576 measured DIC value should be used in the enrichment factor calculation. Likewise, the <sup>14</sup>C-method, which is widely  
577 used for marine primary production and calcification rate measurements due to its sensitivity (Nielsen, 1952), also  
578 requires treatment-specific determination of DIC concentrations. Likewise, slow dissolution of N<sub>2</sub> gas poses a  
579 challenge to accurately estimating isotopic enrichment factor (*A<sub>e</sub>*), and it is advisable to measure this term.

580 Although the analytical precision of C and N isotopes is of order of sub permil levels, many times the low reported  
581 rates (<0.1 nmol N L<sup>-1</sup> d<sup>-1</sup>) are questionable (Gradoville et al., 2017). Therefore, the detection limit of rate  
582 measurements and its proper reporting is a major concern. To overcome this, following the propagation of analytical  
583 and statistical errors in each term of mass balance equation (1), Gradoville et al. (2017) have proposed to report  
584 minimal quantifiable rates (MQR) and the limit of detection (LOD) in triplicate samples. We ought to follow these  
585 protocols in the rates measured in OAE. In addition, we must make sure to sample/filter sufficient water to achieve  
586 35 µg N and 150 µg C in the sample for reliable mass spectrometric measurements.

587  
588  
589  
590

591

592  
593  
594



595  
596  
597 **Figure 2.** A, B: Portable incubator with blue filters to adjust photosynthetically active radiation (PAR). A scalar  
598 PAR sensor (LI-COR) can be observed within the incubator (A, right side). C: laboratory experiment using aeration  
599 and sacrificial replication. Images were taken by James Gately (A, C) and Sylvia Kim (B).

600  
601 **6. Conclusions and recommendations**

602 The field of OAE faces a great diversity of challenges given the continuously evolving experimental  
603 approaches and emerging data availability that will undoubtedly provide new information and ideas to optimize best  
604 practice in laboratory experimentation. This chapter highlights the need for attention to the design, sampling,  
605 performance, and analysis of laboratory procedures used in OAE laboratory experiments. The criteria we present to  
606 achieve best practice in laboratory experimentation and design focus on reproducibility, factors affecting CDR  
607 potential and organism health (e.g., alkalinity conditions leading to flocculation, aggregation), establishing suitable  
608 experimental controls, and identifying the appropriate level of biological organization (physiological, molecular) to  
609 study biotic responses to OAE. Key response variables informing on alterations in seawater chemistry following  
610 alkalization, growth of organisms /biomass buildup/reproductive success, and biogeochemically relevant  
611 properties (e.g., photosynthesis, respiration, calcification) under elevated alkalinity conditions should be measured  
612 and reported. The main recommendations include:

- 613 • Ensure reproducibility through appropriate experimental design and replication.
- 614 • Determine alkalinity thresholds for the formation of precipitates for each experimental approach and
- 615 condition.
- 616 • In addition to the proposed alkalinity target values of 3000-4000  $\mu\text{mol kg}^{-1}$  (Renforth and Henderson,
- 617 2017), use concentrations exceeding these recommended values to mimic responses at the site of
- 618 deployment/non equilibrium and use intermediate alkalinity levels to identify potential nonlinear responses.
- 619 • Establish appropriate experimental design to address questions at specific levels of organization (chemical,
- 620 physiological, molecular) and assuming different scenarios (e.g., mimicking impacts at the site of
- 621 deployment in a non-equilibrated system versus steady state scenarios in an equilibrated system).

622  
 623 Given the emerging nature of ocean alkalinity enhancement as a research field, this chapter will evolve to  
 624 update guidelines as more results become publicly available. Frequent assessments of knowledge acquired from  
 625 emerging and future studies and review of best practices are needed to keep the OAE community engaged and  
 626 forward thinking.

### 627 628 629 **Acknowledgements**

630  
 631 This is a contribution to the “Guide for Best Practices on Ocean Alkalinity Enhancement Research”. We  
 632 thank our funders the ClimateWorks Foundation and the Prince Albert II of Monaco Foundation. Thanks are also  
 633 due to the Villefranche Oceanographic Laboratory for supporting the lead authors' meeting in January 2023.

### 634 635 **References**

636  
 637 Albright, R., Mason, B., Miller, M. and Langdon, C. (2010) Ocean acidification compromises recruitment  
 638 success of the threatened Caribbean coral *Acropora palmata*. PNAS 107, 20400–20404.

639  
 640 Asplund, M.E., Baden, S.P., Russ, S., Ellis, R.P., Gong and N., Hernroth, B.E. (2014) Ocean acidification  
 641 and host–pathogen interactions: blue mussels, *Mytilus edulis*, encountering *Vibrio tubiashii*. Environmental  
 642 Microbiology 16, 1029–1039.

643  
 644 Bach, L.T., Taucher, J., Boxhammer, T., Ludwig, A., The Kristineberg KOSMOS Consortium, Achterberg,  
 645 E.P., et al. (2016) Influence of ocean acidification on a natural winter-to-summer plankton succession: first insights  
 646 from a long-term mesocosm study draw attention to periods of low nutrient concentrations. PLoS ONE 11(8):  
 647 e0159068. doi:10.1371/journal.pone.0159068.

648  
 649 Bacus, S.C., Kelley, A.L. (2023) Effects of ocean acidification and ocean warming on the behavior and  
 650 physiology of a subarctic, intertidal grazer. Mar Ecol Prog Ser 711, 31-45.

651  
 652 Bockmon, E.E. and Dickson, A.G., 2015 An inter-laboratory comparison assessing the quality of seawater  
 653 carbon dioxide measurements. Marine Chemistry 171, 36-43.

654  
 655 Bosch, T.C.G., Krylow, S.M., Bode, H.R. and Steele, R.E. (1988) Thermotolerance and synthesis of heat-  
 656 shock proteins—these responses are present in *Hydra attenuata* but absent in *Hydra oligactis*. Proceedings of the  
 657 National Academy of Sciences U.S.A. 85, 7927–7931.

658  
 659 Brantley, S.L. (2008) Kinetics of mineral dissolution. In S.L. Brantley, J.D. Kubicki, A.F. White (Eds.),  
 660 Kinetics of Water–Rock Interactions, Springer Science (2008), pp. 151-210

661  
 662 Bruno, J.F., Stachowicz, J.J., Bertness, M.D. (2003) Inclusion of facilitation into ecological theory. Trends  
 663 in Ecology and Evolution 18: 119–25.

664  
 665 Brzezinski, M.A. (1985) The Si:C:N ratio of marine diatoms: interspecific variability and the effect of  
 666 some environmental variables. J Phycol. 21, 347–357.

667  
668



669 Byrne, M. and Przeslawski, R. (2013) Multistressor studies of the impacts of warming and acidification of  
670 the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology*, **53**, 582–596.  
671

672 Cailleaud, K., Maillet, G., Budzinski, H., Souissi, S., & Forget-Leray, J. (2007). Effects of salinity and  
673 temperature on the expression of enzymatic biomarkers in *Eurytemora affinis* (Calanoida, Copepoda). *Comparative*  
674 *Biochemistry and Physiology - A Molecular and Integrative Physiology*, **147**, 841–849.  
675

676 Cao, R., Wang, Q., Yang, D, Liu, Y., Ran, et al. (2018) CO<sub>2</sub>-induced ocean acidification impairs the  
677 immune function of the Pacific oyster against *Vibrio splendidus* challenge: An integrated study from a cellular and  
678 proteomic perspective. *Science of The Total Environment* **625**, 1574-1583.  
679

680 Calosi, P., et al. (2013) Adaptation and acclimatization to ocean acidification in marine ectotherms: an in  
681 situ transplant experiment with polychaetes at a shallow CO<sub>2</sub> vent system. *Phil. Trans. R. Soc. B* **368**, 20120444.  
682

683 Casadevall, A., and Fang, F.C. (2010) Reproducible Science. *Infection and Immunity* **78**, 4972–4975.  
684

685 Catlett, D., Matson, P.G., Carlson, C.A., Wilbanks, E.G., Siegel, D.A. and Iglesias-Rodriguez, M.D. (2020)  
686 Evaluation of accuracy and precision in an amplicon sequencing workflow for marine protist communities.  
687 *Limnology and Oceanography Methods* **18**, 20-40.  
688

689 Clair, T.A. and Hindar, A. (2005) Liming for the mitigation of acid rain effects in freshwaters: A review of  
690 recent results. *Environ. Rev.* **13**, 91–128.  
691

692 Cohen, S., Krueger, T., Fine, M. (2017) Measuring coral calcification under ocean acidification:  
693 methodological considerations for the <sup>45</sup>Ca-uptake and total alkalinity anomaly technique. *PeerJ* **5**:e3749.  
694

695 Connell, S.D., Kroeker, K.J., Fabricius, K.E., Kline, D.I., Russell, B.D. (2013) The other ocean  
696 acidification problem: CO<sub>2</sub> as a resource among competitors for ecosystem dominance. *Phil Trans R Soc B* **368**:  
697 20120442.<http://dx.doi.org/10.1098/rstb.2012.0442>.  
698

699 Cornwall, C.E. and Hurd, C.L. (2016) Experimental design in ocean acidification research: problems and  
700 solutions. *ICES Journal of Marine Science* **73**(3): 572–581.  
701

702 Crain CM, Kroeker K, Halpern BS. 2008 Interactive and cumulative effects of multiple human stressors in  
703 marine systems. *Ecol. Lett.* **11**, 1304-1315.  
704

705 Danger, M., Oumarou, C., Benest, D. and Lacroix, G. (2007) Bacteria can control stoichiometry and  
706 nutrient limitation of phytoplankton. *Functional Ecology* **21**, 202-210.  
707

708 Darling ES, Côté IM. 2008 Quantifying the evidence for ecological synergies. *Ecol. Lett.* **11**, 1278-1286.  
709

710 Davis, P.S. (1989) Short-term growth measurements of corals using an accurate buoyant weighing  
711 technique. *Marine Biology* **101**,389-395.  
712

713 DeCarlo, T.M., Comeau, S., Cornwall, C.E., et al. (2019) Investigating marine bio-calcification  
714 mechanisms in a changing ocean with in vivo and high-resolution ex vivo Raman spectroscopy. *Glob Change*  
715 *Biol.* **2019**; **25**: 1877– 1888.  
716

717 de Nooijer, L.J., van Dijk, I., Toyofuku, T., and Reichart, G.J. (2017). The impacts of seawater Mg/Ca and  
718 temperature on element incorporation in benthic foraminiferal calcite. *Geochemistry, Geophysics, Geosystems* **18**,  
719 3617–3630.  
720

721 Dickson, A. G. (2010) The carbon dioxide system in seawater: Equilibrium chemistry and measurements.  
722 In U. Riebesell, V. J. Fabry, & L. Hansson (Eds.), *Guide to best practices for ocean acidification research and data*  
723 *reporting* (pp. 17–40). Luxembourg: Publications Office of the European Union.  
724

725 Dodge, R.E. Wyers, S.C., FRITH, H.R., Knap, A.H., Smith, S.R., Cook, C.B. and Sleeter, T.D. (1984)  
726 Coral calcification rates by the buoyant weight technique—Effects of alizarin staining *J. Exper. Mar. Biol. Ecol.*, 75,  
727 217–232.

728  
729 Dupont, C.L., Barbeau, K., and Palenik, B. (2008) Ni uptake and limitation in marine  
730 *Synechococcus* strains, *Appl. Environ. Microb.*, 74, 23-31.

731  
732 Dupont, C. L., Buck, K.N., Palenik, B., and Barbeau, K. (2010) Nickel utilization in phytoplankton  
733 assemblages from contrasting oceanic regimes, *Deep-Sea Res. Pt. II*, 57, 553-566.

734  
735 Elser, J.J., Dobberfuhl, D.R., MacKay, N.A. and Schampel, J.H. (1996) Organism size, life history, and  
736 N:P stoichiometry: toward a unified view of cellular and ecosystem processes. *BioScience* 46: 674-684.

737  
738 Esposito, M.C., Boni, R., Cuccaro, A., Tosti, E. and Gallo, A. (2020) Sperm Motility Impairment in Free  
739 Spawning Invertebrates Under Near-Future Level of Ocean Acidification: Uncovering the Mechanism. *Front. Mar.*  
740 *Sci.* 6:794. doi: 10.3389/fmars.2019.00794.

741  
742 Farrell, A.P., Eliason, E.J. Sandblom, E., Clark, T.D. (2009) Fish cardiorespiratory physiology in an era of  
743 climate change. *Canadian Journal of Zoology* 87(10), 835-851.

744  
745 Ferderer, A., Chase, Z., Kennedy, F., Schulz, K.G. and Bach, L.T. (2022) Assessing the influence of ocean  
746 alkalinity enhancement on a coastal phytoplankton community. *Biogeosciences* 19, 5375–5399.

747  
748 Figuerola, B., Hancock, A.M., Bax, N., Cummings, V.J., Downey, R., Griffiths, H.J., Smith, J. and Stark,  
749 J.S. (2021) A review and meta-analysis of potential impacts of ocean acidification on marine calcifiers from the  
750 Southern Ocean. *Front. Mar. Sci.* 8:584445. doi: 10.3389/fmars.2021.584445.

751  
752 Forbes, V. and Calow, P. (2002) Population growth rate as a basis for ecological risk assessment of toxic  
753 chemicals. *Philos Trans R Soc Lond B Biol Sci.* 357, 1299–1306.

754  
755 Fuhr, M., Geilert, S., Schmidt, M., Liebetrau, V., Vogt, C., Ledwig, B., Wallmann, K. (2022) Kinetics of  
756 olivine weathering in seawater: an experimental study. *Frontiers in Climate* 4: 831587. doi:  
757 10.3389/fclim.2022.831587.

758  
759 Galic N, Sullivan LL, Grimm V, Forbes VE. 2018 When things don't add up: quantifying impacts of  
760 multiple stressors from individual metabolism to ecosystem processing. *Ecol. Lett.* 21, 568-577.

761  
762 Gerhard, M., Koussoroplis, A.M., Raatz, M. et al. (2023) Environmental variability in aquatic ecosystems:  
763 Avenues for future multifactorial experiments. *Limnology and Oceanography Letters* 8, 247–266.

764  
765 Gately, J.A., Kim, S.M., Jin, B., Brzezinski, M.A. and Iglesias-Rodriguez, M.D. (2023) Coccolithophores  
766 and diatoms resilient to ocean alkalinity enhancement: a glimpse of hope? *Science Advances* 9(24) . doi:  
767 10.1126/sciadv.adg6066.

768  
769 Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W., Holt, R.D. (2010) A framework for community  
770 interactions under climate change. *Trends in Ecology and Evolution* 25, 325–31.

771  
772 Godbold, J.A., Solan, M. (2013) Long-term effects of warming and ocean acidification are modified by  
773 seasonal variation in species responses and environmental conditions. *Philos Trans R Soc Lond B Biol Sci.*  
774 26;368(1627):20130186.

775  
776 Gradoville, M.R., Bombar, D., Crump, B.C., Letelier, R.M., Zehr, J.P., and White, A.E. (2017). Diversity  
777 and activity of nitrogen-fixing communities across ocean basins. *Limnology and Oceanography*, 62(5), 1895-1909.

778

779           Greatorex, R. and Knights, A.M. (2023) Differential effects of ocean acidification and warming on  
780 biological functioning of a predator and prey species may alter future trophic interactions. *Marine Environmental*  
781 *Research* 186, 105903.

782  
783           Gross, M. (2004) Emergency services: a bird's eye perspective on the many different functions of stress  
784 proteins. *Current Protein and Peptide Science* 5, 213–223.

785  
786           Guillard, R.R. and Ryther, J.H. (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* (Hustedt),  
787 and *Detonula confervacea* (Cleve Gran.). *Can J Microbiol*, 8, 229-239.

788  
789           Guo, J.A., Strzepak, R., Willis, A., Ferderer, A. and Bach, L.T. (2022) Investigating the effect of nickel  
790 concentration on phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement.  
791 *Biogeosciences* 19, 3683–3697.

792  
793           Hartmann, J., Suitner, N., Lim, C., Schneider, J., Marín-Samper, L., Arístegui, J., Renforth, P., Taucher, J.  
794 and Riebesell, U. (2023) Stability of alkalinity in ocean alkalinity enhancement (OAE) approaches – consequences  
795 for durability of CO<sub>2</sub> storage. *Biogeosciences*, 20, 781–802.

796  
797  
798           Harvey, B.P., Gwynn-Jones, D. and Moore, P.J. (2013). Meta-analysis reveals complex marine biological  
799 responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution* 3, 1016–1030.

800  
801           Havenhand, J.N., Buttler, F.R., Thorndyke, M.C. and Williamson, J.E. (2008) Near-future levels of ocean  
802 acidification reduce fertilization success in a sea urchin. *Curr. Biol.* 18, R651–R652.

803  
804           Hettinger, A., Sanford, E., Hill, T.M., Russell, A.D., Sato, K.N.S., Hoey, J., Forsch, M., Page, H.N.,  
805 Gaylord, B. (2012) Persistent carry-over effects of planktonic exposure to ocean acidification in the *Olympia* oyster.  
806 *Ecology* 93, 2758–2768.

807  
808           Hilton, J., Lishman, J.P., Mackness, S., Heaney, S.I. (1986) An automated method for the analysis of particulate  
809 carbon and nitrogen in natural waters. *Hydrobiologia* 141, 269-271.

810  
811           Ho, T.-Y. (2013) Nickel limitation of nitrogen fixation in *Trichodesmium*, *Limnol. Oceanogr.*, 58, 112-120.

812  
813           Hu, X. (2020). Effect of organic alkalinity on seawater buffer capacity: A numerical exploration. *Aquatic*  
814 *Geochemistry*. [https://doi.org/ 10.1007/s10498-020-09375-x](https://doi.org/10.1007/s10498-020-09375-x).

815  
816           Hurlbert, S.H. (2009) The Ancient Black Art and Transdisciplinary Extent of Pseudoreplication. *Journal of*  
817 *Comparative Psychology*, American Psychological Association 2009 4, 434–443.

818  
819           Hutchins, D.A., Fu, F.X., Zhang, Y., Warner, M.E., Feng, Y., Portune, K., ... and Mulholland, M.R. (2007).  
820 CO<sub>2</sub> control of *Trichodesmium* N<sub>2</sub> fixation, photosynthesis, growth rates, and elemental ratios: Implications for past,  
821 present, and future ocean biogeochemistry. *Limnology and oceanography*, 52(4), 1293-1304.

822  
823           Hutchins, D.A., Fu, F.X., Yang, S.C., John, S.G., Romaniello, S.J., Andrews, M.G. and Walworth, N.G.  
824 (2023) Responses of globally important phytoplankton groups to olivine dissolution products and implications for  
825 carbon dioxide removal via ocean alkalinity enhancement. *bioRxiv*, pp.2023-04.

826  
827           Iglesias-Rodríguez, M.D., Halloran, P.R., Rickaby, R.E.M., Hall, I.R., Colmenero-Hidalgo, E., et al. (2008)  
828 Phytoplankton calcification in a high CO<sub>2</sub> world. *Science* 320, 336–39.

829  
830           IPCC. 2021. *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the  
831 *Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Summary for Policymakers, V.  
832 Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I.  
833 Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, Yu R. and  
834 B. Zhou, eds. Cambridge University Press.

835  
836 Kelly, M.W., Padilla-Gamiño, Hofmann, G.E. (2013) Natural variation and the capacity to adapt to ocean  
837 acidification in the keystone sea urchin. *Global Change Biology* 19, 2536–2546.  
838  
839 Khalil, M., Doo, S.S., Stuhr, M. *et al.* (2023) Long-term physiological responses to combined ocean  
840 acidification and warming show energetic trade-offs in an asterinid starfish. *Coral Reefs*.  
841 <https://doi.org/10.1007/s00338-023-02388-2>  
842  
843 Kim, H.-C. and Lee, K. (2009) Significant contribution of dissolved organic matter to seawater alkalinity,  
844 *Geophys. Res. Lett.*, 36, L20603, 5 pp..  
845  
846 Koeve, W., Kim, H.-C., Lee, K., and Oschlies, A. (2011) Potential impact of DOC accumulation on  $f\text{CO}_2$   
847 and carbonate ion computations in ocean acidification experiments, *Biogeosciences Discuss.*, 8, 3797–3827,  
848 doi:10.5194/bgd-8-3797-2011.  
849  
850 Kroeker, K.J. and Sanford, E. (2022) Ecological leverage points: species interactions amplify the  
851 physiological effects of global environmental change in the ocean. *Annual Review of Marine Science* 14: 75–103.  
852  
853 Kroeker, K.J., Kordas, R.L., Crim, R.N. and Singh, G.G. (2010) Meta-analysis reveals negative yet variable  
854 effects of ocean acidification on marine organism. *Ecology Letters* 13, 1419-1434.  
855  
856 Kroeker, K.J., Kordas, R.L., Crim, R. *et al.* (2013b) Impacts of ocean acidification on marine organisms:  
857 quantifying sensitivities and interaction with warming. *Global Change Biology*, **19**, 1884–1896.  
858  
859 Kroeker, K., Micheli, F. & Gambi, M. (2013a) Ocean acidification causes ecosystem shifts via altered  
860 competitive interactions. *Nature Clim Change* **3**, 156–159.  
861  
862 LaRoche, J., Rost, B., and Engel, A. (2010). Bioassays, batch culture and chemostat experimentation. In  
863 Approaches and tools to manipulate the carbonate chemistry., Guide for Best Practices in Ocean Acidification  
864 Research and Data Reporting. In: Riebesell U., Fabry VJ, Hansson L., Gattuso J.-P.(Eds.), pp. (pp. 81-94).  
865  
866 LaTerza, A.L., Miceli, C. and Luporini, P. (2001) Divergence between two Antarctic species of the ciliate  
867 *Euplotes*, *E. focardii* and *E. nobilii*, in the expression of heat-shock protein 70 genes. *Molecular Ecology* 10, 1061–  
868 1067.  
869  
870 Lawrence, M.J., Eliason, E.J., Brownscombe, J.W., Gilmour, K.M., Mandelman, J.W. and Cooke, S.J.  
871 (2017) An experimental evaluation of the role of the stress axis in mediating predator-prey interactions in wild  
872 marine fish. *Comparative Biochemistry and Physiology, Part A* 207, 21-29.  
873  
874 Lesser, M.P. (2006) Oxidative stress in marine environments: Biochemistry and physiological ecology.  
875 *Annual Review of Physiology* 68, 253–278.  
876  
877 Lebrato, M., Garbe-Schönberg, D., Müller, M.N., Blanco-Ameijeiras, S., Feely, R.A. et al. (2020) Global  
878 variability in seawater Mg:Ca and Sr:Ca ratios in the modern ocean. *Proc. Natl. Acad. Sci. U.S.A.* 117, 22281–  
879 22292.  
880  
881 Lee, Y.H., Kim, MS., Wang, M. et al. (2022) Epigenetic plasticity enables copepods to cope with ocean  
882 acidification. *Nat. Clim. Chang.* 12, 918–927.  
883  
884 Li, Y., Lieu, Y.J., Cui, G., Cziesielski, M.J. et al. (2018) DNA methylation regulates transcriptional  
885 homeostasis of algal endosymbiosis in the coral model *Aiptasia*. *Sci. Adv.* **4**, eaat2142.  
886  
887 Liu, G., Innes, D. and Thompson, R.J. (2011) Quantitative analysis of sperm plane circular movement in  
888 the blue mussels *Mytilus edulis*, *M. trossulus* and their hybrids. *J. Exp. Zoology*, 315, 280-290.  
889

890 Lushchak, V.I. (2011) Environmentally induced oxidative stress in aquatic animals. *Aquatic Toxicology*  
891 101, 13–30.  
892

893 Ma, K.C.K., Monsinjon, J.R., Froneman, P.W., McQuaid, C.D. (2023) Thermal stress gradient causes  
894 increasingly negative effects towards the range limit of an invasive mussel. *Science of the Total Environment* 865,  
895 (2023), Article 161184.  
896

897 Maranhão, P. and Marques João, C. (2003) The influence of temperature and salinity on the duration of  
898 embryonic development, fecundity and growth of the amphipod *Echinogammarus marinus* Leach (Gammaridae).  
899 *Acta Oecol.*, 24, pp. 5-13.  
900

901 Marañón, E., Cermeño, P., Fernández, E., Rodríguez J., and Zabala, L. (2004) Significance and  
902 mechanisms of photosynthetic production of dissolved organic carbon in coastal eutrophic ecosystems. *Limnol.*  
903 *Oceanogr.* 49, 1652– 1666.  
904

905

906 Martin, B., Jager, T., Nisbet, R.M., Preuss, T.G., Grimm, V. (2014) Limitations of extrapolating  
907 toxic effects on reproduction to the population level. *Ecol Appl.* 24, 1972–1983.  
908

909 Martz, T.R., Daly, K.L., Byrne, R.H., Stillman, J.H. and Turk, D. (2015) Technology for Ocean  
910 Acidification Research: Needs and Availability. *Oceanography* 28 (2), Special Issue on Emerging Themes in Ocean  
911 Acidification Science: 40-47.  
912

913 Matsumoto, K., Tanioka, T. and Rickaby, R. (2020) Linkages between dynamic phytoplankton C: N: P and  
914 the ocean carbon cycle under climate change. *Oceanography* 33, 44–52.  
915

916 Middelburg, J.J., Soetaert, K., and Hagens, M. (2020). Ocean alkalinity, buffering and biogeochemical  
917 processes. *Reviews of Geophysics*, 58, e2019RG000681. [https://doi.org/ 10.1029/2019RG000681](https://doi.org/10.1029/2019RG000681).  
918

919 Millero, F.J., Woosley, R., Ditrolio, B. And Waters, J. (2009) Effect of ocean acidification on the  
920 speciation of metals in seawater. *Oceanography* 22, No. 4, Special Issue On The Future of Ocean Biogeochemistry  
921 in a High-CO<sub>2</sub> World pp. 72-85.  
922

923 Mitchell, A., Hayes, C., Booth, D.J. and Nagelkerken, I. (2023) Future shock: Ocean acidification and  
924 seasonal water temperatures alter the physiology of competing temperate and coral reef fishes. *Science of the Total*  
925 *Environment* 883, 163684.  
926

927 Montoya, J.P., Voss, M., Kahler, P., and Capone, D.G. (1996). A simple, high-precision, high-sensitivity  
928 tracer assay for N (inf2) fixation. *Applied and Environmental Microbiology*, 62(3), 986-993.  
929

930 Montserrat, F., Renforth, P., Hartmann, J., Leermakers, M., Knops, P. and Meysman, F.J.R. (2017) Olivine  
931 dissolution in seawater: implications for CO<sub>2</sub> sequestration through enhanced weathering in coastal environments.  
932 *Environmental Science & Technology* 51, 3960-3972.  
933

934 Moras, C. A., Bach, L. T., Cyronak, T., Joannes-Boyau, R., and Schulz, K. G. (2022) Ocean alkalinity  
935 enhancement – avoiding runaway CaCO<sub>3</sub> precipitation during quick and hydrated lime dissolution, *Biogeosciences*,  
936 19, 3537–3557.  
937

938 Morel, F.M.M., Rueter, J., Anderson, D.M. and Guillard, R.R.L. (1979) Aquil: A chemically defined  
939 phytoplankton culture medium for trace metal studies. *J. Phycol.* 15, 135– 141.  
940

941 Moya, A., Huisman, L., Forêt, S., Gattuso, J.-P., Hayward, D.C., Ball, E.E. and Miller, D.J. (2015) Rapid  
942 acclimation of juvenile corals to CO<sub>2</sub>-mediated acidification by upregulation of heat shock protein and Bcl-2 genes.  
943 *Mol Ecol*, 24: 438-452.  
944

945 NASEM: A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration, National  
946 Academies of Sciences, Engineering, and Medicine, Washington (DC), <https://doi.org/10.17226/26278>, 2022.  
947

948 Nawaz, S., Lezaun, J., Valenzuela, J.M. and Renforth, P. (2023). Broaden Research on Ocean Alkalinity  
949 Enhancement to Better Characterize Social Impacts. *Environmental Science & Technology* 57, 24, 8863–8869.  
950

951 Nielsen, S.E. (1952) The use of radio-active carbon ( $C^{14}$ ) for measuring organic production in the sea. *ICES*  
952 *Journal of Marine Science* 18, 117–140.  
953

954 O'Donnell, M.J., Hammond, L.M. and Hofmann, G.E. (2009) Predicted impact of ocean acidification on a  
955 marine invertebrate: elevated  $CO_2$  alters response to thermal stress in sea urchin larvae. *Mar Biol* 156, 439–446.  
956

957 Paiva, F., Brennecke, D., Pansch, C. and Briski, E. (2021) Consistency of aquatic enclosed experiments:  
958 The importance of scale and ecological complexity. *Diversity and Distributions* 27, 524–532.  
959

960 Paul, A.J., Bach, L.T., Schulz, K.-G., Boxhammer, T., Czerny, J., Achterberg, E.P., et al. (2015) Effect of  
961 elevated  $CO_2$  on organic matter pools and fluxes in a summer Baltic Sea plankton community. *Biogeosciences* 12:  
962 6181–6203.  
963

964 Paul, A.J., Achterberg, E.P., Bach, L.T., Boxhammer, T., Czerny, J., Haunost, M., ... and Riebesell, U.  
965 (2016). No observed effect of ocean acidification on nitrogen biogeochemistry in a summer Baltic Sea plankton  
966 community. *Biogeosciences*, 13(13), 3901-3913.  
967

968 Pistevos, J.C.A., Nagelkerken, I., Rossi, T. and Connell, S.D. (2016) Antagonistic effects of ocean  
969 acidification and warming on hunting sharks. *OIKOS* 126,  
970 <https://doi.org/10.1111/oik.03182>.  
971

972 Place, S.P. and Hofmann, G.E. (2005) Constitutive expression of a stress-inducible heat shock protein gene,  
973 hsp70, in a phylogenetically distant Antarctic fish. *Polar Biology* 28, 261–26.  
974

975 Pujo-Pay, M. and Raimbault, P. (1994) Improvement of the wet-oxidation procedure for simultaneous  
976 determination of particulate organic nitrogen and phosphorus collected on filters. *Mar. Ecol. Prog. Ser.* 105, 203-  
977 207.

978 Ragg, N.L.C., Gale, S.L., Le, D.V., Hawes, N.A., Burritt, D.A., Young, T., Jessica A. Ericson, Zoë  
979 Hilton, Ellie Watts, Jolene Berry, Nick King (2019) The Effects of Aragonite Saturation State on Hatchery-Reared  
980 Larvae of the Greenshell Mussel *Perna canaliculus*  
981

982 Reusch, T.B.H. (2014) Climate change in the oceans: evolutionary versus phenotypically plastic responses  
983 of marine animals and plants. *Evolutionary Applications* 7(2014) 104–122  
984 Reusch Adaptation and global change  
985 in the ocean. *Evolutionary Applications* 7, 104–122.  
986

987 Ridgwell, A., Schmidt, D.N., Turley, C., Brownlee, C., Maldonado, M.T., Tortell, P. et al. (2009). From  
988 laboratory manipulations to Earth system models: scaling calcification impacts of ocean acidification.  
989 *Biogeosciences*, 6, 2611–2623.  
990

991 Ries, J.B. (2004) The effect of ambient Mg/Ca on Mg fractionation in calcareous marine invertebrates: A  
992 record of Phanerozoic Mg/Ca in seawater. *Geology* 32, 981-984.  
993

994 Ries, J.B. 2006. Mg fractionation in crustose coralline algae: Geochemical, biological, and  
995 sedimentological implications of secular variation in the Mg/Ca ratio of seawater. *Geochimica et Cosmochimica*  
996 *Acta* 70, 891-900).  
997

998 Ries, J.B. (2010) Geological and experimental evidence for secular variation in seawater Mg/Ca (calcite-  
999 aragonite seas) and its effects on marine biological calcification. *Biogeosciences* 7, 2795-2849.

1000 Ries, J.B., Cohen, A.L., McCorkle, D.C. (2009) Marine calcifiers exhibit mixed responses to CO<sub>2</sub>-induced  
1001 ocean acidification. *Geology*, 37, 1131-1134.  
1002

1003 Ries, J.B., Ghazaleh, M.N., Connolly, B., Westfield, I., Castillo, K.D., 2016, Impacts of ocean acidification  
1004 and warming on the dissolution kinetics of whole-shell biogenic carbonates. *Geochimica et Cosmochimica Acta*  
1005 192: 318-337. doi: 10.1016/j.gca.2016.07.001.  
1006

1007 Saigusa, M. (1992) Control of hatching in an estuarine terrestrial crab I. Hatching of embryos detached  
1008 from the female and emergence of mature larvae. *Biol. Bull.*, 183, 401-408.  
1009

1010 Sala, M.M., Aparicio, F.L., Balagué, V., Boras, J.A., Borrull, E., Cardelús, C., et al. (2015) Contrasting  
1011 effects of ocean acidification on the microbial food web under different trophic conditions. *Ices J Mar Sci.* 73: 670–  
1012 679.  
1013

1014 Sanders T, Schmittmann L, Nascimento-Schulze JC and Melzner F (2018) High Calcification Costs Limit  
1015 Mussel Growth at Low Salinity. *Front. Mar. Sci.* 5:352. doi: 10.3389/fmars.2018.00352  
1016

1017 Sharp, J.H., Benner, R., Bennett, L., Carlson, C.A., Fitzwater, S.E, Peltzer, E.T. and Tupas, L.M. (1995)  
1018 Analyses of dissolved organic carbon in seawater: the JGOFS EqPac methods comparison. *Mar. Chem.* 48, 91–108.  
1019

1020 Stachowick, J.J. (2001) Mutualism, Facilitation, and the Structure of Ecological Communities. *BioScience*  
1021 51, 235–246.  
1022

1023 Subhas, A.V., Marx, L., Reynolds, S., Flohr, A., Mawji, E.W., Brown, P.J. and Cael, B.B. (2022) Microbial  
1024 ecosystem responses to alkalinity enhancement in the North Atlantic Subtropical Gyre. *Front. Clim.* 4:784997. doi:  
1025 10.3389/fclim.2022.784997.  
1026

1027 Thor, P. and Dupont, S. (2015) Transgenerational effects alleviate severe fecundity loss during ocean  
1028 acidification in a ubiquitous planktonic copepod. *Global Change Biology* 21, 2261–2271.  
1029

1030 Trimborn, S. Thoms, S., Brenneis, T., Heiden, J.P., Beszteri, S. and Bischof, K. (2017) Two Southern  
1031 Ocean diatoms are more sensitive to ocean acidification and changes in irradiance than the prymnesiophyte  
1032 *Phaeocystis antarctica*. *Physiologia Plantarum* 160, 155–170.  
1033

1034 USGCRP (U.S. Global Change Research Program). 2017. Climate Science Special Report: Fourth National  
1035 Climate Assessment, Vol. I, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart and T. K.  
1036 Maycock, eds. Washington, DC:USGCRP.  
1037

1038 Vandamme, D., Pohl, P. I., Beuckels, A., Foubert, I., Brady, P. V., Hewson, J. C., et al. (2015). Alkaline  
1039 flocculation of *Phaeodactylum tricorutum* induced by brucite and calcite. *Bioresour. Technol.* 196, 656–661. doi:  
1040 10.1016/j.biortech.2015.08.042  
1041

1042 Vehmaa, A., Hogfors, H., Gorokhova, E., Brutemark, A., Holmborn, T., & Engström-Öst, J. (2013)  
1043 Projected marine climate change: Effects on copepod oxidative status and reproduction. *Ecology and Evolution* 3,  
1044 4548–4557.  
1045

1046 Verardo, D.J., Froelich, P.N. and McIntyre, A. (1990) Determination of organic carbon and nitrogen in  
1047 marine sediments using the Carlo Erba NA-1500 analyzer. *Deep Sea Res. I* 37, 157-165.  
1048

1049 Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., et al. (2003) Trophic levels are differentially  
1050 sensitive to climate. *Ecology* 84, 2444–53.  
1051

1052 von Weissenberg, E., A. Jansson, K. A. Vuori, and J. EngströmÖst. 2022. Copepod reproductive effort and  
1053 oxidative status as responses to warming in the marine environment. *Ecol. Evol.* 12: e8594. doi:10.1002/ece3.8594  
1054

1055 Wang, H., Pilcher, D.J., Kearney, K.A., Cross, J.N., Shugart, O.M., Eisaman, M.D., and Carter, B.R.  
1056 (2023). Simulated impact of ocean alkalinity enhancement on atmospheric CO<sub>2</sub> removal in the Bering Sea. *Earth's*  
1057 *Future*, 11, e2022EF002816.  
1058  
1059 Wernberg, T., Smale, D.A. and Thomsen, M.S. (2012) A decade of climate change experiments on marine  
1060 organisms: procedures, patterns and problems. *Global Change Biology* 18, 1491–1498.  
1061  
1062 White, A.E., Granger, J., Selden, C., Gradoville, M.R., Potts, L., Bourbonnais, A., ... and Chang, B.X.  
1063 (2020). A critical review of the <sup>15</sup>N<sub>2</sub> tracer method to measure diazotrophic production in pelagic ecosystems.  
1064 *Limnology and Oceanography: Methods*, 18(4), 129-147.  
1065  
1066 Worsfold, P.J., Clough, R., Lohan, M.C., Monbet, P., Ellis, P.S., Quérel, C.R., Floor, G.H. & McKelvie,  
1067 I.D. (2013) Flow injection analysis as a tool for enhancing oceanographic nutrient measurements—A review.  
1068 *Analytical Chimica Acta* 803, 15-40.  
1069  
1070 Yang, B., Leonard, J., Langdon, C. (2023) Seawater alkalinity enhancement with magnesium hydroxide  
1071 and its implication for carbon dioxide removal. *Marine Chemistry* 253, 104251.