

Laboratory experiments in ocean alkalinity enhancement research

Maria D. Iglesias-Rodríguez^{1,2}, Rosalind E.M. Rickaby³, Arvind Singh⁴, James A. Gately^{1,2}

¹ Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara; Santa Barbara, CA 93106, U.S.A.

² Marine Science Institute, University of California, Santa Barbara, CA 93106, U.S.A.

³ Department of Earth Sciences, University of Oxford, Oxford, U.K.

⁴ Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, India.

Correspondence to: M. Débora Iglesias-Rodríguez, iglesias@ucsb.edu

Abstract. Recent concern about the consequences of continuing increases in atmospheric CO₂ as a key heat-trapping agent (USGCRP, 2017; IPCC, 2021) have prompted ocean experts to come together to discuss how to provide science-based solutions. Ocean alkalinity enhancement (OAE) is being considered not only as an ocean carbon dioxide removal (CDR) approach, but also as a potential way to mitigate ocean acidification. Over the last two decades, inter-laboratory comparisons have proven valuable in evaluating the reliability of methodologies associated with sampling and analysis of carbonate chemistry parameters, which have been routinely used in ocean acidification research. Given the complexity of processes and mechanisms related to ecosystem responses to OAE, consolidating protocols to ensure compatibility across studies is fundamental for synthesis and upscaling analysis. This chapter provides an overview of best practice in OAE laboratory experimentation and facilitates awareness of the importance of applying standardized methods to promote data re-use, inter-lab comparisons, meta-analysis and transparency. This chapter provides the reader with the tools to (1) identify the criteria to achieve the best laboratory practice and experimental design; (2) provide guidance on the selection of response variables for various purposes (physiological, biogeochemical, ecological, evolutionary) for inter-lab comparisons; (3) offer recommendation for a minimum set of variables that should be sampled and propose additional variables critical for different types of synthesis and upscaling; and (4) identify protocols for standardized measurements of response variables. Key recommendations include ensuring reproducibility through appropriate experimental design and replication, assessing alkalinity thresholds for secondary precipitates for each experimental approach and condition, using recommended targets of alkalinity (3000-4000 μmol kg⁻¹) and levels exceeding these concentrations to mimic responses at the site of deployment/non equilibrium and use intermediate alkalinity levels to identify potential nonlinear responses, and establish the appropriate experimental design to address questions at specific levels of organization (chemical, physiological, molecular) and assuming different scenarios (e.g., mimicking impacts at the site of deployment in a non-equilibrated system versus steady state scenarios in an equilibrated system).

1. Introduction

Laboratory studies need to be reproducible, consistent and transparent (Box 1) to provide the scientific community and regulators with useful information to move the field forward and facilitate the development of safe guidelines. Based on numerous modeling studies, ocean alkalinity enhancement (OAE) appears to be a promising ocean carbon dioxide removal (CDR) approach, with the likely beneficial side effect of mitigating ocean acidification (Burt et al., 2021; Hartmann et al., 2023; NASEM, 2022; Wang et al., 2023). Laboratory experiments are urgently needed to determine the CDR potential of various OAE methods as well as OAE impacts at various levels of biological organization (ecological, physiological, biochemical, molecular). The emerging empirical studies are offering insight while revealing gaps in our knowledge of the mechanisms governing OAE and its effect on marine biota (e.g., Ferderer et al., 2022; Gately et al., 2023; Yang et al., 2023). For example, the conditions preventing or limiting the formation of secondary precipitates and the pros and cons of various alkali are still under debate. Given that empirical work on OAE is still in its infancy and that some of the assumptions based on modeling studies remain untested, this chapter is an evolving document that will be updated as the OAE community continues to release results.

54
55 Laboratory manipulations allow making observations in a highly controlled environment using model
56 species or subsets of populations (selected species or populations). Results are generally considered highly
57 reproducible (Box 1) and therefore laboratory manipulations are viewed as a necessary step to either generate
58 hypotheses to test in the field or vice versa, when field experimentation is an option. Under the latter, field
59 observations guide the laboratory experiments to validate field results in well-known systems and under tightly
60 controlled conditions.

61
62 A number of approaches – batch, semi-continuous and continuous cultures – have been used to address
63 diverse OAE settings (e.g., at the point of deployment, under steady state conditions, air- versus non air-equilibrated
64 seawater) and various biological scenarios (specific stages of growth, life cycle, and to explore abrupt/short term
65 versus long term responses to manipulations). In some cases, specific stages during the life cycle of organisms can
66 be selected (for example, larval versus adult stage; sexual versus asexual phase). Time series laboratory experiments
67 are less restricted than mesocosm experiments with regards to the duration of experiments because they tend to be
68 ‘cleaner’, with relatively low bacteria numbers and generally without biological confounding factors (viruses,
69 predation, competition for resources, etc.). Therefore, the cause–effect relationships are easier to elucidate as
70 conditions and organisms can be tested in relative isolation, and there is the possibility of extensive replication.

71
72 The main limitation of laboratory experiments is that the dynamic phenomena occurring in the natural
73 environment cannot be captured in the laboratory and, therefore, results may not be applicable to real life scenarios.
74 For example, in laboratory experiments the influence of mixing processes, conditions governing particle flocculation
75 or the linkage to higher levels of biological organization (e.g., predation) are difficult to discern (see Forbes and
76 Calow, 2002; Martin et al., 2014). Portable lab experiments, such as deck incubations aboard research vessels or
77 outdoor incubations, with some influence from the local environment (e.g., diurnal alterations of light, water flow
78 through from the coast to maintain in-situ temperature) as well as community-level mesocosm experiments are the
79 conduit to field manipulations. These large-scale community experimental tanks address the importance of the
80 physico-chemical conditions, space, density-dependent effects, biotic interactions and complexity of natural
81 environments in their response to OAE manipulations\buffering, or boosting, the direct effects of environmental
82 stress on organisms (Paiva et al., 2021).

83
84 This chapter provides best practice guidelines in OAE laboratory experimentation and offers
85 recommendations to enable data re-use, inter-lab comparisons, and transparency. We offer recommendations
86 regarding (1) the criteria to achieve the best laboratory practice and experimental design; (2) the selection of
87 response variables for various purposes (physiological, biogeochemical, ecological, evolutionary) for inter-lab
88 comparisons; (3) a minimum set of variables that should be sampled and additional variables critical for different
89 types of synthesis and upscaling; and (4) protocols for standardized measurements of response variables.

90 91 92 **2. Lessons learned from ocean acidification research**

93 The rich insights obtained in ocean acidification research are key to supporting OAE studies. However, as
94 crucial as it is to follow guidelines when designing laboratory experiments, it is equally important to acknowledge
95 that there may be potential confounders and challenges that may not be accounted for in the guidelines. Being able
96 to conduct quantitative laboratory intercomparisons, including interspecies comparisons, will be critically dependent
97 on identifying recommendations regarding experimental design, sample collection and data analysis. Important
98 considerations include the source of alkalinity, rate of alkalinity addition, testing air-CO₂-equilibrated *versus* non-
99 equilibrated seawater, and the effect of ancillary variables (e.g., temperature) in multifactorial experiments which
100 are known to yield complex and variable results (e.g., see the interactive effects of ocean acidification and warming
101 - Harvey et al., 2013). The guidelines provided in this chapter should significantly improve the quality and impact of
102 the OAE research, which is required to meet the identified societal need for research on OAE and other types of
103 ocean CDR (NASEM, 2021).

104
105 An exploration of procedures, patterns and challenges associated with ocean acidification research has
106 offered ideas on how to design rigorous and reproducible laboratory experiments that enable measuring and
107 monitoring carbonate chemistry shifts and biological responses to ocean acidification (Cornwall and Hurd, 2016).
108 Cornwall and Hurd (2016) reported that 95% of the experimental work between 1993 and 2014 had interdependent
109 or lacked replication in clearly defined treatments, or did not report sufficient methodological detail. More broadly,

110 results from Wernberg et al. (2012) from marine climate change experiments between 2000 and 2009, reported that
111 ~49% of the experiments had identifiable issues with their experimental procedures, and 91% of the experiments
112 reported showed a lack of treatment replication or pseudo-replication. Amongst the studies, 9% included
113 extreme/unrealistic treatments of temperature or pH far beyond worst case scenario projections (Wernberg et al.,
114 2012) although ‘extreme’ pH/alkalinity conditions may prove useful to define thresholds of tolerance and upper
115 limits of alkalinity enhancement, and to understand underlying physiological mechanisms of acclimation to
116 alkalinization. While the urgent need for field trials requires careful consideration of treatment levels, in order to
117 maximize the insight gained from OAE experiments, testing conditions outside the year 2100 IPCC CO₂ emission
118 scenarios are encouraged. These conditions outside worst case scenario projections will further our knowledge on
119 the mechanisms governing biological (e.g., shell production) and abiotic (e.g., particle aggregation, secondary
120 precipitation) responses to applied chemical CDR.
121

122 Like in ocean acidification research, careful attention should be given to the advantages and disadvantages
123 that concern the choices of dissolved inorganic carbon species to measure, and how error propagation will affect the
124 calculated parameters (Martz et al., 2015). Moreover, dissolved organic matter (DOM) is known to contribute to
125 alkalinity (Kim and Lee, 2009; Koeve et al., 2010) although the presence of strong acidic groups in organic matter
126 can decrease net alkalinity (Hu, 2020; Middelburg et al., 2020). Depending on the type of system under
127 investigation, attention should be paid to whether to apply titration alkalinity (typically used in ocean studies) versus
128 the charge balance approach (often used in freshwater systems, with high concentrations of dissolved organic
129 matter) (see Middlebufig et al., 2020). Results from ocean acidification mesocosm experiments focused on
130 phytoplankton revealed that nutrient-limited communities appeared to be more responsive to changing carbonate
131 chemistry than those having access to high inorganic nutrient concentrations (see Paul et al., 2015; Sala et al., 2015;
132 Bach et al., 2016). These observations indicate that trophic state might play a role in the susceptibility of organisms
133 to the changes in carbonate chemistry driven by alkalinization. Also, competition between species has been found to
134 be altered under various carbonate chemistry conditions (see Kroeker et al., 2013a), which merits a focus on
135 experiments that address preferential selection of taxonomic groups under different alkalinity conditions. Although
136 applying nutrient-limiting conditions is experimentally challenging, understanding how species succession and
137 community composition might respond to alkalinization could in part be addressed in a laboratory context.
138

139 While it is fairly straightforward to determine how individual changes in parameters influence chemical
140 and biological responses, understanding impacts of multiple parameters [e.g., increased alkalinity and warming,
141 increased alkalinity and resource availability (nutrients, light, prey)] can be challenging as they can interact in
142 complex ways. Indeed, ocean acidification research revealed antagonistic, synergistic, and additive responses when
143 studying ocean acidification and warming (Byrne and Przeslawski, 2013; Kroeker et al., 2013b; Harvey et al., 2015;
144 Pistevo et al., 2016). Identifying tipping points and interactive effects when other parameters (e.g., temperature) are
145 altered in seawater, in addition to alkalinity, is critical given the capacity of these parameters to drive (otherwise
146 unpredictable) shifts in species abundances, biodiversity and community composition, physiological outputs,
147 survival, and reproduction (Crain et al., 2008; Darling and Côté, 2008; Galic et al., 2018).

148 **Box 1. Criteria for best laboratory practice**

- 149 · **Reproducibility.** From the emerging OAE research (e.g., regarding the formation of secondary
150 precipitates - see Montserrat et al., 2017 versus Fuhr et al., 2022; and Moras et al., 2022) and the ocean
151 acidification literature (e.g., see Ridgwell et al., 2009), we have learned that similar approaches can
152 lead to conflicting and unresolved outcomes. Without appropriate reporting of sample collection,
153 methodology and data processing, it is challenging to re-analyze the data and reconcile the
154 discrepancies. As the field emerges and evolves, it will be required to reevaluate early experiments and
155 possibly re-analyze results with updated protocols.
- 156 · **Defining inclusion and exclusion criteria.** In order to reduce confounding covariates, attention
157 must be paid to factors affecting flocculation, aggregation of particles (e.g., possibly impacted by
158 dissolved organic matter increases after phytoplankton blooms), fluctuations in temperature, which
159 affect mineral dissolution and precipitation rates, and biological and physiological properties,
160 including stage during the life cycle, trophic state, and seasonality, that affect the susceptibility of
161 organisms to OAE (e.g., see Vandamme et al., 2015; Subhas et al., 2022).
- 162 · **Establishing experimental controls.** In OAE experimental designs, controls must be appropriately
163 selected. These could include seawater without added alkalinity, seawater ± nutrients/food, treatments

164 with and without the organisms tested. When mineral dissolution is too slow, an alternative analog that
165 reproduces the basic chemistry is encouraged (for example, the use of salts and alkali; e.g., CaCl_2 and
166 Na_2CO_3 to mimic the effect of limestone-based mineral dissolution). Controls could also contain an
167 alternative form of alkalinity that alters the seawater carbonate chemistry solely, without adding carbon
168 or trace metals (e.g., NaOH).
169 · **Basic biological responses.** Studies on organisms' physiological responses (e.g., growth,
170 respiration, size, reproduction, photosynthesis and calcification) are recommended. These responses
171 can be measured directly; for example, as uptake rates of solutes using traditional assays, mass
172 spectrometric methods for indirect assessment of changes in elements, or molecular responses using
173 markers of functional processes. Rates of growth and calcification can also be measured by changes in
174 dry mass or buoyant mass in many types of organisms, especially in macroinvertebrates and
175 macroalgae (see Dodge et al., 1984; Davis, 1989; Sanders et al., 2018). For organisms that undergo
176 development one must determine which stage of development (e.g., larval vs adult; vegetative vs
177 gamete stage) to target. Also, when altering more than one parameter, particular attention must be paid
178 to potential confounding effects. Multi-factorial experiments can be used to explore the weight of each
179 parameter.

180 181 **3. Seawater media preparation and manipulation of carbonate chemistry**

182 The different steps in experimental design are outlined in Table 1. The process starts with natural or
183 artificially made seawater with or without nutrient additions. One must consider whether adding nutrients/food/prey
184 is required; for example, whether exploring OAE impacts is intended in conjunction with specific scenarios, e.g.,
185 nutrient fertilization, specific stages of growth or population development, and the extent to which nutrient additions
186 or any other basic manipulation of the environmental conditions might impact the interpretation of results. For OAE
187 manipulations where sterilization is required for the experimental set up, autoclaving is discouraged given the
188 alterations in carbonate chemistry, including loss of CO_2 , leading to a decrease in dissolved inorganic carbon and
189 alterations in alkalinity (increase with increasing salinity/decrease with precipitation of carbonate) triggered by
190 autoclaving. Instead, filter-sterilization of seawater through small pore size filters (e.g., $0.22 \mu\text{m}$ filters) is required
191 to remove particles and most bacteria, and produce the stock media where different manipulations are applied to
192 create different alkalinity treatments.
193

194 There are several approaches to simulating the addition of alkalinity that capture different components of
195 any manipulation experiment. The first approach could be testing the impact of instantaneous addition of alkalinity
196 to seawater to mimic the impact on seawater chemistry and ecosystems at the point of deployment. The second
197 involves aeration and equilibration with the atmosphere to explore the physico-chemical response to a steady state/
198 equilibrated scenario. In the latter instance, the medium is aliquoted out to the experimental vessels/tanks where
199 aeration is applied to promote air equilibration. Monitoring carbonate chemistry through time enables determining
200 when equilibration of seawater with air occurs.
201

202 **Table 1. Experimental considerations for OAE experimentation.**

203 **Medium preparation:** the seawater can be obtained from coastal or open ocean sites. Filtered seawater or, when
204 appropriate (e.g., when growing autotrophic organisms), seawater supplemented with nutrients; for example, using
205 f/2 or variations of f/2 media (see Guillard and Ryder, 1962) will be used for growing organisms. Seawater media
206 can also be prepared from artificial recipes (e.g., Aquil medium; Morel et al., 1979) when specific compounds or
207 elements need to be altered in seawater. Media must be sterilized by filtration rather than through autoclaving and
208 nutrients can be added, typically from stock solutions. When possible, moderate aeration should be applied. Types
209 of alkali include adding pulverized mineral directly to the media and promote dissolution physically (e.g., by
210 stirring); dissolving the mineral separately and filter out any particles remaining in the media before
211 experimentation; dissolving salts to mimic the chemistry of the dissolved alkali (e.g., to mimic limestone
212 dissolution, dissolve CaCl_2 and NaCO_3 , which result in higher dissolution rates); and adding liquid alkali such as
213 NaOH . Establishing time series prior to the experiment to determine time frames regarding length of experiment,
214 frequency of sampling, etc. is recommended. **Experimental design:** in addition to optimizing reproducibility by
215 designing enough replication and test the reproducibility of the method, researchers should remain engaged with
216 respect to protocols and experimental design to avoid artifacts and undesirable side effects of methodology. When
217 possible, ensure equilibration of seawater gasses with air and define experimental time frames to test impacts under
218 conditions representative of the site of deployment (where limited gas exchange occurs) and those representative of
219 steady state/equilibrated conditions. Although most laboratory experiments address short term impacts, chronic

220 effects can be tested in long term incubations. **Sampling and analysis:** the parameters to be considered should allow
 221 inter-lab comparisons, address functional properties of organisms (e.g., calcification, silicification, particulate
 222 organic carbon) and fulfill needs to improve model parameterizations. It is important to establish well defined time
 223 windows for sampling as well as frequency of sampling to capture physical, chemical and biological properties of
 224 the studied system. It is advisable to limit the time of sample storage to minimize observations that might confound
 225 interpretation of results (e.g., reverse weathering during storage). Stock solutions (e.g., nutrient and alkalinity
 226 solutions) must be stored in the appropriate vessels to avoid contamination from leachates coming out of the vessel
 227 itself (e.g., silicate contamination from solutions stored in borosilicate containers). Detection limits and accuracy
 228 and precision should be offered for each protocol.
 229

Medium preparation	Experiment design	Sampling and analysis
<p>Natural/artificial seawater</p> <p>Filter sterilization (e.g., 0.22 um)</p> <p>+/- nutrient addition</p> <p>+/- aeration</p> <p>Type of alkalinity treatments</p> <ul style="list-style-type: none"> • Pulverized mineral • Pre-dissolved mineral • Dissolved salts • Liquid alkali <p>Pre-equilibrated vs non-equilibrated seawater with air phase</p> <ul style="list-style-type: none"> • Carbonate chemistry • Flocculation/aggregation • Biology 	<p>Best actions to maximize confidence</p> <ul style="list-style-type: none"> • Within study replication and pseudo-replication • Coordinated networks (teams sharing progress to decide on best protocols) <p>Preliminary time series of TA and carbonate chemistry</p> <ul style="list-style-type: none"> • Define experimental time frames • Assess TA upper limits • Expand the upper limits to address impacts at site of deployment <p>Abrupt vs chronic biology impacts</p> <ul style="list-style-type: none"> • Short-term tests (acclimation) • Long-term experiments (adaptation) 	<p>Criteria for key parameters</p> <ul style="list-style-type: none"> • Inter-lab comparisons • Functional properties • Model parameterization <p>Sampling frequency and timing</p> <ul style="list-style-type: none"> • Select time window for sampling • Identify sampling frequency that captures key chemical, physical or biological features <p>Limit storage to minimize artifacts</p> <p>Identify and report key analytical parameters affecting error</p> <ul style="list-style-type: none"> • Detection limits • Measurement accuracy/precision • Identify any impact of experimental design on uncertainties

230
 231
 232 **3.a Sources of alkalinity**

233 As yet, it is unclear what the optimal method or source of alkalinity enhancement may be in order to
 234 simulate the desired chemistry in seawater media. Proposed sources of alkalinity include silicate minerals (olivine,
 235 basalt), brucite, limestone and its derivatives (quicklime and portlandite), NaOH and mine tailings (NASEM, 2021;
 236 Nawaz et al., 2023). Given the slow dissolution kinetics of the minerals, generating alkaline solutions artificially is
 237 acceptable. For example, Gately et al. (2023) simulated alkalinity enhancement via a limestone-inspired solution by
 238 adding Na₂CO₃ and CaCl₂ or its hydrated form (CaCl₂H₄O₂) to seawater. Adding Na₂CO₃ raises TA and DIC in a
 239 2:1 ratio, with 2 moles of TA added by 2 conservative Na⁺ ions in Na₂CO₃, and 1 mole DIC added by CO₃²⁻. CaCl₂
 240 does not raise alkalinity because it adds equal amounts of positive and negative conservative charge to the solution
 241 from Ca²⁺ and 2 x Cl⁻. However, it does raise the calcium in solution and therefore the saturation state of the
 242 seawater with respect to CaCO₃.
 243

244 Many possibilities for solid or liquid alkalinity additions are being considered (see chapter 3). While adding
 245 minerals as precursors of alkalinity can provide a source of potentially beneficial nutrients (e.g., silicate, iron,
 246 magnesium) (Hartmann et al. 2013), the possible toxic effect of metals leached out of minerals, an example being
 247 nickel (Ni) leached from olivine (Montserrat et al., 2017) is of concern. The use of NaOH is currently gaining
 248 attention given that its environmental footprint is perceived as smaller than the mining of alkaline minerals, which
 249 necessitate an expansion of mining operations, transportation, and industrial processing, which are energetically
 250 costly and can lead to air pollution. Additionally, the amount of Na added to seawater is very small relative to the
 251 large background of NaCl in seawater.
 252

253 The addition of NaOH and other forms of alkalinity to seawater cause initial spikes in pH and a drop in
 254 aqueous CO₂ that can be balanced to a steady state via bubbling with air (Table 1). Determining abiotic and biotic

255 responses to the initial spikes in pH and drops in CO₂ is an important step in addition to understanding responses
256 under steady-state conditions. It may be that large manipulations of alkalinity are needed to elicit a measurable and
257 reproducible response, and the required alkalinity concentrations will be refined with more detailed modeling but,
258 based on current information, proposed targets for alkalinity manipulations are 3000-4000 μmol kg⁻¹ (Renforth and
259 Henderson, 2017). ~4000 μmol/kg is the concentration of alkalinity expected at locations in the ocean where
260 alkalinity is initially added, and ~3000 μmol/kg is the concentration of alkalinity expected once ocean circulation
261 has dispersed the alkalinity over a larger area (Renforth and Henderson, 2017). Alkalinity thresholds for the
262 formation of precipitates will need to be determined for each experimental approach and condition. It is however
263 recommended that researchers consider using alkalinities exceeding the recommended targets, and utilize
264 intermediate treatments (e.g., 2000, 4000, 7000 μatm/kg seawater) rather than just low/high treatments, in order to
265 identify potential nonlinear and even parabolic responses. This approach led to important and unexpected outcomes
266 in ocean acidification research (e.g., Ries et al., 2009).

267 **3.b Impacts of impurities/metal leachates**

268 An important consideration in OAE studies is the impact of metals leached from dissolving minerals and
269 their ecotoxicological potential on marine organisms. For example, although some elements (e.g., Fe and Mg)
270 leached out of minerals could be beneficial micronutrients, the potentially toxic effect of metals such as nickel (Ni)
271 (Montserrat et al., 2017), leached from olivine, is of concern. Diverse responses have however been reported with
272 respect to Ni and it appears that some cyanobacteria rely on Ni more than other photosynthetic organisms (see
273 Dupont et al., 2008, 2010; Ho, 2013). A recent laboratory study testing olivine leachates (containing Si, Ni, Mg, Fe,
274 Cr and Co) in phytoplankton revealed either positive or neutral physiological short term responses in all treatments
275 (Hutchins et al., 2023). However, one should consider the role of long-term experiments to examine organismal and
276 population adaptation of metal exposure as well as potential bioaccumulation and biomagnification impacts in
277 consumers.
278

279
280 Another important consideration is the effect of pH on metal speciation as pH and a change in the
281 concentration of OH⁻ and CO₃²⁻ ions can affect the solubility, adsorption, toxicity, and rates of redox processes of
282 metals in seawater thus altering the interactions of metals with marine organisms (Millero et al., 2009). When
283 dissolving minerals in seawater one must consider nonstoichiometry and incomplete dissolution perhaps as a result
284 of dissolution of impurities, precipitation of secondary minerals, or preferential leaching of elements from the
285 mineral surface (Brantley, 2008, NASEM, 2021). The formation of secondary precipitates has been observed in
286 several studies exploring the dissolution of olivine (Fuhr et al., 2022), and limestone derivatives (Moras et al., 2022;
287 Gately et al., 2023; Hartmann et al., 2023). Using an alkaline solution rather than reactive alkaline particles has been
288 recommended to reduce carbonate precipitation unless seawater critical supersaturation levels are exceeded
289 (Hartmann et al., 2023). In addition, runaway CaCO₃ precipitation, a condition where more alkalinity is removed
290 than initially added, reduces the OAE CO₂ uptake efficiency. More complex precipitates containing Fe, Si, and P
291 were observed in a study using a limestone-inspired OAE approach revealing that mineral precipitation caused by
292 seawater alkalization can also remove inorganic nutrients from solution (Gately et al., 2023).
293


294 Maintaining alkalinity following OAE is critically dependent on the carbonate saturation state, its temporal
295 evolution, and particle surface processes (Hartmann et al., 2023). To minimize the loss of alkalinity and maximize
296 alkalinity enhancement, Hartmann et al (2023) propose the application of an alkaline solution in CO₂ equilibrium
297 with the atmosphere and/or solutions with tested saturation levels to prevent a further increase in supersaturation,
298 and the precipitation of carbonate to avoid loss of alkalinity. A separate reservoir where alkaline solutions have
299 been prepared is desirable for testing upper limits of alkalinity addition and identifying saturation thresholds to
300 minimize precipitation.

301 **4. Experimental design**

302 **4.a Experimental replication**

303
304 Replication is important to determine if results are reproducible although one must consider that when results are so
305 dependent on precise experimental conditions that replicability is needed for reproducibility, the result may be
306 unique and potentially less relevant than a phenomenon that can be reproduced by a variety of independent, non-
307 identical approaches (see Casadevall and Fang, 2010). A number of experimental designs can be used to achieve
308 adequate statistical replication (Fig. 1). For example, simple replication involves experimental units (each of the
309 replicates) per treatment where all the conditions are manipulated independently but in the same way for that
310

311 treatment and where responses to the treatment are measured [defined by Hurlbert (2009) as the “evaluation unit”]
 312 and each experimental unit can be considered as independent. In temporal replication, multiple measurements are
 313 made through time (temporal trends) on the same experimental unit. Sacrificial replication involves the use of
 314 multiple sampling times per treatment (for example, a time series) and multiple experimental units at the time of
 315 samplin. Each approach has distinct strengths and limitations, and the choice of the approach depends on the
 316 scientific questions and the extent of the risk of error propagation. For example, one might choose sacrificial
 317 replication for certain chemical manipulations that require sampling from vessels with comparable volumes but
 318 choose instead temporal replication for monitoring the evolution of a microbial culture or the physiology of fish over
 319 time under certain alkalinity conditions.
 320
 321
 322



323
 324
 325 **Figure 1.** Examples of experimental laboratory design with regards to replication. Each treatment, represented by a
 326 colour, contains experimental units ~~contains~~ (replicates). Each experimental unit is treated as an independent
 327 experiment except in the sacrificial replication approach, where each replicate is treated statistically as an
 328 experimental unit.
 329

330 4.b Preliminary experiments

331 In addition to testing the biological responses to abrupt enhanced alkalinity, marine organisms can be
 332 exposed to enhanced alkalinity conditions after equilibration of seawater pCO₂ with that in the air-phase following
 333 alkalinity addition. Ideally, aeration should be maintained to ensure O₂ levels required by marine animals and also
 334 maintain stable pCO₂ levels in the alkalinity perturbation experiments. Depending on the organism tested (a few
 335 organisms do not tolerate aeration in tanks), aeration might or might not remain for the duration of the experiment
 336 (Table 1). The vessels used in OAE experiments might not be traditional tanks used in aquaria, but rather any type
 337 of container adequate for different type of organisms (e.g., culture flasks for bacteria, conical flasks, carboys for
 338 phytoplankton, open tanks for echinoderms and fish) with air lines to introduce aeration in the media. When running
 339 multifactorial experiments (e.g., temperature and alkalinity), designing an analysis plan and concrete experimental
 340 questions to interrogate can help determine the sample size and minimum number of treatments.
 341

342 An analog to OAE is the use of lime soda and other alkali to combat acid rain, which has caused deleterious
 343 changes in freshwater ecosystems for more than half a century in northern Europe and North America. To reverse
 344 some of these' changes a number of governmental and nongovernmental teams have applied lime and other
 345 neutralizing compounds to streams, rivers, lakes, and catchments in the most affected or most ecologically valuable
 346 regions (see Clair and Hindar, 2005). Another example is the effects of seawater buffering mainly by addition of
 347 Na₂CO₃ addition) utilized by the commercial shellfish industry (e.g., Ragg et al., 2019), which showed a broad
 348 improvement in larval health compared to undersaturated waters.
 349
 350

351 Standardizing technical details in protocols, sampling, sample processing and analyses are crucial to control
352 for variation introduced by reagents, sample storage and other factors. The collection and curation of metadata
353 associated with each sample are critical for data interpretation, inter-lab comparison and drawing conclusions to
354 move forward with planning field deployments for research purposes. For studies involving more than one level of
355 biological organization; i.e., grazing experiments, competition experiments, particular attention should be paid to
356 designing adequate controls.

357
358 The effects of OAE and its interactions with other parameters might differ depending on the duration of the
359 experiments. Indeed, in ocean acidification studies, compensatory metabolic pathways appear to take longer to
360 become established, depending on factors such as the exposure history (Calosi et al., 2013) and phase of the life
361 cycle (Hettinger et al., 2012). In a study testing ocean acidification and warming, biological effects were not
362 detectable in the short term, but were rather manifested over time (Godbold and Solan, 2013). It was suggested that
363 species responses to seasonal variations in environmental conditions might explain these differences that, depending
364 upon timing, can either exacerbate or buffer the long-term directional effects of climatic forcing (Godbold and
365 Solan, 2013).

366 **4.c Recommended minimum set of variables to report**

367 To improve comparability between future work, we recommend a minimum set of variables with the
368 understanding that more variables might be added as new results emerge (Table 2). We recommend to measure and
369 report at least the following variables (shown in bold in Table 2).

- 371 • At least alkalinity and one more parameter of the carbonate system must be measured to calculate key carbonate
372 chemistry parameters including bicarbonate and carbonate ions, CO₂, pH and the saturation state of CaCO₃
373 polymorphs. This information is critical to determine chemical alterations in the dissolved inorganic carbon
374 system as a result of alkalization.
- 375 • Resource availability (e.g., prey, dissolved inorganic nutrients, light) are needed to monitor the growth
376 conditions.
- 377 • Particulate organic carbon (POC), nitrogen (PON), phosphorous (POP) are required to learn about trends in
378 biomass production and stoichiometry.
- 379 • Basic physiological properties (respiration, photosynthesis) should be measured to inform biogeochemical
380 models and learn about biologically-mediated fluxes of elements.
- 381 • Some functional group-specific properties, particularly those involving mineral precipitation (calcification,
382 silicification) and those with environmental effects (e.g., toxin production) and with climate-relevant impacts
383 (nitrogen fixation/denitrification) in context specific cases.
- 384 • Size of offspring and fecundity rates can be used as indicators of transgenerational plasticity and adaptation to
385 alkalization.

386
387 Other variables are important in the exploration of specific questions such as how does seawater alkalization
388 affect biodiversity?; how does metal bioavailability change under increased pH?; what is the role of organic
389 alkalinity in coastal systems? The variables and protocols listed in this chapter is not exhaustive and only provides a
390 proxy sample largely based on the literature on climate impacts on marine systems and ocean acidification.

391
392
393
394
395
396
397
398
399
400
401
402
403
404

405
406
407
408
409
410

Table 2. Examples of responses to ocean alkalinity enhancement to be measured in experimental manipulation studies. Knowledge need (M=medium, H=high; measurement mode (MM=manual mode; S=sensor; SD=sensor in development). A minimum variable set is highlighted in bold. Selected references are provided as examples of protocols.

Type of response	Variable	Knowledge need	Measurement mode	Protocol reference
Chemical and environmental	Carbonate chemistry parameters {[HCO ₃ ⁻], [CO ₃ ²⁻], [CO ₂], pCO ₂ , Ω}	H	MM, S, SD	Dickson (2010); Bockmon and Dickson (2015)
	Dissolved organic matter	M	MM	Marañón et al. (2004); Sharp et al. (1995)
	Dissolved inorganic nutrients	H	MM	Worsfold et al. (2013)
	Resource availability (prey, light)	H	MM, S	Lawrence et al. (2017)
	Particulate organic matter (C, N, P)	H	MM	Verardo et al. (1990); Hilton et al. (1996); Pujo-Pay and Raimbault (1994); Fu et al. (2008)
	Trace metals (in solution and in aggregates)	M	MM	Guo et al. (2022); Hutchins et al. (2023)
	Biologically and biogeochemically relevant elements (e.g., Si, Mg:Ca)	M	MM	Brzezinski (1985); de Nooijer et al. (2017)
Physiological	Basic physiology (respiration, photosynthetic, growth rates; morphometric measurements)	H	MM, S	Iglesias-Rodriguez et al. (2008); Kelly et al. (2013); Farrell et al. (2009)
	Some functional group-specific physiology (e.g., calcification, silicification, nitrification/denitri	H	MM, S	Cohen et al. (2017); DeCarlo et al. (2019)

	fication, toxin production)			
	Physiological stress [e.g., heat shock proteins, oxidative stress-related proteins, photosynthetic stress (shifts in quantum yield), morphological alterations (e.g., cyst formation)	M	MM	O'Donnell et al. (2009); Moya et al. (2015); Trimborn et al. (2017)
	Incidence of pathogens and disease	H	MM	Asplund et al. (2014)
Reproduction	Spawning success	M	MM	Liu et al. (2011)
	Size of offspring	M	MM	Cao et al. (2018); Johnson (2022); Albright et al. (2010)
	Sperm motility	M	MM	Esposito et al. (2020); Havenhand et al. (2008)
	Epigenetic analysis	M	MM	Li et al. (2018); Lee et al. (2022)
	Fecundity	M	MM	Maranhão and Marques (2003); Thor and Dupont (2015)
	Hatching success	M	MM	Saigusa (1992)
Species interactions	Competition for resources	M	MM	Connell et al. (2013); Guo et al. (2022)
	Predation and species interactions	M	MM	Greatorex and Knights (2023); Bacus and Kelley (2023); Mitchell et al. (2023)

	Synergistic/antagonistic effects of other environmental parameters	M	MM, S	Gerhard et al. (2023); Khalil et al. (2023)
--	--	---	-------	---

411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461

4.d Type of experiments

Laboratory experiments can be designed to both address short term responses and to explore the longer-term adaptation to chronic exposure to enhanced alkalinity conditions. Filtered natural seawater should be used, when possible, in incubations unless artificial seawater is required (for example, when studying the effect of metal concentrations). Short term manipulations involve the use of batch, semi-continuous and continuous incubation experiments. In *batch incubation experiments*, all resources are provided at the beginning of the incubation, without further addition and sampling takes place during a short time period (hours, days, weeks). Only gases and alkali can be added during the course of the experiment. When biological processes are measured, a phase during the life cycle (e.g., larva/adult; vegetative cells/gametes) or growth (healthy, exponentially-growing/resource-limited, stationary growing organisms /senescent organisms) is typically targeted. Sampling is conducted until the nutrients are consumed and beyond if decaying populations are the focus of the investigation.

Given that resources (light, nutrients) are the limiting factor in batch incubation experiments, the organisms are in the exponential growth phase for a limited time period. To expand sampling and replication during the exponential growth phase, resupply of nutrients using a *semi-continuous culturing* approach can prevent food/nutrients from becoming a limiting factor. When the studied organism is phototrophic, one must ensure subculturing (microbial cultures) or appropriate arrangement of organisms to prevent light limitation. The advantage of semi-continuous culturing is that it allows investigating trends over extended time periods, increase replication and higher yield. Generally, the resource is added manually or pumped from the nutrient supply vessel into the culture vessel during exponential growth or when specific conditions are met (e.g., when a certain biomass concentration is reached).

In *continuous cultures*, the rate of addition of nutrient is controlled to maintain steady state cell growth. This system is known as chemostat, where typically, a volume of culture medium is added and the same volume is removed from the growing culture. A challenge with this type of ‘bioreactors’ is that, over long time periods, they can be more susceptible to microbial contamination and long-term phenotypic and genotypic variance in the cultures (Reusch, 2013).

Portable incubation experiments that simulate regional in situ alkalinity deployments are an important step in understanding seawater alkalization and its impact on marine organisms prior to field testing. This type of incubation experiments, which simulate alkalinity additions under diverse local in situ parameters (e.g., temperature, irradiance, nutrients), can be accomplished using portable incubators onboard research vessels (i.e., deck incubations) or outdoors, at coastal research facilities (Fig. 2).

When studying photosynthetic organisms high-quality light filters should be attached to the acrylic tank to adjust photosynthetically active radiation (PAR) within the incubator (e.g., Fig. 2). To maintain in situ seawater temperatures, an inflow port can supply seawater to the incubator. Effort should be taken to ensure movement of seawater quickly through the incubator to maintain a uniform temperature.

When collecting natural seawater, one must consider how biological interactions (e.g., grazing) could confound results and filter accordingly. Unlike laboratory experiment, that allow for seawater-air phase CO₂ equilibration, portable incubation experiments require instantaneous alkalinity additions; thus, careful consideration should be given to the method of alkalinity addition used. When adding liquid alkalinity, e.g., solutions (e.g., 1 M) of NaOH one must consider that flocculation commonly occurs upon alkalinity addition (Subhas et al., 2022). Adding pulverized minerals directly to the treatment vessels is another option although this method may yield incomplete dissolution or slow dissolution (e.g., Fuhr et al., 2022) with undesirable effects including secondary precipitation, particle aggregation and detrimental biological impacts (NASEM, 2022). Some researchers have opted for mimicking mineral dissolution instead (see Gately et al., 2023). As in the traditional laboratory experiments described above, vessels within the incubator should ideally be aerated during experimentation. In addition to

462 chemical and biological parameters, PAR and temperature data should be collected throughout the experimental
463 timeframe through discrete sampling or semi-continuously using sensors and data loggers. The best practices
464 outlined in Box 1 should be adhered to when planning portable incubation experiments. Effort should be taken to
465 position the incubator in a way that avoids confounding factors such as light contamination (e.g., from the ship).

466 **5. Sampling and analysis**

467 Technical variability amongst experimental methods ranging from sampling and sample processing can
468 propagate through the various steps before analysis; for example, chemical analysis and molecular work/sequencing
469 can be error-prone (e.g., Catlett et al., 2020). The use of blanks every time sampling is conducted is essential for
470 detecting contamination originating from the experiment itself or from the adjacent environment (e.g., exogenous
471 sources such as surface contamination, flagellates in droplets through aeration, etc.). When possible, several barriers
472 to contamination are recommended (e.g., filters at various points of aeration). Additionally, for samples (other than
473 those preserved for analysis of alkalinity, dissolved inorganic carbon analysis or pH) that are kept for further
474 analyses, contaminants that grow during shipping or while samples are being stored can sometimes be reduced by
475 freezing at -80 °C, when possible, or by using the appropriate preservatives when storing at ambient temperature is
476 required (e.g., ethanol, paraformaldehyde, glutaraldehyde). Attention should be paid to the material of vessels where
477 samples and solutions are stored; for example, avoid borosilicate bottles to store nutrients or alkalinity solutions as
478 silicate can be leached into solution.

481 Establishing time series prior to the experiment to determine time frames regarding the appropriate length
482 of the experiment and frequency of sampling is recommended. It is important to establish well defined time
483 windows for sampling as well as frequency of sampling to capture physical, chemical and biological properties of
484 the studied system. It is advisable to limit the time of sample storage to minimize observations that might confound
485 interpretation of results (e.g., reverse weathering during storage) (Subhas et al., 2022).

487 **5.a Criteria for key parameters**

488 For the most part, laboratory experiments are aimed at elucidating the physiological performance and
489 biogeochemical responses of organisms (rather than communities) to physical or chemical alterations in the
490 environment although responses in ecological fitness could be drawn from laboratory experiments (Table 2).
491 Importantly, environmental change can affect species differently and interactions between species that are sensitive
492 to environmental change can function as ecological leverage points through which modest changes in abiotic
493 conditions are amplified into large changes in marine ecosystems (see Kroeker and Sanford, 2022). These
494 interactions can be measured as competition, predation, and symbiotic relationships (mutualism, commensalism and
495 parasitism) that can vary along environmental gradients that cause stress (Stachowick, 2001; Bruno et al., 2003; Ma
496 et al., 2023).

497 Criteria for selection of species should include whether the organism is amenable to laboratory
498 experimentation, the amount of background knowledge on the organism's physiology and biogeochemistry,
499 ecological importance of the organism, and local and global impacts. Considerations when selecting organisms
500 should also include geographic origin (e.g., temperate/tropical/polar) and ecosystem type (e.g., benthic vs pelagic).
501 Special attention should be paid to those species that (1) significantly impact or respond biogeochemically to
502 chemical changes caused by alkalinity addition (e.g., possibly calcifiers, photosynthetic organisms); (2) keystone
503 organisms (e.g., corals, salmon, sea stars, toxin-producing phytoplankton); and (3) organisms/functional groups of
504 known vulnerability to climate change (corals, urchins).

505 Calcium carbonate producing organisms are particularly interesting because of their known sensitivity to
506 changes in carbonate chemistry and because any alteration in their abundance or calcification rates could have
507 implications in the CDR potential of alkalization. Mineralogical composition of carbonate containing organisms
508 might possibly be affected by alkalization. For example, recent meta-analysis of studies exploring the effects of
509 the carbonate chemistry shifts caused by ocean acidification revealed effects on shell state, development and growth
510 rate (Figuerola et al., 2021). Biomineralization studies should explore species-specific responses driven by
511 mineralogical composition (calcite, aragonitic, high/low Mg calcite) of their tests, shells and skeletons.
512 Environmental and biological control on calcification particularly any changes in the Mg content in calcite driven by
513 the use of brucite and other minerals potentially adding Mg to calcite must be reported as calcite with a high Mg
514 content is less stable in aqueous solutions (Ries et al., 2016). Empirical studies have shown that the Mg/Ca ratio of
515
516
517

518 Mg-calcite producing organisms generally varies proportionally with seawater Mg/Ca (e.g., Ries, 2004; Ries, 2006)
519 and therefore particular attention should be paid to the Mg content (and solubility) of biomineralized calcite. The
520 addition to proposed Ca and Mg containing minerals - Ca(OH)₂ (slaked lime), Mg(OH)₂ (brucite), CaCO₃
521 (limestone) or (Mg,Ca)CO₃ (dolomite) - will alter the Mg/Ca ratio of the seawater. An extensive body of literature
522 reports biogenic and abiotic precipitation of low-Mg calcite when seawater Mg/Ca falls within the calcite stability
523 field (seawater molar Mg/Ca < 2) and the biogenic and abiogenic precipitation of aragonite and high-Mg calcite
524 when seawater Mg/Ca falls within the aragonite stability field (seawater molar Mg/Ca > 2) (Ries, 2010). Thus,
525 modification of local seawater Mg/Ca ratios by OAE has the potential to favor aragonite and high-Mg calcite
526 organisms if seawater Mg/Ca is increased, and low-Mg calcite organisms if seawater Mg/Ca is decreased. This is an
527 important area of future OAE research.

528
529 Central to OAE laboratory experimentation is our ability to measure any possible stress induced by
530 alkalization and learn about underlying mechanisms behind acclimation to the chemical alterations of seawater
531 caused by OAE. This can be achieved by measuring basic functions (growth rates, size, reproductive success),
532 sensitivities to alkalization might be organism-specific and possibly trophic level-specific (e.g., Voigt et al. 2003,
533 Gilman et al. 2010) although most laboratory experiments do not address the complexity of trophic interactions.
534 Similarly, measuring adaptation and diversity in acclimation between and within related organisms is a challenge
535 and the ocean acidification literature revealed how important it is to pay attention to diversity of responses (see
536 Kroeker et al., 2010).

537
538 Stress is often measured as a reduction in organismal performance or fitness caused by environmental
539 change (Schulte, 2014). In addition to these general physiological or behavioral responses, markers of stress such as
540 oxidative stress are often used. For example, it is well established that the production of reactive oxygen species
541 (ROS) can increase due to environmental stress including ocean acidification (Lesser, 2006; Lushchak, 2011). Many
542 biomarkers are commonly used for studying oxidative stress in marine organisms (Cailleaud et al., 2007; Vehmaa et
543 al., 2013) and an increase in ROS and superoxide dismutase and catalase activities have been reported in marine
544 animals under stress (von Weissenberg et al., 2022). Heat shock proteins (HSPs) are also used as molecular markers
545 of stress because of their abundance, high sensitivity to stress and being ubiquitously expressed (Gross, 2004).
546 Among all HSPs, HSP70s are the most studied as a strong up-regulation of HSP70 production has been
547 demonstrated broadly with the exception of *Hydra oligactis* (Bosch et al., 1998), and some Antarctic animals (La
548 Terza et al. 2001; Place and Hofmann, 2005).

549 **5.b Measurements of nutrient uptake rates**

551 The uptake rate of carbon and other nutrients that results in the observed standing stocks of particulate
552 matter involve many physiological processes that are sensitive to changes in inorganic carbon chemistry and pH
553 (Matsumoto et al., 2020). Chemical changes following the addition of alkalinity might alter physiological processes
554 that represent sources (calcification, respiration) and sinks (photosynthesis) of CO₂. One should also pay attention to
555 the reciprocal interactions between these physiological processes and the chemically altered environment as even
556 minor changes in biological processes, or in the balance between them, can have implications for the CDR potential
557 and biodiversity.

558 One of the most unknown effects of OAE is the fate of biological fixation rates of different elements (e.g.,
559 carbon and N₂ fixation rates). Such rates are measured in batch cultures and bioassay (mixed natural community)
560 incubation experiments (LaRoche et al., 2010). While the objective of culture experiments is to understand the effect
561 of environmental parameters on the elemental uptake by particular species in a lab, bioassay experiments have to
562 deal with a rather complex species interaction in the field or after subsampling of mesocosms in a lab (Hutchins et
563 al., 2007; Paul et al., 2016). Labelled/enriched (~99%) stable isotope tracers is the most used method for rate
564 estimation these days. The rate calculation is based on isotopic mass balance equation (Montoya et al., 1996):

$$565 \quad \text{C or N}_2 \text{ fixation rate} = \frac{[POM]}{t} \left(\frac{A_f - A_0}{A_e - A_0} \right) \quad (1)$$

566 where, [POM] is the concentration of element of interest (C or N) at the end of the incubation. Likewise, A_f
567 = atom% in POM at the end of incubation, A_0 = atom% in POM at the start of the incubation, t is time of incubation,
568 and A_e = isotopic enrichment in the dissolved form after the tracer addition at the start of the incubation

569 This equation/method is sensitive to analytical protocols in routine incubations (White et al., 2020), and
570 might be even more sensitive in OAE incubations due to the issue of gas equilibration in tightly capped bottles. While
571 the C substrate-based incubations are supposedly straightforward in incubations, N₂ gas incubations face the challenge
572 of under-equilibration leading to underestimation of rates. But OAE incubations can produce larger errors in the C
573 fixation estimates as well. This is because NaHCO₃ is generally used as a C substrate. To estimate ¹³C isotopic
574 enrichment after tracer addition (term in equation 1), a DIC value is normally assumed (as it does not change much at
575 a given region). But OAE is expected to increase (or fluctuate) DIC during the experimental period, and thus a
576 measured DIC value should be used in the enrichment factor calculation. Likewise, the ¹⁴C-method, which is widely
577 used for marine primary production and calcification rate measurements due to its sensitivity (Nielsen, 1952), also
578 requires treatment-specific determination of DIC concentrations. Likewise, slow dissolution of N₂ gas poses a
579 challenge to accurately estimating isotopic enrichment factor (*A_e*), and it is advisable to measure this term.

580 Although the analytical precision of C and N isotopes is of order of sub permil levels, many times the low reported
581 rates (<0.1 nmol N L⁻¹ d⁻¹) are questionable (Gradoville et al., 2017). Therefore, the detection limit of rate
582 measurements and its proper reporting is a major concern. To overcome this, following the propagation of analytical
583 and statistical errors in each term of mass balance equation (1), Gradoville et al. (2017) have proposed to report
584 minimal quantifiable rates (MQR) and the limit of detection (LOD) in triplicate samples. We ought to follow these
585 protocols in the rates measured in OAE. In addition, we must make sure to sample/filter sufficient water to achieve
586 35 µg N and 150 µg C in the sample for reliable mass spectrometric measurements.

587
588
589
590

591

592
593
594



595
596
597 **Figure 2.** A, B: Portable incubator with blue filters to adjust photosynthetically active radiation (PAR). A scalar
598 PAR sensor (LI-COR) can be observed within the incubator (A, right side). C: laboratory experiment using aeration
599 and sacrificial replication. Images were taken by James Gately (A, C) and Sylvia Kim (B).

600
601 **6. Conclusions and recommendations**

602 The field of OAE faces a great diversity of challenges given the continuously evolving experimental
603 approaches and emerging data availability that will undoubtedly provide new information and ideas to optimize best
604 practice in laboratory experimentation. This chapter highlights the need for attention to the design, sampling,
605 performance, and analysis of laboratory procedures used in OAE laboratory experiments. The criteria we present to
606 achieve best practice in laboratory experimentation and design focus on reproducibility, factors affecting CDR
607 potential and organism health (e.g., alkalinity conditions leading to flocculation, aggregation), establishing suitable
608 experimental controls, and identifying the appropriate level of biological organization (physiological, molecular) to
609 study biotic responses to OAE. Key response variables informing on alterations in seawater chemistry following
610 alkalization, growth of organisms /biomass buildup/reproductive success, and biogeochemically relevant
611 properties (e.g., photosynthesis, respiration, calcification) under elevated alkalinity conditions should be measured
612 and reported. The main recommendations include:

- 613
- Ensure reproducibility through appropriate experimental design and replication.
 - Determine alkalinity thresholds for the formation of precipitates for each experimental approach and condition.
 - In addition to the proposed alkalinity target values of 3000-4000 $\mu\text{mol kg}^{-1}$ (Renforth and Henderson, 2017), use concentrations exceeding these recommended values to mimic responses at the site of deployment/non equilibrium and use intermediate alkalinity levels to identify potential nonlinear responses.
 - Establish appropriate experimental design to address questions at specific levels of organization (chemical, physiological, molecular) and assuming different scenarios (e.g., mimicking impacts at the site of deployment in a non-equilibrated system versus steady state scenarios in an equilibrated system).

622

623 Given the emerging nature of ocean alkalinity enhancement as a research field, this chapter will evolve to
624 update guidelines as more results become publicly available. Frequent assessments of knowledge acquired from
625 emerging and future studies and review of best practices are needed to keep the OAE community engaged and
626 forward thinking.

627 Author contribution

628 M. Débora Iglesias-Rodriguez wrote the original manuscript with contributions from the authors. Rosalind E.M.
629 Rickaby, Arvind Sing and James A. Gately contributed with the critical review, commentary and revision of sections
630 and general comments on the manuscript.

633 Competing interests

634 Competing interests are declared in a summary for the entire volume at:
635 [url{https://sp.copernicus.org/articles/sp-bpoae-ci-summary.zip}](https://sp.copernicus.org/articles/sp-bpoae-ci-summary.zip).

638 Acknowledgements

639

640 This is a contribution to the “Guide for Best Practices on Ocean Alkalinity Enhancement Research”. We
641 thank our funders the ClimateWorks Foundation and the Prince Albert II of Monaco Foundation. Thanks are also
642 due to the Villefranche Oceanographic Laboratory for supporting the lead authors' meeting in January 2023.

644 References

645

646 Albright, R., Mason, B., Miller, M. and Langdon, C. (2010) Ocean acidification compromises recruitment
647 success of the threatened Caribbean coral *Acropora palmata*. PNAS 107, 20400–20404.

648

649 Asplund, M.E., Baden, S.P., Russ, S., Ellis, R.P., Gong and N., Hernroth, B.E. (2014) Ocean acidification
650 and host–pathogen interactions: blue mussels, *Mytilus edulis*, encountering *Vibrio tubiashii*. Environmental
651 Microbiology 16, 1029–1039.

652

653 Bach, L.T., Taucher, J., Boxhammer, T., Ludwig, A., The Kristineberg KOSMOS Consortium, Achterberg,
654 E.P., Algueró-Muñiz, M., Anderson, L.G., Bellworthy, J., Büdenbender, J., Czerny, J., Ericson, Y., Esposito, M.,
655 Fischer, M., Haunost, M., Hellemann, D., Horn, H.G., Hornick, T., Meyer, J., Sswat, M., Zark, M. and Riebesell, U.
656 (2016) Influence of ocean acidification on a natural winter-to-summer plankton succession: first insights from a
657 long-term mesocosm study draw attention to periods of low nutrient concentrations. PLoS ONE 11(8): e0159068.
658 doi:10.1371/journal.pone.0159068.

659

660 Bacus, S.C., Kelley, A.L. (2023) Effects of ocean acidification and ocean warming on the behavior and
661 physiology of a subarctic, intertidal grazer. Mar Ecol Prog Ser 711, 31-45.

662

663 Bockmon, E.E. and Dickson, A.G., 2015 An inter-laboratory comparison assessing the quality of seawater
664 carbon dioxide measurements. Marine Chemistry 171, 36-43.

665

666 Bosch, T.C.G., Krylow, S.M., Bode, H.R. and Steele, R.E. (1988) Thermotolerance and synthesis of heat-
667 shock proteins—these responses are present in *Hydra attenuata* but absent in *Hydra oligactis*. Proceedings of the
668 National Academy of Sciences U.S.A. 85, 7927–7931.

669
670 Brantley, S.L. (2008) Kinetics of mineral dissolution. In S.L. Brantley, J.D. Kubicki, A.F. White (Eds.),
671 Kinetics of Water–Rock Interactions, Springer Science (2008), pp. 151-210

672
673 Bruno, J.F., Stachowicz, J.J., Bertness, M.D. (2003) Inclusion of facilitation into ecological theory. Trends
674 in Ecology and Evolution 18: 119–25.

675
676 Brzezinski, M.A. (1985) The Si:C:N ratio of marine diatoms: interspecific variability and the effect of
677 some environmental variables. J Phycol. 21, 347–357.

678
679 Byrne, M. and Przeslawski, R. (2013) Multistressor studies of the impacts of warming and acidification of
680 the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology*, **53**, 582–596.

681
682 Cailleaud, K., Maillet, G., Budzinski, H., Souissi, S., & Forget-Leray, J. (2007). Effects of salinity and
683 temperature on the expression of enzymatic biomarkers in Eurytemora affinis (Calanoida, Copepoda). Comparative
684 Biochemistry and Physiology - A Molecular and Integrative Physiology, 147, 841–849.

685
686 Cao, R., Wang, Q., Yang, D, Liu, Y., Ran, W., Qu Y, Wu H, Cong M, Li F, Ji C, Zhao J. (2018) CO₂-
687 induced ocean acidification impairs the immune function of the Pacific oyster against *Vibrio splendidus* challenge:
688 An integrated study from a cellular and proteomic perspective. Science of The Total Environment 625, 1574-1583.
689

690 Calosi, P., Rastrick, S.P.S., Lombardi, C., de Guzman, H.J., Davidson, L., Jahnke, M., Giangrande, A.,
691 Hardege, J.D., Schulze, A., Spicer, J.I. and Gambi, M-C. (2013) Adaptation and acclimatization to ocean
692 acidification in marine ectotherms: an in situ transplant experiment with polychaetes at a shallow CO₂ vent system.
693 Phil. Trans. R. Soc. B 368, 20120444.

694
695 Casadevall, A., and Fang, F.C. (2010) Reproducible Science. Infection and Immunity 78, 4972–4975.

696
697 Catlett, D., Matson, P.G., Carlson, C.A., Wilbanks, E.G., Siegel, D.A. and Iglesias-Rodriguez, M.D. (2020)
698 Evaluation of accuracy and precision in an amplicon sequencing workflow for marine protist communities.
699 Limnology and Oceanography Methods 18, 20-40.

700
701 Clair, T.A. and Hindar, A. (2005) Liming for the mitigation of acid rain effects in freshwaters: A review of
702 recent results. Environ. Rev. 13, 91–128.

703
704 Cohen, S., Krueger, T., Fine, M. (2017) Measuring coral calcification under ocean acidification:
705 methodological considerations for the ⁴⁵Ca-uptake and total alkalinity anomaly technique. PeerJ 5:e3749.

706
707 Connell, S.D., Kroeker, K.J., Fabricius, K.E., Kline, D.I., Russell, B.D. (2013) The other ocean
708 acidification problem: CO₂ as a resource among competitors for ecosystem dominance. Phil Trans R Soc B 368:
709 20120442.<http://dx.doi.org/10.1098/rstb.2012.0442>.

710
711 Cornwall, C.E. and Hurd, C.L. (2016) Experimental design in ocean acidification research: problems and
712 solutions. ICES Journal of Marine Science 73(3): 572–581.

713
714 Crain CM, Kroeker K, Halpern BS. 2008 Interactive and cumulative effects of multiple human stressors in
715 marine systems. Ecol. Lett. 11, 1304-1315.

716
717 Danger, M., Oumarou, C., Benest, D. and Lacroix, G. (2007) Bacteria can control stoichiometry and
718 nutrient limitation of phytoplankton. Functional Ecology 21, 202-210.

719
720 Darling ES, Côté IM. 2008 Quantifying the evidence for ecological synergies. Ecol. Lett. 11, 1278-1286.

721

722 Davis, P.S. (1989) Short-term growth measurements of corals using an accurate buoyant weighing
723 technique. *Marine Biology* 101,389-395.
724

725 DeCarlo, T.M., Comeau, S., Cornwall, C.E., Gajdzik, L., Guagliardo, P., Sadekov, A., Thillainath, E.C.,
726 Trotter, J. and McCulloch, M.T. (2019) Investigating marine bio-calcification mechanisms in a changing ocean with
727 in vivo and high-resolution ex vivo Raman spectroscopy. *Glob Change Biol.* 2019; 25: 1877– 1888.
728

729 de Nooijer, L.J., van Dijk, I., Toyofuku, T., and Reichart, G.J. (2017). The impacts of seawater Mg/Ca and
730 temperature on element incorporation in benthic foraminiferal calcite. *Geochemistry, Geophysics, Geosystems* 18,
731 3617–3630.
732

733 Dickson, A. G. (2010) The carbon dioxide system in seawater: Equilibrium chemistry and measurements.
734 In U. Riebesell, V. J. Fabry, & L. Hansson (Eds.), *Guide to best practices for ocean acidification research and data*
735 *reporting* (pp. 17–40). Luxembourg: Publications Office of the European Union.
736

737 Dodge, R.E. Wyers, S.C., FRITH, H.R., Knap, A.H., Smith, S.R., Cook, C.B. and Sleeter, T.D. (1984)
738 Coral calcification rates by the buoyant weight technique—Effects of alizarin staining *J. Exper. Mar. Biol. Ecol.*, 75,
739 217–232.
740

741 Dupont, C.L., Barbeau, K., and Palenik, B. (2008) Ni uptake and limitation in marine
742 *Synechococcus* strains, *Appl. Environ. Microb.*, 74, 23-31.
743

744 Dupont, C. L., Buck, K.N., Palenik, B., and Barbeau, K. (2010) Nickel utilization in phytoplankton
745 assemblages from contrasting oceanic regimes, *Deep-Sea Res. Pt. II* , 57, 553-566.
746

747 Elser, J.J., Dobberfuhl, D.R., MacKay, N.A. and Schampel, J.H. (1996) Organism size, life history, and
748 N:P stoichiometry: toward a unified view of cellular and ecosystem processes. *BioScience* 46: 674-684.
749

750 Esposito, M.C., Boni, R., Cuccaro, A., Tosti, E. and Gallo, A. (2020) Sperm Motility Impairment in Free
751 Spawning Invertebrates Under Near-Future Level of Ocean Acidification: Uncovering the Mechanism. *Front. Mar.*
752 *Sci.* 6:794. doi: 10.3389/fmars.2019.00794.
753

754 Farrell, A.P., Eliason, E.J. Sandblom, E., Clark, T.D. (2009) Fish cardiorespiratory physiology in an era of
755 climate change. *Canadian Journal of Zoology* 87(10), 835-851.
756

757 Ferderer, A., Chase, Z., Kennedy, F., Schulz, K.G. and Bach, L.T. (2022) Assessing the influence of ocean
758 alkalinity enhancement on a coastal phytoplankton community. *Biogeosciences* 19, 5375–5399.
759

760 Figuerola, B., Hancock, A.M., Bax, N., Cummings, V.J., Downey, R., Griffiths, H.J., Smith, J. and Stark,
761 J.S. (2021) A review and meta-analysis of potential impacts of ocean acidification on marine calcifiers from the
762 Southern Ocean. *Front. Mar. Sci.* 8:584445. doi: 10.3389/fmars.2021.584445.
763

764 Forbes, V. and Calow, P. (2002) Population growth rate as a basis for ecological risk assessment of toxic
765 chemicals. *Philos Trans R Soc Lond B Biol Sci.* 357, 1299–1306.
766

767 Fuhr, M., Geilert, S., Schmidt, M., Liebetrau, V., Vogt, C., Ledwig, B., Wallmann, K. (2022) Kinetics of
768 olivine weathering in seawater: an experimental study. *Frontiers in Climate* 4: 831587. doi:
769 10.3389/fclim.2022.831587.
770

771 Galic N, Sullivan LL, Grimm V, Forbes VE. 2018 When things don't add up: quantifying impacts of
772 multiple stressors from individual metabolism to ecosystem processing. *Ecol. Lett.* 21, 568-577.
773

774 Gerhard, M., Koussoroplis, A.M., Raatz, M., Pansch, C., Fey, S.B., Vajedsamiei, J., Calderó-Pascual, M.,
775 Cunillera-Montcusí, D., Juvigny-Khenafou, N.P.D., Polazzo, F., Thomas, P.K., Symons, C.C., Beklioğlu, M.,
776 Berger, S.A., Chefaoui, R.M., Ger, K.A., Langenheder, S., Nejstgaard, J.C., Ptacnik, R. and Striebel, M. (2023)

777 Environmental variability in aquatic ecosystems: Avenues for future multifactorial experiments. *Limnology and*
778 *Oceanography Letters* 8, 247–266.

779

780 Gately, J.A., Kim, S.M., Jin, B., Brzezinski, M.A. and Iglesias-Rodriguez, M.D. (2023) Coccolithophores
781 and diatoms resilient to ocean alkalinity enhancement: a glimpse of hope? *Science Advances* 9(24) . doi:
782 10.1126/sciadv.adg6066.

783

784 Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W., Holt, R.D. (2010) A framework for community
785 interactions under climate change. *Trends in Ecology and Evolution* 25, 325–31.

786

787 Godbold, J.A., Solan, M. (2013) Long-term effects of warming and ocean acidification are modified by
788 seasonal variation in species responses and environmental conditions. *Philos Trans R Soc Lond B Biol Sci.*
789 26;368(1627):20130186.

790

791 Gradoville, M.R., Bombar, D., Crump, B.C., Letelier, R.M., Zehr, J.P., and White, A.E. (2017). Diversity
792 and activity of nitrogen-fixing communities across ocean basins. *Limnology and Oceanography*, 62(5), 1895-1909.

793

794 Greatorex, R. and Knights, A.M. (2023) Differential effects of ocean acidification and warming on
795 biological functioning of a predator and prey species may alter future trophic interactions. *Marine Environmental*
796 *Research* 186, 105903.

797

798 Gross, M. (2004) Emergency services: a bird’s eye perspective on the many different functions of stress
799 proteins. *Current Protein and Peptide Science* 5, 213–223.

800

801 Guillard, R.R. and Ryther, J.H. (1962) Studies of marine planktonic diatoms. I. *Cyclotella nana* (Hustedt),
802 and *Detonula confervacea* (Cleve Gran.). *Can J Microbiol*, 8, 229-239.

803

804 Guo, J.A., Strzepak, R., Willis, A., Ferderer, A. and Bach, L.T. (2022) Investigating the effect of nickel
805 concentration on phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement.
806 *Biogeosciences* 19, 3683–3697.

807

808 Hartmann, J., Suitner, N., Lim, C., Schneider, J., Marín-Samper, L., Arístegui, J., Renforth, P., Taucher, J.
809 and Riebesell, U. (2023) Stability of alkalinity in ocean alkalinity enhancement (OAE) approaches – consequences
810 for durability of CO₂ storage. *Biogeosciences*, 20, 781–802.

811

812

813 Harvey, B.P., Gwynn-Jones, D. and Moore, P.J. (2013). Meta-analysis reveals complex marine biological
814 responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution* 3, 1016–1030.

815

816 Havenhand, J.N., Buttler, F.R., Thorndyke, M.C. and Williamson, J.E. (2008) Near-future levels of ocean
817 acidification reduce fertilization success in a sea urchin. *Curr. Biol.* 18, R651–R652.

818

819 Hettinger, A., Sanford, E., Hill, T.M., Russell, A.D., Sato, K.N.S., Hoey, J., Forsch, M., Page, H.N.,
820 Gaylord, B. (2012) Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster.
821 *Ecology* 93, 2758–2768.

822

823 Hilton, J., Lishman, J.P., Mackness, S., Heaney, S.I. (1986) An automated method for the analysis of particulate
824 carbon and nitrogen in natural waters. *Hydrobiologia* 141, 269-271.

825

826 Ho, T.-Y. (2013) Nickel limitation of nitrogen fixation in *Trichodesmium*, *Limnol. Oceanogr.*, 58, 112-120.

827

828 Hu, X. (2020). Effect of organic alkalinity on seawater buffer capacity: A numerical exploration. *Aquatic*
829 *Geochemistry*. [https://doi.org/ 10.1007/s10498-020-09375-x](https://doi.org/10.1007/s10498-020-09375-x).

830

831 Hurlbert, S.H. (2009) The Ancient Black Art and Transdisciplinary Extent of Pseudoreplication. *Journal of*
832 *Comparative Psychology*, American Psychological Association 2009 4, 434–443.

833
834 Hutchins, D.A., Fu, F.X., Zhang, Y., Warner, M.E., Feng, Y., Portune, K., ... and Mulholland, M.R. (2007).
835 CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth rates, and elemental ratios: Implications for past,
836 present, and future ocean biogeochemistry. *Limnology and oceanography*, 52(4), 1293-1304.
837
838 Hutchins, D.A., Fu, F.X., Yang, S.C., John, S.G., Romaniello, S.J., Andrews, M.G. and Walworth, N.G.
839 (2023) Responses of globally important phytoplankton groups to olivine dissolution products and implications for
840 carbon dioxide removal via ocean alkalinity enhancement. bioRxiv, pp.2023-04.
841
842 Iglesias-Rodríguez, M.D., Halloran, P.R., Rickaby, R.E.M., Hall, I.R., Colmenero-Hidalgo, E., Gittins,
843 J.R., Green, D.R.H., Tyrrell, T., Gibbs, S.J., von Dassow, P., Rehm, E., Armbrust, E.V. and Boessenkool, K.P.
844 (2008) Phytoplankton calcification in a high CO₂ world. *Science* 320, 336–39.
845
846 IPCC. 2021. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the
847 Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Summary for Policymakers, V.
848 Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I.
849 Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, Yu R. and
850 B. Zhou, eds. Cambridge University Press.
851
852 Kelly, M.W., Padilla-Gamiño, Hofmann, G.E. (2013) Natural variation and the capacity to adapt to ocean
853 acidification in the keystone sea urchin. *Global Change Biology* 19, 2536–2546.
854
855 Khalil, M., Doo, S.S., Stuhr, M. and Westphal , H. (2023) Long-term physiological responses to combined
856 ocean acidification and warming show energetic trade-offs in an asterinid starfish. *Coral Reefs*.
857 <https://doi.org/10.1007/s00338-023-02388-2>
858
859 Kim, H.-C. and Lee, K. (2009) Significant contribution of dissolved organic matter to seawater alkalinity,
860 *Geophys. Res. Lett.*, 36, L20603, 5 pp..
861
862 Koeve, W., Kim, H.-C., Lee, K., and Oschlies, A. (2011) Potential impact of DOC accumulation on *f*CO₂
863 and carbonate ion computations in ocean acidification experiments, *Biogeosciences Discuss.*, 8, 3797–3827,
864 doi:10.5194/bgd-8-3797-2011.
865
866 Kroeker, K.J. and Sanford, E. (2022) Ecological leverage points: species interactions amplify the
867 physiological effects of global environmental change in the ocean. *Annual Review of Marine Science* 14: 75–103.
868
869 Kroeker, K.J., Kordas, R.L., Crim, R.N. and Singh, G.G. (2010) Meta-analysis reveals negative yet variable
870 effects of ocean acidification on marine organism. *Ecology Letters* 13, 1419-1434.
871
872 Kroeker, K.J., Kordas, R.L., Crim, R., Hendriks, I.E., Ramajo, L., Singh, G.S., Duarte, C.M. and Gattuso,
873 J.-P. (2013b) Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with
874 warming. *Global Change Biology*, 19, 1884–1896.
875
876 Kroeker, K., Micheli, F. & Gambi, M. (2013a) Ocean acidification causes ecosystem shifts via altered
877 competitive interactions. *Nature Clim Change* 3, 156–159.
878
879 LaRoche, J., Rost, B., and Engel, A. (2010). Bioassays, batch culture and chemostat experimentation. In
880 Approaches and tools to manipulate the carbonate chemistry., Guide for Best Practices in Ocean Acidification
881 Research and Data Reporting. In: Riebesell U., Fabry VJ, Hansson L., Gattuso J.-P.(Eds.), pp. (pp. 81-94).
882
883 LaTerza, A.L., Miceli, C. and Luporini, P. (2001) Divergence between two Antarctic species of the ciliate
884 *Euplotes*, *E. focardii* and *E. nobilii*, in the expression of heat-shock protein 70 genes. *Molecular Ecology* 10, 1061–
885 1067.
886

887 Lawrence, M.J., Eliason, E.J., Brownscombe, J.W., Gilmour, K.M., Mandelman, J.W. and Cooke, S.J.
888 (2017) An experimental evaluation of the role of the stress axis in mediating predator-prey interactions in wild
889 marine fish. *Comparative Biochemistry and Physiology, Part A* 207, 21-29.
890

891 Lesser, M.P. (2006) Oxidative stress in marine environments: Biochemistry and physiological ecology.
892 *Annual Review of Physiology* 68, 253–278.
893

894 Lebrato, M., Garbe-Schönberg, D., Müller, M.N., Blanco-Ameijeiras, S., Feely, R.A. Lorenzoni, L.,
895 Molinero, J.C., Bremer, K., Jones, D.O.B., Iglesias-Rodriguez, D., Greeley, D., Lamare, M.D., Paulmier, A., Graco,
896 M., Cartes, J., Barcelos e Ramos, J., de Lara, A., Sanchez-Leal, R., Jimenez, P., Paparazzo, F.E., Hartman, S.E.,
897 Westernströer, U., Küter, M., Benavides, R., da Silva, A.F., Bell, S., Payne, C., Olafsdottir, S., Robinson, K.,
898 Jantunen, L.M., Korablev, A., Webster, R.J., Jones, E.M., Gilg, O., Bailly du Bois, P., Beldowski, J., Ashjian, C.,
899 Yahia, N.D., Twining, B., Chen, X.G., Tseng, L.C., Hwang, J.S., Dahms, H.U., Oschlies, A. (2020) Global
900 variability in seawater Mg:Ca and Sr:Ca ratios in the modern ocean. *Proc. Natl. Acad. Sci. U.S.A.* 117, 22281–
901 22292.
902

903 Lee, Y.H., Kim, M.S., Wang, M., Bhandari, R.K., Park, H.G., Wu, R.S-S. and Lee, J.-S. (2022) Epigenetic
904 plasticity enables copepods to cope with ocean acidification. *Nat. Clim. Chang.* 12, 918–927.
905

906 Li, Y., Lieu, Y.J., Cui, G., Cziesielski, M.J., Zahran, N., Michell, C.T., Voolstra, C.R. and Aranda, M.
907 (2018) DNA methylation regulates transcriptional homeostasis of algal endosymbiosis in the coral model *Aiptasia*.
908 *Sci. Adv.* 4, eaat2142.
909

910 Liu, G., Innes, D. and Thompson, R.J. (2011) Quantitative analysis of sperm plane circular movement in
911 the blue mussels *Mytilus edulis*, *M. trossulus* and their hybrids. *J. Exp. Zoology*, 315, 280-290.
912

913 Lushchak, V.I. (2011) Environmentally induced oxidative stress in aquatic animals. *Aquatic Toxicology*
914 101, 13–30.
915

916 Ma, K.C.K., Monsinjon, J.R., Froneman, P.W., McQuaid, C.D. (2023) Thermal stress gradient causes
917 increasingly negative effects towards the range limit of an invasive mussel. *Science of the Total Environment* 865,
918 (2023), Article 161184.
919

920 Maranhão, P. and Marques João, C. (2003) The influence of temperature and salinity on the duration of
921 embryonic development, fecundity and growth of the amphipod *Echinogammarus marinus* Leach (Gammaridae).
922 *Acta Oecol.*, 24, pp. 5-13.
923

924 Marañón, E., Cermeño, P., Fernández, E., Rodríguez J., and Zabala, L. (2004) Significance and
925 mechanisms of photosynthetic production of dissolved organic carbon in coastal eutrophic ecosystems. *Limnol.*
926 *Oceanogr.* 49, 1652– 1666.
927

928

929 Martin, B., Jager, T., Nisbet, R.M., Preuss, T.G., Grimm, V. (2014) Limitations of extrapolating
930 toxic effects on reproduction to the population level. *Ecol Appl.* 24, 1972–1983.
931

932 Martz, T.R., Daly, K.L., Byrne, R.H., Stillman, J.H. and Turk, D. (2015) Technology for Ocean
933 Acidification Research: Needs and Availability. *Oceanography* 28 (2), Special Issue on Emerging Themes in Ocean
934 Acidification Science: 40-47.
935

936 Matsumoto, K., Tanioka, T. and Rickaby, R. (2020) Linkages between dynamic phytoplankton C: N: P and
937 the ocean carbon cycle under climate change. *Oceanography* 33, 44–52.
938

939 Middelburg, J.J., Soetaert, K., and Hagens, M. (2020). Ocean alkalinity, buffering and biogeochemical
940 processes. *Reviews of Geophysics*, 58, e2019RG000681. [https://doi.org/ 10.1029/2019RG000681](https://doi.org/10.1029/2019RG000681).
941

942 Millero, F.J., Woosley, R., Ditrolio, B. And Waters, J. (2009) Effect of ocean acidification on the
943 speciation of metals in seawater. *Oceanography* 22, No. 4, Special Issue On The Future of Ocean Biogeochemistry
944 in a High-CO₂ World pp. 72-85.
945

946 Mitchell, A., Hayes, C., Booth, D.J. and Nagelkerken, I. (2023) Future shock: Ocean acidification and
947 seasonal water temperatures alter the physiology of competing temperate and coral reef fishes. *Science of the Total*
948 *Environment* 883, 163684.
949

950 Montoya, J.P., Voss, M., Kahler, P., and Capone, D.G. (1996). A simple, high-precision, high-sensitivity
951 tracer assay for N (inf2) fixation. *Applied and Environmental Microbiology*, 62(3), 986-993.
952

953 Montserrat, F., Renforth, P., Hartmann, J., Leermakers, M., Knops, P. and Meysman, F.J.R. (2017) Olivine
954 dissolution in seawater: implications for CO₂ sequestration through enhanced weathering in coastal environments.
955 *Environmental Science & Technology* 51, 3960-3972.
956

957 Moras, C. A., Bach, L. T., Cyronak, T., Joannes-Boyau, R., and Schulz, K. G. (2022) Ocean alkalinity
958 enhancement – avoiding runaway CaCO₃ precipitation during quick and hydrated lime dissolution, *Biogeosciences*,
959 19, 3537–3557.
960

961 Morel, F.M.M., Rueter, J., Anderson, D.M. and Guillard, R.R.L. (1979) Aquil: A chemically defined
962 phytoplankton culture medium for trace metal studies. *J. Phycol.* 15, 135– 141.
963

964 Moya, A., Huisman, L., Forêt, S., Gattuso, J.-P., Hayward, D.C., Ball, E.E. and Miller, D.J. (2015) Rapid
965 acclimation of juvenile corals to CO₂-mediated acidification by upregulation of heat shock protein and Bcl-2 genes.
966 *Mol Ecol*, 24: 438-452.
967

968 NASEM: A Research Strategy for Ocean-based Carbon Dioxide Removal and Sequestration, National
969 Academies of Sciences, Engineering, and Medicine, Washington (DC), <https://doi.org/10.17226/26278>, 2022.
970

971 Nawaz, S., Lezaun, J., Valenzuela, J.M. and Renforth, P. (2023). Broaden Research on Ocean Alkalinity
972 Enhancement to Better Characterize Social Impacts. *Environmental Science & Technology* 57, 24, 8863–8869.
973

974 Nielsen, S.E. (1952) The use of radio-active carbon (C¹⁴) for measuring organic production in the sea. *ICES*
975 *Journal of Marine Science* 18, 117–140.
976

977 O'Donnell, M.J., Hammond, L.M. and Hofmann, G.E. (2009) Predicted impact of ocean acidification on a
978 marine invertebrate: elevated CO₂ alters response to thermal stress in sea urchin larvae. *Mar Biol* 156, 439–446.
979

980 Paiva, F., Brennecke, D., Pansch, C. and Briski, E. (2021) Consistency of aquatic enclosed experiments:
981 The importance of scale and ecological complexity. *Diversity and Distributions* 27, 524–532.
982

983 Paul, A.J., Bach, L.T., Schulz, K.-G., Boxhammer, T., Czerny, J., Achterberg, E.P., Hellemann, D., Trense,
984 Y., Nausch, M., Sswat, M. and Riebesell, U. (2015) Effect of elevated CO₂ on organic matter pools and fluxes in a
985 summer Baltic Sea plankton community. *Biogeosciences* 12: 6181–6203.
986

987 Paul, A.J., Achterberg, E.P., Bach, L.T., Boxhammer, T., Czerny, J., Haunost, M., ... and Riebesell, U.
988 (2016). No observed effect of ocean acidification on nitrogen biogeochemistry in a summer Baltic Sea plankton
989 community. *Biogeosciences*, 13(13), 3901-3913.
990

991 Pisteos, J.C.A., Nagelkerken, I., Rossi, T. and Connell, S.D. (2016) Antagonistic effects of ocean
992 acidification and warming on hunting sharks. *OIKOS* 126,
993 <https://doi.org/10.1111/oik.03182>.
994

995 Place, S.P. and Hofmann, G.E. (2005) Constitutive expression of a stress-inducible heat shock protein gene,
996 hsp70, in a phylogenetically distant Antarctic fish. *Polar Biology* 28, 261–26.
997

998 Pujó-Pay, M. and Raimbault, P. (1994) Improvement of the wet-oxidation procedure for simultaneous
999 determination of particulate organic nitrogen and phosphorus collected on filters. *Mar. Ecol. Prog. Ser.* 105, 203-
1000 207.

1001 Ragg, N.L.C., Gale, S.L., Le, D.V., Hawes, N.A., Burritt, D.A., Young, T., Jessica A. Ericson, Zoë
1002 Hilton, Ellie Watts, Jolene Berry, Nick King (2019) The Effects of Aragonite Saturation State on Hatchery-Reared
1003 Larvae of the Greenshell Mussel *Perna canaliculus*

1004

1005 Reusch, T.B.H. (2014) Climate change in the oceans: evolutionary versus phenotypically plastic responses
1006 of marine animals and plants. *Evolutionary Applications* 7(2014) 104–122
1007 Reusch Adaptation and global change
1008 in the ocean. *Evolutionary Applications* 7, 104–122.

1009 Ridgwell, A., Schmidt, D.N., Turley, C., Brownlee, C., Maldonado, M.T., Tortell, P. and Young, J.R.
1010 (2009). From laboratory manipulations to Earth system models: scaling calcification impacts of ocean acidification.
1011 *Biogeosciences*, 6, 2611–2623.

1012

1013 Ries, J.B. (2004) The effect of ambient Mg/Ca on Mg fractionation in calcareous marine invertebrates: A
1014 record of Phanerozoic Mg/Ca in seawater. *Geology* 32, 981-984.

1015

1016 Ries, J.B. 2006. Mg fractionation in crustose coralline algae: Geochemical, biological, and
1017 sedimentological implications of secular variation in the Mg/Ca ratio of seawater. *Geochimica et Cosmochimica*
1018 *Acta* 70, 891-900).

1019

1020 Ries, J.B. (2010) Geological and experimental evidence for secular variation in seawater Mg/Ca (calcite-
1021 aragonite seas) and its effects on marine biological calcification. *Biogeosciences* 7, 2795-2849.

1022

1023 Ries, J.B., Cohen, A.L., McCorkle, D.C. (2009) Marine calcifiers exhibit mixed responses to CO₂-induced
1024 ocean acidification. *Geology*, 37, 1131-1134.

1025

1026 Ries, J.B., Ghazaleh, M.N., Connolly, B., Westfield, I., Castillo, K.D., 2016, Impacts of ocean acidification
1027 and warming on the dissolution kinetics of whole-shell biogenic carbonates. *Geochimica et Cosmochimica Acta*
1028 192: 318-337. doi: 10.1016/j.gca.2016.07.001.

1029

1030 Saigusa, M. (1992) Control of hatching in an estuarine terrestrial crab I. Hatching of embryos detached
1031 from the female and emergence of mature larvae. *Biol. Bull.*, 183, 401-408.

1032

1033 Sala, M.M., Aparicio, F.L., Balagué, V., Boras, J.A., Borrull, E., Cardelús, C., Cros, L., Gomes, A., López-
1034 Sanz, A., Malits, A., Martínez, R.A., Mestre, M., Movilla, J., Sarmiento, H., Vázquez-Domínguez, E., Vaqué, D.,
1035 Pinhassi, J., Calbet, A., Calvo, E., Gasol, J.M., Pelejero, C. and Marrasé, C. (2015) Contrasting effects of ocean
1036 acidification on the microbial food web under different trophic conditions. *Ices J Mar Sci.* 73: 670–679.

1037

1038 Sanders T, Schmittmann L, Nascimento-Schulze JC and Melzner F (2018) High Calcification Costs Limit
1039 Mussel Growth at Low Salinity. *Front. Mar. Sci.* 5:352. doi: 10.3389/fmars.2018.00352

1040

1041 Sharp, J.H., Benner, R., Bennett, L., Carlson, C.A., Fitzwater, S.E, Peltzer, E.T. and Tupas, L.M. (1995)
1042 Analyses of dissolved organic carbon in seawater: the JGOFS EqPac methods comparison. *Mar. Chem.* 48, 91–108.

1043

1044 Stachowick, J.J. (2001) Mutualism, Facilitation, and the Structure of Ecological Communities. *BioScience*
1045 51, 235–246.

1046

1047 Subhas, A.V., Marx, L., Reynolds, S., Flohr, A., Mawji, E.W., Brown, P.J. and Cael, B.B. (2022) Microbial
1048 ecosystem responses to alkalinity enhancement in the North Atlantic Subtropical Gyre. *Front. Clim.* 4:784997. doi:
1049 10.3389/fclim.2022.784997.

1050

1051 Thor, P. and Dupont, S. (2015) Transgenerational effects alleviate severe fecundity loss during ocean
1052 acidification in a ubiquitous planktonic copepod. *Global Change Biology* 21, 2261–2271.

1053

1054 Trimborn, S. Thoms, S., Brenneis, T., Heiden, J.P., Beszteri, S. and Bischof, K. (2017) Two Southern
1055 Ocean diatoms are more sensitive to ocean acidification and changes in irradiance than the prymnesiophyte
1056 *Phaeocystis antarctica*. *Physiologia Plantarum* 160, 155–170.
1057
1058 USGCRP (U.S. Global Change Research Program). 2017. Climate Science Special Report: Fourth National
1059 Climate Assessment, Vol. I, D. J. Wuebbles, D. W. Fahey, K. A. Hibbard, D. J. Dokken, B. C. Stewart and T. K.
1060 Maycock, eds. Washington, DC:USGCRP.
1061
1062 Vandamme, D., Pohl, P. I., Beuckels, A., Foubert, I., Brady, P. V., Hewson, J. C., Muylaert, K. (2015).
1063 Alkaline flocculation of *Phaeodactylum tricornutum* induced by brucite and calcite. *Bioresour. Technol.* 196, 656–
1064 661. doi: 10.1016/j.biortech.2015.08.042
1065
1066 Vehmaa, A., Hogfors, H., Gorokhova, E., Brutemark, A., Holmborn, T., & Engström-Öst, J. (2013)
1067 Projected marine climate change: Effects on copepod oxidative status and reproduction. *Ecology and Evolution* 3,
1068 4548–4557.
1069
1070 Verardo, D.J., Froelich, P.N. and McIntyre, A. (1990) Determination of organic carbon and nitrogen in
1071 marine sediments using the Carlo Erba NA-1500 analyzer. *Deep Sea Res. I* 37, 157-165.
1072
1073 Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bährmann, R., Fabian, B., Heinrich, W.,
1074 Köhler, G., Lichter, D., Marstaller, R. and Sander, F.W. (2003) Trophic levels are differentially sensitive to climate.
1075 *Ecology* 84, 2444–53.
1076
1077 von Weissenberg, E., A. Jansson, K. A. Vuori, and J. Engström-Öst. 2022. Copepod reproductive effort and
1078 oxidative status as responses to warming in the marine environment. *Ecol. Evol.* 12: e8594. doi:10.1002/ece3.8594
1079
1080 Wang, H., Pilcher, D.J., Kearney, K.A., Cross, J.N., Shugart, O.M., Eisaman, M.D., and Carter, B.R.
1081 (2023). Simulated impact of ocean alkalinity enhancement on atmospheric CO₂ removal in the Bering Sea. *Earth's*
1082 *Future*, 11, e2022EF002816.
1083
1084 Wernberg, T., Smale, D.A. and Thomsen, M.S. (2012) A decade of climate change experiments on marine
1085 organisms: procedures, patterns and problems. *Global Change Biology* 18, 1491–1498.
1086
1087 White, A.E., Granger, J., Selden, C., Gradoville, M.R., Potts, L., Bourbonnais, A., ... and Chang, B.X.
1088 (2020). A critical review of the ¹⁵N₂ tracer method to measure diazotrophic production in pelagic ecosystems.
1089 *Limnology and Oceanography: Methods*, 18(4), 129-147.
1090
1091 Worsfold, P.J., Clough, R., Lohan, M.C., Monbet, P., Ellis, P.S., Quérel, C.R., Floor, G.H. & McKelvie,
1092 I.D. (2013) Flow injection analysis as a tool for enhancing oceanographic nutrient measurements—A review.
1093 *Analytical Chimica Acta* 803, 15-40.
1094
1095 Yang, B., Leonard, J., Langdon, C. (2023) Seawater alkalinity enhancement with magnesium hydroxide
1096 and its implication for carbon dioxide removal. *Marine Chemistry* 253, 104251.