# **A description of Ocean Forecasting Applications around the Globe**

Jennifer Veitch<sup>1,19</sup>, Enrique Alvarez-Fanjul<sup>2</sup>, Arthur Capet<sup>3,4</sup>, Stefania Ciliberti<sup>5</sup>, Mauro Cirano<sup>6</sup>, 2 Emanuela Clementi<sup>7</sup>, Fraser Davidson<sup>16</sup>, Ghada el Serafy<sup>8</sup>, Guilherme Franz<sup>9</sup>, Patrick Hogan<sup>10</sup>, Sudheer 3 Joseph<sup>11</sup>, Svitlana Liubartseva<sup>7</sup>, Yasumasa Miyazawa<sup>12</sup>, Heather Regan<sup>17</sup>, Katerina Spanadouki<sup>14,15</sup> 4 5 6 <sup>1</sup>Egagasini Node, South African Environmental Observation Network (SAEON), Cape Town, South Africa 7 <sup>2</sup>Mercator Ocean International, Toulouse, France 8 <sup>3</sup>ECOMOD, Royal Belgian Institutes of Natural Sciences, Brussels, Belgium 9 <sup>4</sup> BGEOSYS, Free University of Brussels, Brussels, Belgium 10 <sup>5</sup>Nologin Oceanic Weather Systems, Santiago de Compostela, Spain 11 <sup>6</sup>Department of Meteorology, Institute of Geosciences, Federal University of Rio de Janeiro (UFRJ), Brazil 12 <sup>7</sup>CMCC Foundation - Euro-Mediterranean Center on Climate Change, Italy 13 <sup>8</sup>Data Science and Water Quality, Deltares, Delft, the Netherlands 14 <sup>9</sup>Federal University of Bahia (UFBA), Brazil <sup>10</sup>NOAA, National Centers for Environment Information, Stennis Space Center, Hancock County, Mississippi, United States 15 <sup>11</sup>Indian National Centre for Ocean Information Services (INCOIS), Pragathi Nagar, Nizampet, Hyderabad, Telangana 16 17 500090. India <sup>12</sup>Application Laboratory, Japan Agency for Marine-Earth Science and Technology, Kanagawa, Japan 18 <sup>13</sup>Instituto Geofisico del Perú, Lima – Perú 19 20 <sup>14</sup>Coastal and Marine Research Laboratory of the Institute of Applied & Computational Mathematics of the Foundation for 21 the Research and Technology (FORTH-IACM), Heraklion, Crete, Greece 22 <sup>15</sup>Institute of GeoEnergy (IG) of the Foundation for Research and Technology (FORTH), Chania, Crete – Greece 23 <sup>16</sup> Environment and Climate Change Canada <sup>17</sup>Nansen Environmental and Remote Sensing Center, Bergen, Norway 24 <sup>18</sup>The Nansen-Tutu Centre, Department of Oceanography, University of Cape Town 25 26 27 28 *Correspondence to*: Jennifer Veitch (ja.veitch@saeon.nrf.ac.za) 29 30 Abstract. Operational oceanography can be considered the backbone of Blue Economy: it offers solutions that can support multiple UN Sustainable Development Goals by promoting sustainable use of ocean resources for economic growth, 31 32 livelihoods and job creation. Given this strategic challenge, the community worldwide has started to develop science-based 33 and user-oriented downstream services and applications that use ocean products as provided by forecasting systems as main 34 input. This paper provides examples of stakeholder support tools offered by such applications and includes sea state 35 awareness, oil spill forecasting, port services and fishing and aquaculture, among others. Also emphasized is the important

- 36 role of ocean literacy and citizen science to increase awareness and education in these critical topics. Snapshots of various
- 37 applications in key world ocean regions, within the framework of the OceanPrediction DCC, are illustrated, with emphasis
- 38 given on their level of maturity. Fully operational examples can be used as inspiration for export to other areas.

#### 39 **1 Introduction**

40 The World Bank defines the Blue Economy as the sustainable use of ocean resources for economic growth, improved 41 livelihoods and jobs while preserving the health of the ecosystem. The Blue Economy has the potential to help address many 42 of the UN sustainable development goals including: no poverty, zero hunger, affordable and clean energy, decent work and economic growth, climate action and life below water. Various programs and associated actions of the UN Decade of Ocean 43 44 Science for Sustainable Development<sup>1</sup> are designed to provide the science to support the Blue Economy as well as to ensure the resilience of both marine ecosystems as well as coastal populations. A key objective of several of the programs is the 45 46 development of improved coast-to-ocean forecasts and predictions and, most essentially, their uptake and usefulness to 47 coastal stakeholders. To achieve this and to support the development of a sustainable Blue Economy, the operational oceanography community should be able to support the development of downstream applications in which model data is 48 49 transformed into tailored information for the end users. These applications are intended to create applied solutions to various societal, environmental and scientific challenges from which both public entities and private companies can benefit and 50 actively take part in the implementation of the so-called "value chain". The ETOOFS (Expert Team on Operational Ocean 51 52 Forecasting Systems) guide on Implementing Operational Ocean Monitoring and Forecasting Systems (Alvarez Fanjul et al., 2022) provides a thorough overview of the need for downstream services as well as examples of advanced systems that 53 54 includes: portals for the dissemination of sea state awareness (e.g. https://data.marine.copernicus.eu/); oil spill forecasting (e.g. MOTHY<sup>2</sup>, WITOIL<sup>3</sup>; MEDSLIK-II<sup>4</sup>), port services (e.g. SAMOA<sup>5</sup> and Aquasafe<sup>6</sup>) ; voyage planning (e.g. VISIR<sup>7</sup>) and 55 56 fishing and aquaculture.

57

In this chapter, we provide only some examples of existing downstream services for eight of the nine regions identified by the OceanPrediction DCC: the West Pacific and Marginal Seas of South and East Asia, Indian Seas, African Seas, Mediterranean and Black Seas, North-East Atlantic, South and Central America, North America and the Arctic. The Antarctic region is not included in this review of downstream services due to the lack of services provided there. The distribution of the regions is based on both the UNEP (United Nations Environmental Programme) as well as the GOOS Regional Alliances, with some clustering.

64

65 The regional sections have been prepared by each of the regional teams of the OceanPrediction DCC
66 (<u>https://www.unoceanprediction.org/en/about/community</u>) and, though not comprehensive, each provide a flavour of the

6 <sup>2</sup> <u>http://www.meteorologie.eu.org/mothy/</u>

- 10 <sup>6</sup> <u>https://hidromod.com/?s=aquasafe</u>
- 11 <sup>7</sup> <u>https://www.visir-model.net/</u>
- 12 13

<sup>5 &</sup>lt;sup>1</sup> <u>https://oceandecade.org/</u>

<sup>7 &</sup>lt;sup>3</sup> <u>http://www.witoil.com/</u>

<sup>8 &</sup>lt;sup>4</sup> <u>https://www.medslik-ii.org/</u>

<sup>9 &</sup>lt;sup>5</sup> <u>https://www.puertos.es/</u>

67 needs in each region as well as some of the downstream application services developed to meet them and their maturity 68 levels. The downstream applications have been broadly grouped as follows: Extremes, Hazards and Safety: Natural Resources and Energy; Shipping, Ports and Navigation and Climate Adaptation and specific contributions for each grouping 69 70 may differ per region. Extremes, Hazards and Safety refers to all extreme events, both offshore (such as marine heat waves) and coastal (such as storm surges), marine pollution (that includes water quality and oil spills) and search and rescue 71 72 operations. Natural Resources and Energy refers to all downstream applications associated with the sustainable exploitation 73 of marine resources (we include aguaculture), renewable energy, tourism and recreation as well as conservation efforts. 74 Shipping, Ports and Navigation includes operational support for research activities (including cruise-track optimization as well as deploying equipment) and Climate Adaptation focuses on longer time-scale tools that are provided to support coastal 75 76 and ecosystem resilience. The examples provided are primarily based on public sector forecasting systems and services, 77 with a few exceptions. The OceanPrediction DCC Atlas of Services, https://www.unoceanprediction.org/, will contain a 78 more complete list of downstream services in each of the regions.

#### 79 2 The West Pacific and Marginal Seas of South and East Asia

In the West Pacific and its marginal sea region, development of operational ocean forecast systems were initiated by governmental operational/research agencies related to meteorology, hydrography, and oceanography in several countries including Australia, China, Japan, Korea, Indonesia and New Zealand. Several downstream services led by the governmental operational agencies have been developed with focusing on support to search and rescue operations and preparation for marine disasters. Recently some industrial applications for fishery and shipping operations have been developed based on close collaborations between scientists and targeted users.

86

87 As a one-stop-shop for the provision of downstream applications with support from the Ocean Decade Collaborative Centre 88 on Ocean- Climate Nexus and Coordination (DCC-OCC) and the Ocean to Climate Seamless Forecasting (OSF) 89 Programme,, China is developing a COAST Toolkit as a knowledge hub and information platform for decision-makers and scientists to obtain information services for action. The Toolkit aims to address the challenge of marine and coastal disasters 90 91 prevention and resources development based on ocean solutions. There are six main modules included in the COAST Toolkit: Module 1: Marine disasters prevention and mitigation: Module 2: Maritime navigation safety, including in the 92 93 Arctic; Module 3: Coastal ecosystem health; Module 4: Integrated coastal zone management; Module 5: Blue economy support; Module 6: Ocean literacy. COAST will deliver predictive capacities, services, and products for marine and coastal 94 95 systems. The products will link field data with complex models and applications with visualization.

96

Examples of various dowstream applications in the West Pacific and marginal seas of South and East Asia is provided in thesections below.

3

#### 99 2.1 Extreme, Hazards and Safety

New Zealand's Moana project (<u>https://www.moanaproject.org/</u>) has developed an interactive particle tracking tool (<u>https://www.moanaproject.org/particle-tracker</u>) on their web portal that allows users to release particles, plankton or larvae into either hindcast or forecast models, based on global or their regionally optimized simulations. This tool supports not only offshore safety operations and oil spill response but also fisheries.

104

105 The Ocean and Climate Early Warning Universal System (OCEANUS), developed by the First Institute of Oceanography 106 (FIO) in China, with the support of the Ocean to Climate Seamless Forecasting System (OSF) Ocean Decade Program, is a 107 similar example of a platform that supports various early warning downstream applications. The OCEANUS platform 108 automatically integrates multi-source observational data, an operational forecast system developed by FIO (the Global Ocean 109 Environment Forecast System: for more information refer to Qiao et al., 2018), automatic post-processing of forecast results, and real-time transmission and release of forecast products. The forecast system supports three downstream applications on 110 111 the OCEANUS platform: Global Coral Reef Bleaching Early Warning System, Global Maritime Search and Rescue Forecast 112 System, Global Oil Spill Response System. Detailed information can be found in the OCEANUS Brochure https://osf-un-113 ocean-decade.com/pdfPreview?id=6401.

114

The FIO-Malaysian Meteorological Department (MMD, also known as Met Malaysia) Ocean Forecasting System, developed in collaboration with the FIO provides 5 day forecasts of surface wave heights, wave period, sea level, ocean currents, sea temperature and salinity for the Malaysian and adjacent seas. These forecasts are operationally disseminated through a web portal hosted by the MMD (Figure-1) and provide early warning to ensure the safety and well-being of marine socioeconomic activities in Malaysia through, for example, oil spill and search and rescue responses.

120

Below some examples specific to particular applications within the West Pacific and Marginal Seas of South and East Asiaare highlighted.

123

#### 124 Search and Rescue

Korea Ocean Observing and Forecasting System (KOOFS) led by Korea Hydrographic and Oceanography Agency (KHOA) provides forecast information required for S&R operations (Republic of Korea/OceanPredict, 2022). The Japan Coast Guard is operating a support system for S&R using an ocean forecasting product provided from the Japan Meteorological Agency (JMA) (Asahara et al. 2015). While also providing ongoing support for S&R, the Australian Bluelink forecast system assisted in the high profile case of the disappearance of Malaysia Airlines flight MH370 (Schiller et al., 2019).

- 130
- 131 Oil Spills

Oil spill tracking models utilizing ocean forecasting products are also developed in several countries including China, Korea, and Japan. For example, an oil spill tracking model coupled with an ocean circulation-tide-wave coupling model was applied for evaluating potential contamination caused by an accident of an oil tanker Sanchi in 2018 around the East China Sea (Qiao et al., 2019). Indonesian Agency for Meteorology, Climatology and Geophysics (Badan Meteorologi, Klimatologi, dan Geofisika, BMKG) is operating downscaled model products for forecasting storm surge and coastal inundation hazards around Jakarta and other port cities in Indonesia (Ramdhani, 2019). Coupling of high-resolution coastal ocean current, wave, river flood models are required for forecasting in real-time and evaluating potential inundation locations in the target cites.

140 Marine Heatwaves

The Moana project in New Zealand aims to improve understanding of ocean circulation, connectivity and marine heatwaves to provide information that supports New Zealand's seafood industry. It provides an operational marine heatwave indicator (<u>https://www.moanaproject.org/marine-heatwave-forecast</u>), as well as sea surface temperature anomalies, based on their regionally optimized operational forecast model.

145



146 147

Figure-1: A snap-shot of the Malaysian Meteorological Departments web portal on which the FIO-MMD Ocean ForecastingSystem is disseminated.

5

150

#### 151 2.2 Natural Resources and Energy

Decadal time scale reanalysis products of ocean and wave models are used for assessing feasibility of ocean renewable energy development around Japan coastal seas and their adjacent Asian Seas (Webb et al., 2020). Reliable estimation of the renewable energy potential associated with wave, ocean current, and thermal energy requires sufficiently long-time duration periods for adequately considering the possible time-dependent natural variability. They have evaluated minimum time duration periods of 20-year for wave and 10-year for ocean current and thermal energy conversion around Japan. The highresolution wave (NOAA WAVEWATCH III) and ocean and tidal current forecast (JAMSTEC JCOPE) models driven by the atmospheric reanalysis forcing were used for calculation of the energy potential reanalysis.

159

In some cases, ocean forecasting data (JCOPE) has been used for marine environmental assessment for exploration of seafloor resources in the Northwestern Pacific such as cobalt-rich ferromanganese crusts (Nagao et al., 2018). Direct velocity measurement using acoustic Doppler current profilers (ADCPs) in deep oceans presents some technical challenges, and combined use of ocean forecasting data and ADCP measurement could be effective for reliable assessment of ocean current variability around the targeted areas (Nagao et al., 2018).

165

In Japan, industrial/commercial use of ocean forecasting is being developed for supporting trade ship navigation (Sato and Horiuchi, 2022), and fishery activities (e.g., <u>https://oceaneyes.co.jp/en/home-2</u>). An early warning system of the abrupt occurrences of strong currents damaging set-net fisheries is operated under intensive collaboration between universities and local fishery agencies in Japan (Hirose et al., 2017). Close collaboration among universities, research institutes, instruments companies, and fishermen demonstrates significant enhancement of marine observation networks through exchange of ocean forecasting information and in-situ observation among them (Nakada et al., 2014; Takikawa et al., 2019). In Oceanian Seas, Bluelink<sup>8</sup> forecast products are widely utilized for maritime transport providers, fishing industries, and tourism operators.

173

## 174 2.3 Shipping, Ports and Navigation

175 Defence

The Royal Australian Navy ingest forecast data produced by Bluelink into their system for Acoustic Geo-environmental Exemplification (SAGE) to calculate range predictions (Schiller et al., 2019). These calculate, for a specific ship, the distance they could expect to detect a submarine or be detected by a submarine, based on the current ocean conditions, estimated from the forecasts provided.

180

181 Sea Level

<sup>24 &</sup>lt;sup>8</sup> https://www.csiro.au/bluelink/

Sea level is vital for port operations. The Australian Bureau of Meteorology provides aggregated sea level forecasts based on the Bluelink operational systems, superimposed with other factors that influence coastal sea-level. Additionally, these forecasts have proven most beneficial when incorporated into existing decision tools that include the BOM river flood warning interface where ocean boundary conditions are improved by the forecasts (Schiller at al., 2019).

186

#### 187 2.4 Climate Adaptation

CSIRO, BOM and the Australian Government's Department of Climate Change, Energy, the Environment and Water have produced a web portal (<u>https://climatechangeinaustralia.gov.au/en/</u>) that provides climate information, projections, tools and data to inform decision-making related to climate change in Australia. The portal incorporates both observational datasets as well as climate projections.

192

#### 193 3 Indian Seas

194 Operational ocean forecast systems and downstream services in the Indian Ocean have several stakeholders, including 195 government agencies, maritime industries, research institutions, and the public. The operational oceanographic services for 196 the Indian Seas underwent significant progress during the past 25 years. These functional systems have several components, 197 which include observation networks designed to collect and research teams to analyze, and disseminate oceanographic data, 198 assimilate the data to numerical models, and provide forecasts to support decision-making, improve safety, and enhance the 199 understanding of the Indian Ocean environment. The Indian Ocean forecasting system, operational at the Indian National 200 Centre for Ocean Information Services (INCOIS) helps several regional small island countries in the Indian Ocean under 201 regional alliances such as Regional Integrated Multi-Hazard Early Warning System for Africa and Asia (RIMES) and the 202 Columbo Security Conclave (CSC). INCOIS serves as the Regional Specialized Meteorological Centre (RSMC) for Global 203 Numerical Ocean and Wave Prediction for the Indian Region as per the WMO mandate. RSMC services are provided to the region through a web portal which can be accessed at (https://incois.gov.in/oceanservices/rsmc\_ocean.jsp), with an example 204 205 of their ocean and wave prediction service provided in Figure-2. Provided below are some key components and applications 206 of these systems.

- 207
- 208



- 210
- 211 Figure-2: Web interface of RSMC for numerical ocean prediction (left) and the same for wave prediction (right)
- 212

#### 213 **3.1 Extremes, Hazards and Safety**

#### 214 Search and Rescue Aid Tool (SARAT)

The Search And Rescue Aid Tool (SARAT<sup>9</sup>) is developed for facilitating individuals/vessels in distress in the shortest possible time. This has been initiated and developed under the Make in India program. The tool uses model ensembles that account for uncertainties in the initial location and last known time of the missing object to locate the person or object with high probability - the movement of the lost objects is governed mainly by the currents and winds.

## 219 Oil Spill Trajectory Prediction

The oil spill prediction system (OOSA<sup>10</sup>) operational at INCOIS works based on the GNOME model which uses ocean currents from an ocean general circulation model and winds from an atmospheric general circulation model to simulate the Lagrangian drift of oil spills which needs initial location of spill and quantity of the oil and type of oil if available for producing movement of oil under the influence of winds and currents.

224 Marine Heat Wave Advisory Services (MHAS)

Marine Heat Waves refers to the anomalous (above 90 percentile ) increase of sea surface temperature compared to the historical (past 30 years) values persistent over consecutive 5 days. These heat waves have profound impact on marine ecology and fisheries and marine biodiversity. In view of the environmental significance of marine heatwaves, India started generating marine heat wave advisories and made it available as a service through the web portal<sup>11</sup>. It also issues special bulletins during excessive and persistent heat waves.

<sup>31 &</sup>lt;sup>9</sup> <u>https://sarat.incois.gov.in/sarat/home.jsp</u>

<sup>32 &</sup>lt;sup>10</sup>https://incois.gov.in/portal/osf/oosa.jsp

<sup>33 &</sup>lt;sup>11</sup> <u>https://incois.gov.in/portal/mhw/index.jsp</u>

## 232 15 3.2 Natural Resources and Energy

#### 233 Potential Fishing Zone (PFZ) Advisories

Using satellite derived SST and Chlorophyll and tapping the habitat preference of fishes, advisories to fishers are provided through a wide range of communication channels such as web-portal<sup>12</sup>, Short Message Services (SMS), Radio, mobile applications and electronic display boards for the past couple of decades and there is positive feedback from fishers about this service. As the fishermen community are of diverse ethnic background and speak multiple languages the services are provided as multilingual texts. There are about 700,000 registered users for this service at present.

239

## 240 Coral Bleach Alert System (CBA)

Coral reefs play a pivotal role in marine ecosystems and are vital for the habitats of flora and fauna in Ocean. Ecologically, coral reefs are significant as they provide a conducive environment for several marine species and thereby contribute to the biological productivity in the Ocean. However, coral reefs are sensitive to Sea Surface Temperature (SST) and sustained thermal stress can cause severe damage to the coral reefs and they get bleached proportionate to the intensity and duration of the thermal stress. India has developed a satellite based operational system for assessing the thermal stress on corals from satellite SST corroborated with ground truthing through field examination of coral damage. This service is for assessing the degree of damage caused to the coral environments within the Indian seas and is made available through a web portal<sup>13</sup>.

#### 248 **3.3 Shipping, Ports and Navigation**

#### 249 Small Vessel Advisory Services (SVAS)

Small Vessel Advisory and Forecast Services System (SVAS<sup>14</sup>) is an innovative impact-based advisory and forecast service system for small vessels operating in the Indian coastal waters. SVAS warns users against potential zones where vessel overturning can take place, ten days in advance. This warning system is based on 'Boat Safety Index' (BSI) derived from wave model forecast outputs such as significant wave height, wave steepness, directional spread and the rapid development of wind sea.

#### 255 **3.4 Climate Adaptation**

#### 256 Climate Indices

Climate indices such as El Nino/ La Nina conditions and Indian Ocean Dipole conditions are computed based on model simulations and made available through the webportal<sup>15</sup>. The status of the above-mentioned inter-annual climate modes are regularly updated and provided to the end users along the indices for the past 12 months. These indices are widely used by policy makers and the agricultural sector as they have significant impact on Indian Monsoon and annual rainfall patterns in the region.

262

<sup>37 &</sup>lt;sup>12</sup> <u>https://incois.gov.in/MarineFisheries/PfzAdvisory</u>

<sup>38 &</sup>lt;sup>13</sup> <u>https://incois.gov.in/portal/coralwarning</u>

<sup>39 &</sup>lt;sup>14</sup> <u>https://incois.gov.in/portal/osf/SVA.jsp</u>

<sup>40 &</sup>lt;sup>15</sup> <u>https://incois.gov.in/portal/ElNino</u>





#### 274 4 African Seas

275 While the development of operational ocean forecast systems and downstream services, optimized for African regional seas 276 and coastal regions is limited, it is ongoing (Uba et al., 2020; de Vos et al., 2021; Hart-Davies and Backeberg, 2023) and 277 various strategies exist to support stakeholders. In the simplest example, local met offices use global services to package alerts for subscribed users via text messages or emails, while others add value to global services by customizing solutions for 278 279 stakeholders. The most advanced services are in the North of the continent, where downstream applications benefit from the 280 advanced Mediterranean Sea operational systems (Cirano et al., 2025), in the Red Sea area where an optimized regional 281 system has been developed (Cirano et al., 2025, Hoteit et al., 2021) and in the far South where a co-designed decision 282 support portal is well established for stakeholders. Examples of approaches to various downstream applications will be 283 provided below.

284 A more cohesive, regional approach to the provision of operational information to support marine and coastal operations in 285 Africa has been established by GMES and Africa (GMES: Global Monitoring for Environment and Security<sup>16</sup>) via 286 MarCOSIO (Marine and Coastal Operations for Southern Africa and the Indian Ocean<sup>17</sup>) and MarCNoWA (Marine and Coastal Areas Management in North and West Africa<sup>18</sup>). These platforms currently make use of global services for earth 287 288 observations as well as marine forecast products that in some cases are optimized for local conditions. Linked to MarCOSIO 289 is the National Oceans and Coastal Information Management System (OCIMS<sup>19</sup>), developed by the South African 290 Department of Forestry, Fisheries and the Environment (DFFE) in collaboration with the Council for Scientific and Industrial Research (CSIR). OCIMS provides customized decision support tools that include coastal flood hazard, operations 291 292 at sea, fisheries and aquaculture, integrated vessel tracking, marine spatial planning, water quality, marine predators. These 293 tools are co-designed with the key stakeholder groups in annual stakeholder engagement workshops that bring together the 294 developers as well as the end-users that include the aquaculture industry. National Sea Rescue Institute (NSRI), marine

<sup>44 &</sup>lt;sup>16</sup> https://au.int/GMESAfrica

<sup>45 &</sup>lt;sup>17</sup> <u>https://marcosio.org/</u>

<sup>46 &</sup>lt;sup>18</sup> <u>https://gmes.rmc.africa/</u>

<sup>47 &</sup>lt;sup>19</sup> <u>https://ocims.environment.gov.za/</u>

<sup>48</sup> 49

authorities and navy, municipalities etc. These tools currently make use of operational satellite products, optimized for the
South African coastline, as well as global forecast models that are not locally optimized. Limited area operational forecast
models are in development (https://somisana.ac.za) and will be integrated into the OCIMS DeSTs within the next year.

## 298 4.1 Extremes, Hazards and Safety

299 Oil Spill

In the case of an oil spill in African waters, global services are generally called upon to assist with the mitigation effort. For example, in the case of the devastating oil spill in the Indian Ocean on 25 July 2020 when the Wakashio Bulk carrier ran aground off Mauritius (Seveso et al., 2021), Mercator Ocean International provided Meteo-France with ocean current forecasts to feed the MOTHY pollutant drift model and the CMCC (Euro-Mediterranean Centre on Climate Change) used Copernicus Marine Service near real time products like forecasted currents and ECMWF winds to forecast the weathering and transport of the oil slick.

The SOMISANA team in South Africa have developed a pre-emptive approach in which they release a 'virtual' oil spill at each of the ship-to-ship refueling locations within their high-resolution bay-scale models. They use a simple lagrangian particle tracking approach to allow the hypothetical oil spill to be tracked 5 days into the future. Additionally, their oil spill tracking functionality, developed using the OpenDrift software, allows for seamless tracking between the global and coastal/bay-scale forecast models and can be launched on demand.

The iRED-M1 system (Hoteit et al., 2021) developed at the King Abdullah University of Science and Technology provides an ocean-wave-atmosphere coupled forecast system with dedicated web servers for interactive visualization, analytics and queries. These forecasts are used mainly for oil-spill trajectories as well as to provide assessments on extreme weather and wave conditions.

315 Storm surge

316 Storm surge information was highlighted as being important all of the time in Eastern African countries due to the frequent 317 flooding events events that occur in association with cut-off low events and tropical cyclones and that have serious 318 ecosystem, socio-economic and health impacts (Mather and Stretch, 2012; Ravela et al., 2013; Cambaza et al., 2019, Molua 319 et al, 2020; Singh et al., 2023). In South Africa and Mozambique the met services and a local municipality have developed 320 downscaled storm surge models (Cirano et al., 2025) in order to provide early warnings to coastal stakeholders. These 321 forecasts are provided either on an operational web portal (e.g. https://marine.weathersa.co.za/Forecasts Surge.html) and/or 322 by early warnings that come in the form of emails or text messages to subscribed users that include port authorities, fishing 323 communities, NGOs and consultants.

- 324
- 325 Search and Rescue

The South African OCIMS provides an Operations at Sea decision support tool (<u>https://www.ocims.gov.za/coastops/</u>), that operationally disseminates marine weather information that includes NOAAs GFS wind and wave forecasts, historic winds and waves based on the downscaled atmospheric models that are run by the South African Weather Service (SAWS). As an additional tool that has been custom-built for and requires a login from the National Sea Rescue Institute (NSRI), allows the user to use global wind, wave and current forecasts to optimize search domains.

## 331 4.2 Natural Resources and Energy

## 332 Fisheries management

333 Despite fisheries being consistently identified as the most essential coastal activity requiring operational forecast services 334 throughout the African Seas regions, relatively few downstream applications exist to support the industry. One example is 335 ABALOBI (https://abalobi.org/) that is a South African-based enterprise that aims to support the sustainability of small-scale 336 fishing communities through technology. ABALOBI provides a mobile application that is designed for users that span the 337 value-chain from small-scale fishers to consumers. The application provides forecast information about marine weather 338 (from the NCEP Global Forecast System) and also notification about red tide events (derived from CMEMS satellite 339 information), but also provides various logging and business management tools. ABALOBI supports the traceability of 340 seafood, fully documented fisheries, fair and transparent supply chains and community cohesion and entrepreneurship 341 (ABALOBI Impact Report).

The fundamental triad of enrichment, concentration and retention along with the transport of fish eggs and larvae from their spawning to nursery areas is critical for the sustainability of the high productivity that supports the lucrative South African fishing industry. Furthermore, connectivity between marine protected areas is an essential component in the health and longevity of marine ecosystems. To this end, many studies have made use of numerical ocean models to force lagrangian particle experiments in order to understand these transport and retention processes and their various impacts (Pfaff et al., 2022; Heye, 2021).

#### 348

#### 349 Aquaculture

In order to reduce the impact of harmful algal blooms (HABs) on the South African aquaculture industry such as the extreme event that occurred on the South West Cape Coast in 2017 and that caused the mortality of ~250 tonnes of farmed abalone (Groom et al., 2019), OCIMS has incorporated a HAB decision support tool (https://www.ocims.gov.za/hab/app/). This operational tool provides a matrix of probability of HABs occurring in key locations along the South African coastline. The spatial and and temporal extent of the bloom is captured by remotely sensed chlorophyll data that is provided by the EUMETSAT datastore (Sentinel 3 OLCl & SLSTR) and the Copernicus Marine Service (Global Color chl-a) and chl-a estimates are optimized for high biomass bloom water types (Smith et al., 2018).

## 357 4.3 Shipping, Ports and Navigation

The South African Weather Service provides regionally optimized wind and wave forecasts to support port operations. The CSIR's Vessel Motion Forecast Tool (Troch et al., 2023) utilizes numerical models to predict long-period wave climates and subsequent moored ship motions, providing port authorities with important information regarding vessel stability. This tool enables port operators to assess the suitability of different vessel sizes at berths for both current and forecasted wave conditions, directly improving operational efficiency and safety. By linking numerical models and providing an intuitive user interface, the tool delivers actionable insights into potential berth-specific issues, allowing for proactive planning and minimization of downtime.

## 365 4.4 Climate Adaptation

Digital Earth Africa (DE Africa: <u>https://www.digitalearthafrica.org/platform-resources/services/coastlines</u>) significantly
 supports climate adaptation along African coastlines through its Coastlines application. This tool leverages satellite imagery

368 and data analysis to monitor coastal erosion, inundation, and shoreline changes, critical factors influenced by climate change. 369 By providing time-series data, DE Africa helps identify vulnerable areas and track the impact of rising sea levels and 370 increased storm surges. While the Coastlines application primarily utilizes satellite data, it can be enhanced by incorporating 371 predictive models. For example, hydrodynamic models forecasting wave action and sea level rise can be integrated to project 372 future coastal changes. Additionally, climate models that predict changes in rainfall patterns and storm frequency can inform 373 the interpretation of observed coastal shifts, allowing for more robust risk assessments and adaptation planning. This 374 integration of data and models enables informed decision-making for coastal management, infrastructure planning, and 375 community resilience in the face of a changing climate.

376

#### 377 5 Mediterranean and Black Seas

During the last decades, the constant evolution of increasingly accurate operational forecasting systems in particular in the Mediterranean Sea and, at a lower extent in the Black Sea, from regional to local and coastal scales providing systematic information of the essential ocean variables, has led to the consolidation and to the development of a wide range of scientific and societal applications in the area.

Mediterranean and Black Sea analysis and forecast operational numerical products, such as the ones delivered through the Copernicus Marine Service (https://marine.copernicus.eu) by the Med- (https://marine.copernicus.eu/about/producers/medmfc; Coppini et al., 2023) and BLK- (https://marine.copernicus.eu/about/producers/bs-mfc; Ciliberti et al., 2022) MFCs (Monitoring and Forecasting Centers) are essential to provide a 3 dimensional state of the sea including: currents, temperature, salinity, mixed layer thickness, sea level, wind waves, and biogeochemistry to support many downstream applications and activities.

Considering that the two basins are characterized by a large variety of complex physical processes occurring on a wide range of spatiotemporal scales, it is required to develop models that can reproduce specific ocean variables evolutions and to focus on specific processes representation (from wind driven and thermohaline circulation to water mass formation, coastal processes such as upwelling and storm surge, extreme and fast events such as medicanes). Following all these needs, the Mediterranean and Black Sea communities have been implementing models based on different codes and parameterizations properly designed to solve specific problems.

Several downstream applications developed and implemented in the Mediterranean and Black seas are presented hereafter considering: climate change studies, oil spill, ship routing, search and rescue, marine litter, ports, water quality, fish and larvae dispersion, fisheries and aquaculture management as well as adaptation and management strategies. Most of the listed applications are described in a recent book from Schroeder and Chiggiato (2022) who edited an introductory guide on the oceanography of the Mediterranean Sea and in the ETOOFS (Expert Team on Operational Ocean Forecasting Systems) guide from Alvarez Fanjul et al. (2022).

## 400 **5.1 Extremes, Hazards and Safety**

401 Oil spills

Oil spill models are forced by meteo-ocean forecasting products providing ocean currents, wind and waves which should be
 available on a regular basis. Several oil spill models are operated in the Mediterranean and Black seas and specific
 forecasting systems have also been implemented in areas of oil spill emergencies such as those presented in Cucco et al.

405 (2012). Moreover, oil spill modeling in harbor and port areas have been developed, such as in the Port of Taranto in south 406 Italy (Liubartseva et al., 2021), the Limassol port areas in Cyprus (Zodiatis et al., 2024), the Port of Tarragona in Spain 407 (Morell Villalonga et al., 2020), the Spanish harbors through the SAMOA project launched by Puertos del Estado (PdE). 408 Additionally, MEDSLIK (Zodiatis et al., 2021) and MEDSLIK-II (De Dominicis et al., 2013), Lagrangian oil spill models 409 for short term forecasting, were applied in various areas. Several Decision Support System (DSS) dedicated to oil slicks 410 emergencies and predictions in the Mediterranean Sea have been developed such as: the French MOTHY (Daniel, 1996) 411 drift system, the Italian the WITOIL (Where Is The Oil) multi-model DSS, the MEDESS4MS (Zodiatis et al. 2016: 412 Sorgente et al., 2020). The OILTOX lagrangian oil spill model adapted for the Black Sea environment for oil spill transport 413 and fate has been implemented in the North-western shelf of the Black Sea and Dnipro-Boog Estuary (Broychenko et al., 414 2003). The POSEIDON Oil Spill fate and trajectory model is based on the PARCEL model (Pollani et al., 2001) which is 415 able to simulate not only the drift of the oil but also the chemical transformations under the specific environmental 416 conditions.

417 Search and Rescue

An advanced web-based and mobile decision support system for search-and-rescue (SAR) in the Mediterranean has been developed by Coppini et al. (2016). The system simulates drifting objects at sea, using the met-ocean data provided by the Copernicus Marine Service as an input. The performance of the service is evaluated by comparing simulations to data from the Italian Coast Guard pertaining to actual incidents in the Mediterranean Sea.

422 At the national and international level, the National Forecasting Centre of Météo-France provides met-ocean support and 423 drift forecasts to assist authorities in charge of search and rescue operations. The aforementioned MOTHY system can 424 resolve not only search and rescue targets but it also computes the drift of lost cargo containers (Coppini et al., 2022). The 425 system uses the Copernicus Marine Service data among several forcing fields.

- The Hellenic Centre for Marine Research (HCMR) has an agreement with the Hellenic Coast Guard for a SAR service in the Greek seas. The application is developed and hosted at the POSEIDON operational system and provides forecasting of drifting objects.
- Currently, under the ever-increasing flow of people trying to reach Europe by crossing the Mediterranean Sea, the efficiency
   of SAR calls for an enhancement. That requires both improved modeling of drifting objects and optimized search assets
   allocation.
- In the Adriatic basin, Slovenian Environment Agency provides met-ocean support and drift forecasts to assist authorities in charge of search and rescue operations (Ličer et al., 2020) and is based on high-resolution wind forecasts and ocean modeling downscaling of Copernicus Marine Service forecasts for the Med Sea. The system can resolve search and rescue targets, oil spills and cargo containers.
- 436 Marine litter

Marine plastic pollution, usually from anthropogenic sources, is increasingly recognized as an emerging threat to the Mediterranean environment, biodiversity, human health, and well-being (Schroeder and Chiggiato, 2022). Recently, an important shift has been conducted for the Mediterranean Sea from the spatially-uniform distributions of plastic sources to a more realistic representation of land-based and offshore inputs (Liubartseva et al., 2018; Macias et al., 2019; Soto-Navarro et al., 2020; Kaandorp et al., 2020; Tsiaras et al., 2021; Tsiaras et al., 2022a) and for the Black Sea (Miladinova et al., 2020; Stanev and Ricker, 2019, Gonzalez-Fernandez et al., 2022) to identify the accumulation and dissipation of floating litter in such semi-enclosed sea basins.

14

## 444 Water quality

445 The physical-biogeochemical forecasting system for the Northern Adriatic Sea developed in the framework of the

CADEAU project (Bruschi et al., 2021) is based on a high resolution (up to around 750m) implementation of the MITgcmBFM coupled model (Cossarini et al., 2017) targeting water quality and eutrophication, and it uses the daily Med-MFC
products for initialization and to constrain the open boundary.

The trophic index (TRIX) eutrophication assessment indicator has been calculated both on in situ data and with a coupled circulation and biogeochemical numerical modeling system. TRIX is defined by four state variables: chlorophyll-a, oxygen, dissolved inorganic nitrogen, and total phosphorus. As an example, the trophic index differences have been computed to evaluate the trophic state of marine waters along the Emilia-Romagna coastlines (Italy) and over the whole Adriatic Sea (Fiori et. al, 2016).

A relocatable modelling system for describing and forecasting the microbial contamination that affects the quality of bathing waters was implemented at five coastal areas in the Adriatic Sea, which differ for urban, oceanographic and morphological conditions (Ferrarin et al., 2021). The modelling systems are all based on the hydrodynamic finite element model SHYFEM (Umgiesser et al., 2022). Pollution events are mainly triggered by urban sewer outflows during massive rainy events, with relevant negative consequences on the marine environment and tourism and related activities of coastal towns.

## 459 **5.2 Natural Resources and Energy**

## 460 Fish larvae dispersion, fishery and marine aquaculture management

The study of larvae dispersion, regional connectivity and their impact on the structure of species populations and fisheries are generally provided using lagrangian models (van Sebille et al., 2018; Laurent et al., 2020; Melaku Canu et al., 2020) and in the Mediterranean sea these have been carried out thanks to the availability of information provided by operational forecasting systems (more information on such applications can be found in Schroeder and Chiggiato, 2022).

Being strongly supported by the policies and initiatives of the European Union, marine aquaculture guarantees food security and reduces the fishing pressure on wild fish stocks. Farm site selection strategy based on an aquaculture suitability index has been developed for the Central Mediterranean (Porporato et al., 2020). The index is based on the outputs of ecophysiological models which were forced using time series of sea surface temperature, significant wave height, distance to harbor, current sea uses, and cumulative impacts. Tyrrhenian and Ionian coastal areas are found to be more suitable, compared to the Northern Adriatic and southern Sicilian ones.

- 471 Small pelagic fish play a key role in marine food webs, being the trophic link between plankton and larger fish. Given their 472 pronounced sensitivity to environmental changes, end-to-end (physics-plankton-fish) small pelagic fish two-way coupled 473 models (Gkanasos et al., 2021) are unique tools that can be used to study the impact of climate change and fisheries in a 474 single modeling framework.
- 475 Coupled hydrodynamic/biogeochemical models can also be used to evaluate the environmental impact of aquaculture waste
  476 and investigate the carrying capacity of coastal marine ecosystems (Tsiaras et al., 2022b; Tsagaraki et al., 2011).

Moreover, Dynamic Energy Budget (DEB) models (Hatzonikolakis et al., 2017), forced with hydrodynamic/biochemical
model output (temperature, Chl-a), can be also implemented to simulate the growth of farmed mussels (Mytilus
galloprovincialis) and the potential impact of future climate on their habitat suitability.

480 Adaptation and management strategies to address harmful algal blooms and jellyfish outbreaks

In recent years, eutrophication phenomena, prompted by global warming and population increase, have stimulated the proliferation of potentially harmful algal taxa resulting in the prevalence of frequent and intense harmful algal blooms (HABs) in coastal areas of the Mediterranean and Black seas. Drivers of HABs in coastal areas of Eastern Mediterranean were studied by means of a machine learning methodological approach (Tamvakis et al., 2021). Water temperature has been found to have the most powerful effect on genera's presences.

A jellyfish outbreak forecasting system has been developed for the Mediterranean Sea as a preventive and mitigation tool for citizens and coastal stakeholders, aiming to reduce the jellyfish blooms socio-economic impact in coastal areas through a feasible and powerful management strategy (Marambio et al., 2021). The system explores the Copernicus Marine Service output to predict the jellyfish spatio-temporal distributions.

Previously, the high-resolution ocean modeling was applied to examine the transport and stranding of the pelagic stinging jellyfish Pelagia noctiluca on the Ligurian Sea coast (Berline et al., 2013). Jellyfishes were modeled as Lagrangian particles transported by sea currents with a diel vertical migration. Two environmental factors were found to be critical: the position of the Northern Current and the wind regime.

## 494 **5.3 Shipping, Ports and Navigation**

## 495 Ship routing

The GUTTA-VISIR system is a tactical, global-optimization, single-objective, deterministic model system for ship route planning (Mannarini et al., 2016; Mannarini and Carelli, 2019), which has been implemented in the Mediterranean Sea for several applications (i.e. in the Adriatic Sea, Mannarini et al., 2021) using the analysis and forecast wave and current fields from the Med-MFC.

#### 500 Ports

To respond to the need for information on wind, waves and sea level at the scale of ports and harbor, a Spanish initiative has been developed and operationally implemented called SAMOA-2 (Álvarez Fanjul et al., 2018; Sotillo et al., 2019; Garcia-Leon et al. 2022) operating in 31 ports. It is an integrated system based on Copernicus Marine data, the service provides daily forecasts of sea-level, circulation, temperature and salinity fields at horizontal resolution that range from 350 m (coastal domains) to 70 m (port domains). Another example implemented along the Spanish coastal waters is provided by PORTUS (https://portus.puertos.es/), an early warning system that employs both the in-situ data and the operational forecasts (Álvarez Fanjul et al., 2018).

## 508 5.4 Climate Adaptation

509 Over the last decades, marine heat waves (MHWs) are expected to become more intense, longer and more frequent through 510 anthropogenic warming. Combining high-resolution satellite data and a regional reanalysis, Dayan et al. (2023) have studied 511 MHWs to understand how much each Mediterranean country's Exclusive Economic Zone waters may be affected.

As was stated in the 2nd Edition of the Copernicus Marine Service Ocean State Report, ocean deoxygenation is found to be one of the most pernicious, yet under-reported side-effects of human-induced climate change. This problem is particularly acute in the Black Sea, where Capet et al. (2016) have found the decline of the Black Sea oxygen inventory. The reason for this is that atmospheric warming reduces the ventilation of the lower oxic layer by lowering cold intermediate layer formation rates.

## 517 6 North-East Atlantic

518 The structured provision of mature regional core services and coastal operational forecasting systems in the North-East

519 Atlantic (Cirano et al., 2025) enabled a significant deployment of downstream operational services addressing a wide variety 520 of sectors (Figure-4).

A rich portfolio documenting use-cases of downstream services uptake can be found for instance at the Copernicus Marine Service User Uptake portal and the ETOOFS Guide (Alvarez Fanjul et al., 2022). In particular, the EuroGOOS coastal working group roadmap for operational coastal services (El Serafy et al., 2023) details components of the coastal services value chain in Europe and reviews the status, gaps, and steps needed to improve these services and the sustainability of their provision. A full review of the downstream services that are presently active or upcoming in the established sectors of the European Blue Economy is given in El Serafy et al. (2023). Here we highlight a few examples for selected sectors.



527

Figure-4: Principal characteristics of CMS regional core services for the North-East Atlantic region and its relation to its downstream use in sectors.

## 530 6.1 Extremes, Hazards and Safety

## 531 Oil spills

532 Coastal areas with industrial ports and harbors are among the locations most at risk from oil spill pollution, which heavily 533 impacts aquatic life and ecology, the coastal infrastructures, and the local economy. This underlines the need for timely and 534 accurate coastal services for operations and disaster response. Oil spill models predicting the fate and the transport of the oil 535 slick have been recently enhanced by downscaling from state-of-art regional models (e.g. Copernicus Marine Service) and to very high-resolution hydrodynamic models for coastal and harbor areas. A coastal service in water monitoring and oil spill 536 537 pollution is the OKEANOS project (https://parsec-accelerator.eu/portfolio-items/okeanos/), a web-based integrated and 538 intuitive service combining open-source satellite observations (i.e., affordable), artificial intelligence and high-resolution 539 ocean modelling (i.e., accurate). Another example of oil spill forecasting is the drift model MOTHY developed by Meteo 540 France, which uses ocean currents from the Copernicus Marine Global Ocean Forecast model. This system allows

541 predictions of the possible trajectory of oil spills and estimates the resulting impacts several hours or days in advance. 542 MOTHY has been operational since 1994 and is frequently activated for actual spills or search and rescue operations.

## 543 6.2 Natural Resources and Energy

## 544 Aquaculture sector

Novel coastal services, including mapping of suitable fishing areas, fronts detection, marine conditions and scheduler, land pollution, site prospection, spat capture assistance, and contaminant source retrieval, are provided by FORCOAST (https://forcoast.eu/) in aquaculture pilot sites among others regional waters the North Sea, Baltic Sea, and the coastal Atlantic Ocean. These services are Copernicus-based services that incorporate Copernicus products, local monitoring data, and advanced modeling.

Recent projects that aimed at the co-development with end users and demonstration of Harmful Algal Blooms (HAB) forecasting services as one of the societal needs from the coastal observing and forecasting systems include the FP7 Asimuth (Cusack et al., 2016), H2020 AtlantOS (Cusack et al. 2018) and Interreg Atlantic Area PRIMROSE (<u>https://pml.ac.uk/projects/primrose-predicting-risk-and-impact-of-harmful-eve/</u>), all providing near real-time and forecast information for the aquaculture industry along Europe's Atlantic coast.

Last, but not least, all the data and information produced by operational coastal services may be used in the framework of the Maritime Spatial Planning Directive to identify Allocated Zones for Aquaculture (AZA), following national and international guidelines (e.g. FAO, Macias et al. 2019), as shown by use cases as AQUAGIS (European Aquaculture Society - ePoster Viewer).

559 Coastal tourism sector

Various coastal services have been developed following inquiries from the coastal tourism sector. A good example is a tailored product based on the North East Atlantic operational forecasting model in Ireland developed by Irish Marine Institute (IMI). Surface currents subsets are provided over five geographical areas around the Irish waters and the English Channel and published in a GRIB format via an ftp site (https://sftp.marine.ie/), while ensuring low data volume. The service was developed in collaboration with the sailing community that contacted the IMI to request its development and was notably used during the Fastnet sailing race.

Another Irish example serves beach goers. The Irish Environmental Protection Agency in collaboration with Local Authorities and the Department of Housing, Planning and Local Government run a webpage https://www.beaches.ie, where the latest information on water quality and others is presented for 204 beaches in Ireland. Met Eireann (the Irish national met service) and the Marine Institute contribute to the information provided with current weather and weather forecasts and tidal information, respectively.

571 Among the services that provide the latest water quality information, the service carried out in the framework of the 572 CADEAU project (Bruschi et al., 2021) provides data and information to assess the potential impact of bacterial pollution 573 sources on bathing waters (as defined in the EU Bathing Water Directive) and help bathing waters' managers in identifying 574 potential sources of impact and planning mitigation measures.

National marine forecasting agencies also serve the coastal tourism sector. The Marine Forecasting Centre of Belgium of the
Royal Belgian Institute of Natural Sciences (RBINS) issues 5-day forecasts of the marine conditions in the North Sea twice a
day with a high resolution for the Belgian part of the North Sea. These forecasts are used in numerous applications among

them the tourism and leisure industries. Surfers use the application for mobile devices to schedule their sessions for good

579 waves and current conditions.

580 Renewable energy sector

The renewable energy sector is a prominent player in the Blue Economy and therefore one of the main potential users of coastal services. Indeed, EU hosted 70% of global ocean energy (wave and tidal) installed capacity, and 86% of the world's total installed offshore wind capacity at the end of 2018 (Díaz and Soares, 2020), while jobs in the offshore wind energy sector have multiplied nine-fold in less than 10 years (European Commission, 2020).

585 Current bottlenecks relating to the large-scale installation of ocean multi-use activities are addressed by the UNITED project 586 (https://www.h2020united.eu/), which demonstrates business synergies and benefits of ocean multi-use; provides a roadmap 587 for deployment in future multi-use sites and potential scaling barriers to be addressed through best practices and lessons 588 learnt. Another example of coastal services for the renewable energy sector is Ireland's Marine Renewable Energy Portal 589 (http://www.oceanenergyireland.ie/), an online access point for all relevant information and data related to Irish marine 590 renewable energy activity and resources including maps, tools, and information for renewable energy site assessment, 591 development, and management.

## 592 **6.3 Shipping, Ports and Navigation**

593 Coastal information services tailored to the needs of the port sector are provided by the HiSea project 594 (https://hiseaproject.com/). The services include early warning service on potential risk factors issuing alerts on storms, 595 harmful algal blooms, faecal contamination, and other hazards regarding pollution accidents to identify the appropriate 596 responses. It provides key performance indicators regarding fish growth rates, environmental conditions, or the level of 597 vulnerability to storms for vessels, and information for planning operations including accurate and reliable meteorological, hydrodynamic, and water quality forecasts. Further examples of platforms and services for ports are SAMOA and 598 599 AQUASAFE. The SAMOA service from Puertos del Estado aims to provide high-resolution coastal operational prediction 600 systems in domains such as harbours and nearby coastal waters, for different Spanish Port Authorities (Sotillo et al., 2019). 601 Similarly, the AQUASAFE platform is operational for all Portuguese Ports and in the Port of Santos (Brazil). This platform 602 aims to increase efficiency and safety in port operations, by providing access to real time and forecast information. It is also 603 used to support aquacultures, inland navigation, irrigation, and water utilities.

## 604 6.4 Climate Adaptation

605 Climate adaptation is central to the efforts in the North East Atlantic region, where regional core services and operational 606 forecasting systems play a vital role in responding to the impacts of climate change, such as rising sea levels, extreme 607 weather, and changes in marine ecosystems. Key systems like the Copernicus Marine Environment Monitoring Service 608 (CMEMS), the European Centre for Medium-Range Weather Forecasts (ECMWF), and the UK Met Office's coastal 609 forecasting systems provide essential data on oceanographic and atmospheric conditions, aiding climate resilience in marine 610 sectors like fisheries, shipping, and coastal infrastructure. Initiatives such as the Atlantic Action Plan for a Sustainable Blue 611 Economy, the Interreg North Sea Region Programme, and the European Maritime and Fisheries Fund (EMFF) are focused 612 on enhancing climate resilience, offering solutions like adaptive coastal management, improved early warning systems, and 613 sustainable practices.

#### 614 **7 South and Central America**

The lack of available regional core services and coastal operational forecasting systems in South and Central America (Cirano et al., 2025) makes the development of downstream applications difficult. For instance, very few use case demos are described in the Copernicus Marine Service User Uptake for this region. Normally, downstream applications are only developed in partnership with universities or specialized companies capable of implementing operational systems based on a downscale approach from global models.

620 Despite the general lack of regional systems for coastal operational forecast systems in South and Central America, smaller-621 scale services exist and provide useful information for stakeholders. For example, the Baía Digital project 622 (http://baiadigital.com/en/) in Brazil, is a portal that integrates various data sources, including regional model forecasts 623 focuses on developing an operational digital platform to provide environmental, social, and economic information in the 624 region of Guanabara Bay and its surroundings. The diagnostic and prognostic information generated comes from different 625 sources, such as historical databases, data collection platforms, and numerical computational models. Atmospheric and 626 oceanic regional model forecasts represent temporally and spatially the marine and atmospheric dynamics of the Guanabara 627 Bay region. The digital platform has been developed and improved from the interaction between professionals from different 628 areas of science and students from different educational levels, investing in the technical and scientific training of 629 researchers. In addition, extension activities involving students from the school segment will be planned to aim at promoting 630 a scientific culture based on knowledge of Guanabara Bay. The project base is the Laboratory of Computational Methods in 631 Engineering (LAMCE), located in the UFRJ Technological Park, in partnership with other laboratories and teaching and 632 research institutions. The project represents a pioneering effort associated with the regional initiatives of the Atlantic 633 International Research Center (AIR Centre).

634 In the next sections we showcase a number of bespoke downstream applications based on specific needs.

#### 635 7.1 Extremes, Hazards and Safety

636 Oil spills

637 The Brazilian Oil Research Group (BROIL) was created in response to the oil spill disaster that impacted more than 3,000 638 km along the north-northeastern Brazilian coastline in 2019, with significant environmental, economic, and social impacts. 639 BROIL comprises institutions in Brazil (e.g., UFBA, UFPE, UFRJ, INPE and PUC-Rio) and abroad (e.g., OOM-Portugal; 640 IRD/LEGOS-France, HZG-Germany), BROIL works upon three main pillars: (i) detection, through remote sensing 641 techniques; (ii) control, through a set of hydrodynamic and oil spill models; and (iii) remediation, through a set of biota oil-642 exposure case studies (Franz et al., 2021). Numerical models used to predict oil spill trajectory include the Regional Ocean 643 Modeling System (ROMS) and the Lagrangian model MEDSLIK-II. Recently, a partnership with the Brazilian Sea 644 Observatory will enable the use of forecasts with higher resolution hydrodynamic models and to predict the oil spill 645 trajectory automatically through the MOHID modeling system.

646

The North Coast Project (http://www.projetocostanorte.eco.br/) also integrated research groups with different expertise for the development of a method for determining the vulnerability of mangroves to contamination by oil and for producing knowledge about the Brazilian North Coast, in cooperation among ENAUTA, the Nucleus of Studies in Geochemistry and Marine and Coastal Ecology (NEGEMC) of UERJ, the Laboratory of Computational Methods in Engineering (LAMCE) of COPPE/UFRJ, the Laboratory of Research in Marine Environmental Monitoring (LAPMAR) of UFPA and PROOCEANO,

- a Brazilian company of oceanographic technology. The largest continuous area of mangrove forests in the world is found on
- 78 79

the north coast of Brazil – located between the states of Maranhão and Pará – totaling around 7,400 km2, which corresponds to 4.3% of the entire area of mangrove forests in the world. The main objective of the project was to determine the vulnerability, sensitivity, and susceptibility to oil contamination of the mangroves, based on the development of numerical hydrodynamic models with multiple resolution scales and the use of data assimilation techniques to represent large and mesoscale oceanographic phenomena, with seasonal and interannual variability, to small-scale phenomena with daily variability, such as tidal currents in floodplains. The hydrodynamic modeling results were used as input data for the modeling of the transport and dispersion of oil.

660

661 Civil protection

The water level increase due to storm surges can be of the same order of magnitude as tide amplitude along the south-eastern Brazilian coast (Franz et al., 2016). Following a downscaling approach, water level forecasts are available to this region, aiming to help civil protection actions. Water level forecasts, as well as data from several tide gauges along the Santa Catarina coast, are available for the public in general on the EPAGRI's company website<sup>20</sup>. The water level forecasts of high-resolution models (e.g., Babitonga Bay) are also available for port operation. The operational models developed by the Brazilian Sea Observatory initiative (Franz et al., 2021) were updated in collaboration with EPAGRI, considering GEOGloWS<sup>21</sup> flow predictions for major rivers.

669

#### 670 Coastal Engineering

671 Coastal models developed by the Centre for Marine Studies (CEM - UFPR) within the scope of the Brazilian Sea 672 Observatory initiative, through the application of the MOHID modeling system, were used to support local companies in the 673 design of submarine outfalls and study of environmental impacts of bridge construction.

#### 674 7.2 Natural Resources and Energy

675 Aquaculture

676 Information on water quality in bays and estuaries is essential for planning and managing bivalve mollusc production (e.g., 677 water temperature, microbiological contamination, salinity and nutrients). These parameters are influenced by marine 678 currents, river flows, solar radiation and winds, as well as by urbanization pressure and consequent contamination of water 679 bodies (Cabral et al., 2020). The numerical modeling system MOHID was applied to the main aquaculture production zone of shellfish in Brazil, located in the bay of Ilha de Santa Catarina, with the objective of integrating the range of 680 681 environmental data in a hydrodynamic and water guality model capable of simulating the variables of greatest interest in the 682 production of bivalve molluscs, thus serving as a powerful management tool (Garbossa et al., 2023; Garbossa et al., 2021; 683 Lapa et al., 2021). The model was recently implemented in operational mode by the company EPAGRI to provide forecasts, 684 nested within a regional model developed in partnership with universities (e.g., UFPR), as a continuation of the Brazilian Sea 685 Observatory initiative (Franz et al., 2021).

<sup>81 &</sup>lt;sup>20</sup> <u>https://ciram.epagri.sc.gov.br/index.php/maregrafos/</u>

<sup>82 &</sup>lt;sup>21</sup> <u>https://geoglows.ecmwf.int/</u>

#### 686 **7.3 Shipping, Ports and Navigation**

#### 687 Ports

Within the objective of increasing navigation security, São Paulo (Brazil) Harbor Pilots (Praticagem de São Paulo in Portuguese) has been using the AquaSafe platform (https://aquasafe.hidromod.com/landing-page/about), developed by the Portuguese company HIDROMOD and locally implemented in partnership with the University of Santa Cecília (Unisanta) (Ribeiro et al., 2016). The data provided by the platform assists in choosing the better entering and leaving periods of the harbor. The AquaSafe platform is connected to a real-time sensor data stream (tide gauge, weather station, and ADCPs) from Praticagem's Center for Coordination, Communication, and Traffic Operations (C3OT). Furthermore, are also available high-resolution forecast solutions for wave parameters, sea level, wind, and other meteo-oceanographic parameters.

#### 695 7.4 Climate Adaptation

696 The BASIC Cartagena is an applied research project on Basin Sea Interactions with Communities in the coastal zone of 697 Cartagena (Colombia). Located on the Caribbean coast in the north of Colombia, Cartagena and its surrounding beaches 698 represent the Country's principal touristic destination. The first phase of the project started in July 2014 and was completed 699 in June 2017 under the title "Reducing the risk of water pollution in vulnerable coastal communities of Cartagena, Colombia: 700 Responding to climate change." The second phase of the project, titled "Building Resilience in Cartagena Bay," is currently 701 being implemented since February 2018. Its general objective is to contribute to the improved environmental governance of 702 Cartagena Bay by providing scientifically based advice toward climate-compatible and sustainable development policies. 703 Studies of fluvial hydrology are dedicated to the research of the Magdalena River basin, with a focus on surface waters that 704 flow from the Dique Canal towards Cartagena Bay. Analysis of the watershed's human development and climatic conditions 705 permit modeling of the watershed's runoff processes. Future scenarios of climate change and human development will be used to generate prognostics of freshwater discharge from the Dique Canal into Cartagena Bay. In the coastal zone, studies 706 707 focus on the monitoring of water quality and sediment in Cartagena Bay. Analysis of physicochemical and microbiological 708 parameters, as well as contaminants, will permit an impact assessment of human activities and climate variation on the sea, 709 as well as the generation of vulnerability maps. Hydrodynamic modeling will be used for prognostics of the dispersion of 710 fresh water from the Digue Canal into Cartagena Bay under future watershed scenarios.

#### 711 8 North America

712 North America is a vast continent with lengthy continental coastlines that include densely populated areas with busy harbors 713 and vast remote isolated coastlines. Core ocean forecasting services are anchored by national meteorological centers that 714 increasingly trend towards prediction services of the full earth system. This includes the US National Oceanographic and 715 Atmospheric Agency (NOAA), as well as the Canadian Meteorological and Environmental Prediction Center within the 716 federal department of Environment and Climate Change Canada (ECCC). Benefiting ocean forecasting services in North 717 America are mature collaborations between government departments, universities and industry including the US Integrated 718 Ocean Observing System (IOOS) (https://ioos.us) partnership with 11 regional associations and the CIOOS, the Canadian 719 IOOS (https://cioos.ca) networks with 3 regional associations. In Canada, the CONCEPTS initiative coordinates ocean 720 prediction that regroups several federal government departments together including National Defense, Fisheries and Oceans 721 Canada (DFO), the Canadian Coast Guard, the Canadian Hydrographic Service, and the Meteorological Service of Canada.

722 In North America ocean forecast systems are advanced and relatively abundant. They provide a wide range of downstream 723 applications, some of which are described below.

#### 724 8.1 Extremes, Hazards and Safety

725 In the US, the U.S. Coast Guard (USCG) is the primary federal agency for responding to maritime safety and security 726 (including search and rescue and marine pollution) in navigable waters and deep water ports, although other agencies also 727 play prominent roles, including the Environmental Protection Agency (EPA), NOAA, the Federal Emergency Management 728 Agency (FEMA), and State Agencies. The USCG relies on several ocean forecast systems to monitor and predict 729 oceanographic and meteorological conditions critical for navigation, search and rescue, marine pollution and environmental 730 protection, primarily those run by various NOAA entities (National Weather Service, Ocean Prediction Center, OFS, and 731 NCEP). These systems provide data on currents, wave heights, sea surface temperatures, and other factors that impact 732 maritime operations.

In Canada, the Canadian Coast Guard (CCG) make use of the Canadian Operational Network of Coupled Environmental
 Prediction Systems (CONCEPTS) that is collaboratively produced by Environment and Climate Change Change (ECCC),
 Fisheries and Oceans Canada (DFO) and the Department of National Defence (DND) to support their offshore operations.

- 736
- 737 Storm Surge

738 While the coast guards in the respective countries are responsible for the dangers associated with storm surges, storm surge 739 warnings are issued by ECCC in Canada and by the National Hurricane Centre (NHC) and the National Weather Service in 740 the US. The NHC focus on the broader regional picture and use both weather forecasts as well as the SLOSH (Sea, Lake and 741 Overland Surges from Hurricanes: https://vlab.noaa.gov/web/mdl/slosh) model with real-time data to issue warnings via graphical maps and advisories through NOAA websites, television and radio broadcasts, mobile alerts and social media. In 742 743 Canada, the ECCC's Meteorological Service of Canada (MSC) monitor and forecast conditions, based on both global and 744 their own regionally optimized models, that lead to storm surge and coastal flooding. They have recently implemented a 745 comprehensive coastal flooding prediction and alerting program that provides maps that display an index of the probability 746 of storm surges or coastal flooding occurring.

747

#### 748 Oil Spills

749 The Emergency Response Division (ERD) of the Office of Response and Restoration (OR&R) within NOAA provide 750 Environmental Sensitivity Index (ESI) maps and data, which are used to identify vulnerable resources and habitats in 751 advance of emergencies so that appropriate response actions can be planned. ERD works with local experts to develop or 752 update ESI maps throughout the country. Another is the CAMEO® software suite (EPA) which helps emergency planners 753 and responders deal with chemical incidents. ADIOS (Automated Data Inquiry for Oil Spills), developed by NOAA, 754 provides rapid analysis of how different oil types weather in various marine conditions. By predicting how oil properties 755 change (e.g., evaporation, dispersion), ADIOS helps responders plan effective cleanup strategies. GNOME (General NOAA 756 Operational Modeling Environment) is a critical software suite developed by NOAA to predict the movement and fate of oil 757 spills in water. It incorporates information from forecast systems, like currents and winds to forecast spill trajectories, while 758 also modeling the weathering processes that alter oil's properties over time. Through its components like WebGNOME, PyGNOME, and the ADIOS oil database, GNOME provides mapping and visualization tools, enabling responders to assess 759 760 situations, plan contingencies, and minimize environmental impact. It uses output from various forecast systems produced by 761 the NOAA/NWS's (National Weather Services) Environmental Modeling Center including RTOFS (Real-Time Ocean

- 91
- Forecast System), GFS (Global Forecast System), among others and serves as a vital tool for real-time emergency response,
   contingency planning, and research related to oil spill science.

764 In Canada, while the CCG is the lead agency for coordinating responses to oil spills, their principle is that the 'polluter' pays 765 and should report the spill, take the initial action and fund the cleanup. Industry-funded response organizations, certified by 766 Transport Canada, provide spill response services on behalf of the polluter that would include modelling systems that predict 767 the trajectory and fate of spilled oil.

- 768
- 769 Search and Rescue

770 NOAA's National Environmental, Satellite, Data, and Information Services (NESDIS) Line Office operates the Search And 771 Rescue Satellite Aided Tracking (SARSAT) System to detect and locate people in distress. Mariners, aviators, and 772 recreational enthusiasts can all access the satellite system in an emergency using a portable radio transmitter that can send an 773 SOS signal from anywhere on earth, at any time, including in most extreme weather conditions. This is coupled with the 774 Search and Rescue Optimal Planning System (SAROPS) tool, used by the USCG for maritime search planning. SAROPS 775 uses an Environmental Data Server (EDS) that ingests real-time and forecast environmental data (produced by agencies such 776 as NOAA) to predict the drift of a person or object in the water. This is done by simulating thousands of possible drift 777 scenarios providing probability maps that help to focus the search efforts. The success of this tool is strongly dependent on 778 the quality of the forecast models that it ingests.

The Canadian Coast Guard makes use of observations and models produced by Fisheries and Oceans Canada (DFO) and weather and oceanographic forecasts produced by the ECCC in order to optimize their search operations.

- 781
- 782 Water quality

783 Several U.S. government agencies are involved in supporting marine water quality. Key agencies include (a) the 784 Environmental Protection Agency (EPA), which sets water quality standards, regulates pollutants, and monitors coastal and 785 marine waters; (b) the National Oceanic and Atmospheric Administration (NOAA), which conducts research on ocean 786 health, manages marine resources, and supports programs like the National Estuarine Research Reserve System; (c) the U.S. 787 Coast Guard (USCG), which Coast Guard monitors and responds to marine pollution incidents and ensures maritime safety; 788 (d) the U.S. Army Corps of Engineers (USACE), which Corps manages coastal projects and assesses impacts on water 789 quality from dredging and construction; (e) the Fish and Wildlife Service (FWS), which FWS protects fish and wildlife 790 habitats and works to restore ecosystems, which directly impacts water quality; and (e), the National Park Service (NPS): 791 The NPS manages marine protected areas and conducts water quality monitoring within national parks.

Ocean forecast systems play a key role in monitoring and managing water quality in North America, particularly in coastal
 and nearshore areas. Various water quality models are used by the EPA (<u>https://www.epa.gov/beaches/models-predicting-</u>
 <u>beach-water-quality</u>). These incorporate hydrodynamic forecasts that that are essential for accurately simulating the transport
 and mixing of pollutants.

## 796 8.2 Natural Resources and Energy

#### 797 Fisheries

798 Both the U.S. National Marine Fisheries Service (NMFS) as well as Fisheries and Oceans Canada (DFO) heavily rely on 799 numerical ocean models to support their operations, particularly for fisheries management and protected species 800 conservation. NMFS use models like HYCOM (Hybrid Coordinate Ocean Model) and RTOFS (Real-Time Ocean Forecast 801 System), while DFO use HYCOM as well as regionally tailored models developed by them and in collaboration with ECCC. 802 These models provide crucial data on ocean currents, temperature, and salinity, enabling predictions of fish distribution and 803 marine species movements as well as assessments of habitat suitability. This information is then used to set sustainable catch 804 limits, protect endangered species from human activities, and forecast environmental impacts, thereby informing critical 805 decisions regarding the management and preservation of marine resources.

The NMFS disseminates information through a variety of channels, including their official website (http://fisheries.noaa.gov/), scientific publications, and direct communication with stakeholders. They provide online access to oceanographic data, habitat suitability maps, and species distribution forecasts, ensuring that researchers, resource managers, and the public have access to vital information. NMFS also collaborates with other agencies and organizations to share data and findings, fostering a collaborative approach to marine resource management.

#### 811 Recreation and Tourism

In the US, NOAAs Operational Forecasts Systems (OFS) as well as the NWSs maritime forecasts cover various regions (including the Great Lakes) and provide information on water-levels, currents temperature, salinity and winds that are essential for safe navigation, recreational boating and fishing. The Regional Ocean Modelling System is used by various institutes to provide high resolution forecasts for specific regions, for example the Gulf of Maine Operational Forecast System (GoMOFS) uses ROMS to predict ocean conditions to support tourism and marine recreational activities.

817 In Canada, CONCEPTS as well as the Regional Ice Ocean Prediction System (RIOPS) are used to support tourism by 818 providing forecasts that support safe navigation, ice prediction and ecosystem modelling. A Port Ocean Prediction System 819 (POPS) is being developed by the DFO for major Canadian ports and waterways and provides high resolution forecasts that 820 support marine recreation.

The forecast information is provided through a number of different apps, some examples are: the NOAA Weather Radar &
Live Alerts, PredictWind, Windy, SailFlow, Surfline, MagicSeaweed.

823 Offshore Energy

824 For the offshore energy sector in North America, ocean forecast systems are essential to ensure the safety and efficiency of 825 operations, particularly for oil, gas, and renewable energy projects like offshore wind farms. These systems provide critical 826 information on ocean currents, waves, winds, and other environmental conditions. In addition, research centers, like the 827 National Renewable Energy Laboratory (NREL) and Woods Hole Oceanographic Institution, produce specialized models for 828 specific energy projects. Hindcast data help model historical ocean conditions, and operational forecasts aid in planning and 829 real-time decision-making. Companies like Fugro, Woods Hole Group, DNV GL, and RPS Group offer tailored ocean 830 forecasting and metocean services that provide high-resolution, localized ocean and weather forecasts to support the offshore 831 energy industry. These forecasts are often customized for specific platforms, rigs, or turbines.

The oil and gas energy industry have specific ocean forecast requirements depending on the application, such as diver operations, unmanned vehicles operations, rig installation, production, etc. In the Gulf of Mexico, a leading area for exploration and production, the Loop Current Eddy (LCE) shedding is a process of great interest, as current speeds of extended or detached LCE's often have current speeds in excess of 2-3 m/s, speeds which often require repositioning of equipment or temporary cessation of operations.

#### 837 8.3 Shipping, Ports and Navigation

838 With the advent of new standards for marine navigation. Implementations and applications of ocean prediction systems for 839 E-Navigation and port management are expanding in North America. In the US NOAA's Physical Oceanographic Real-840 Time System (PORTS) provides real-time water level, current, and meteorological information for major U.S. ports and 841 harbours. While the National Operational Coastal Modeling Program (NOCMP) develops and operates a network of 842 Operational Nowcast and Forecast Hydrodynamic Model Systems (OFS) for critical U.S. ports, harbors, and coastal waters. 843 These systems provide predictions of water levels, currents, and other oceanographic variables, aiding in navigation, harbor 844 management, and coastal hazard mitigation. In Canada, CONCEPTS (ECCC/DFO) provides oceanographic forecasts for 845 various regions, including the St. Lawrence Seaway and major Canadian ports and the DFO is developing the Port Ocean 846 Prediction System (POPS) for major Canadian ports and waterways.

These forecasts are starting to be integrated into various Vessel Traffic Management Systems (VTMS) that are used throughout North America. For example, the Canadian Coast Guards Vessel Traffic Services (VTS) are increasingly using data from CONCEPTS and other forecast models and Port specific VTMS in the US (e.g. the Port of New York and New Jersey) integrate data from NOAAs Operational Forecast System.

#### 851 8.4 Climate Adaptation

852 The United States leverages ocean models extensively to bolster climate adaptation strategies for both coastal and ecosystem resilience. A network of federal agencies, including NOAA, EPA, USFWS (U.S. Fish and Wildlife Service), NPS (National 853 854 Park Service), USACE (Army Corps of Engineers), DOI (Department of the Interior), and FEMA (Federal Emergency 855 Management Agency), utilize these models to understand and respond to the impacts of climate change on marine 856 environments. NOAA plays a central role, conducting research on ocean temperature, sea-level rise, and habitat changes, 857 while also collecting and disseminating crucial data to stakeholders. Models provide critical information on sea-level rise, 858 coastal erosion, extreme weather events, and ocean warming, informing the development of resilience strategies and 859 enabling communities, governments, and industries to make informed decisions.

860 Specifically for ecosystem resilience, ocean models support a variety of ecological and biological studies. Agencies like 861 NOAA, through programs like NMFS and OAR (Office of Oceanic and Atmospheric Research), and USFWS, with its 862 Endangered Species Program and National Wildlife Refuge System, use model outputs to monitor marine biodiversity, track 863 species, understand ecosystem dynamics, and manage resources. These models, providing real-time and forecasted data on 864 ocean conditions, help researchers study the effects of climate change, track biological events, and inform conservation and 865 restoration efforts, including those focused on coral reefs and endangered species. Furthermore, for coastal resilience, these 866 models are essential for engineering projects, providing critical predictions of oceanographic and atmospheric conditions that 867 inform the design and maintenance of coastal infrastructure, erosion management, and preparedness for extreme events. In 868 particular, the USGS provides a suite of tools for predicting coastal changes, especially during storms. These tools forecast 869 factors like coastal erosion, overwash, and inundation, which help engineers evaluate potential changes in shoreline position

and design resilient coastal infrastructure. Their Coastal Change Hazards Portal integrates data on sea-level rise, coastal erosion, and sediment transport, which are critical for long-term coastal engineering projects.

872

## 873 **9 Arctic**

874 The Arctic environment is evolving quickly. Short-term models allow users to monitor changes to the landscape, particularly 875 at the ice edge and responses to short-term events (such as storms). This information is valuable for national environment agencies, especially those with Arctic coastlines. As detailed in Cirano et al., 2025, there are a number of short-term (up to 876 877 10 day) forecasting systems available in the Arctic. Nine of these are global models, eight are regional, and five are coastal. 878 It is important to note that many of the Arctic forecast system outputs are used as inputs to other models. This can be specific 879 modelling in response to an event - for example, oil spill trajectory modeling, as described in Nordam et al. (2019) - or for 880 monitoring the state of a specific parameter that is not present in the main forecasting system, such as the use of TOPAZ4 to 881 force a coastal 800 m resolution ocean model for a weekly monitoring and assessment of the sea-louse 882 (https://www.globalseafood.org/advocate/norwegian-researchers-develop-sea-lice-tracking-model/). The latter example is 883 currently only applied to the coastline of mainland Norway at present, but as fishing extends further and further north, such 884 forecasts may also become more relevant further into the Arctic.

They are also used to feed into weather forecast models, an Arctic-specific application mirroring the standard process of forcing ocean models with weather forecast outputs that is often used in other regions. This is because ice conditions can have important feedback to the atmosphere, and models developed specifically for ice can represent these conditions well. The NOAA (the US National Ocean and Atmospheric Administration) ice drift is primarily used for this purpose (https://mag.ncep.noaa.gov/docs/NCEP\_PDD\_MAG.pdf), to provide sea ice conditions for the NWS (the US National Weather Service) global atmospheric model: this has been the case since 1998.

In the following subsections, the other main applications of Arctic forecasts are provided, focusing on direct applications of the forecasts themselves. Note that in most cases the downstream applications are suggested by providers but there is little data available in the public domain about user uptake for a given usage.

## 894 9.1 Extremes, Hazards and Safety

895 As more activities happen at the ice edge and in the marginal ice zone, there is an increase in the risk of both harm to humans 896 and negative consequences of their activities, and there have been some incidents in the last decade (for example, 897 https://barentsobserver.com/en/nature/2013/09/tanker-accident-northern-sea-route-09-09). Marchenko et al. (2015) note "the main operational risk factors faced include geographical remoteness, climate-change related aspects and weather, electronic 898 899 communications challenges, sea ice, lack of precise maps or hydrographic and meteorological data". Forecasting models can 900 be used both to reduce risk and to target the response to an incident. For example, the Barents-2.5km model, used by MET 901 Norway, acts as one of the main inputs to further modeling of pollutants (such as drift of oil spills from ships) and iceberg 902 drifting, which are all based on the same type of Lagrangian drift calculations (Sutherland et al., 2020). It is also used in 903 search and rescue operations, where information on where a lost person or vessel may drift in the short term is very 904 important.

905 Storm Surge

906 Coastal models play an important role in understanding the short-term behaviour of a region. One such example is the storm 907 surge model, which provides both coastal forecasts (useful for those with activities in coastal waters, such as fishing) and a 908 warning system for storm surges along the coast of mainland Norway and Svalbard. Users receive an alert when an extreme 909 weather event is likely; for example, during the storm "Elsa" in February 2020, it was found to be a useful tool to both 901 monitor the development and to send warnings out (Kristensen et al., 2024).

## 911 9.2 Natural Resources and Energy

As sea ice declines, more opportunities to exploit natural resources such as oil and gas extraction arise, although the safety of fixed assets and persons will still be at risk of storms, high waves, sea ice and incoming icebergs. To reduce ocean pollution and carbon footprint from transportation of people/resources to and from destinations, as well as minimise risk from ending up in thick ice, companies must choose the best routes for transportation. Short-term forecasts in conjunction with available real-time observations can be very important for this (Grigoryev et al., 2022). While no specific operational downstream applications have been identified in this category for the Arctic, in the sections below are described the growing needs specific to the region.

919 Fisheries

920 The Agreement to prevent Unregulated High Seas Fisheries in the central Arctic Ocean has been in place since 25th June 921 (<u>https://arctic-council.org/news/introduction-to-international-agreement-to-prevent-unregulated-fishing-in-the-high-922 seas-of-the-central-arctic-ocean/</u>) and aims to ensure that future fishing in the Arctic as sea ice declines can be carried out 923 sustainably.

Short-term forecasts could help to support this agreement as well as to inform users about conditions suited to fish stocks and to reduce the chance of operating in risky conditions which could lead to oil spills. As noted by Neis et al., (2020), "When harvesters adjust their activity or move into new fishing grounds, forecasts become critical tools for anticipating dangerous conditions and 'learning' an unknown environment or working context (e.g., different gear)", which suggests that even if the central Arctic Ocean remains tightly controlled, an increase in fishing activities in the northern peripheral seas as ice declines (Fauchald et al., 2021) may increase the need for forecasts of environmental conditions for a new set of users in the future

930

931 Tourism

Arctic tourism has been increasing in recent decades (Larsen and Fondahl, 2014f), particularly the concept of "last chance tourism" (Eijgelaar et al., 2010). As well as requiring forecasts for navigation in waters where ships have been built for comfort rather than operational purposes, tourism is often focused on reaching the ice edge or ecosystems to spot wildlife. This can require accurate forecasts of sea ice conditions and the limit of the Marginal Ice Zone which is a hotspot for biological activity in the Arctic (and attracts the more audacious fishermen as a result). Search and rescue-based forecasts for such purposes is also relevant as ships aim to get close to the ice rather than avoid it.

938

## 939 9.3 Shipping, Ports and Navigation

Reductions in summer sea ice, and thinner ice, open new routes to traverse the Arctic (for example, the Northeast Passage),
providing more efficient routes across the globe, as well as providing opportunities for many of the above users to work
further into the Arctic Basin away from the coast. In all the cases currently described, there is an aspect of navigation driving

104

943 a need for forecasts. One of the main considerations when navigating is sea ice jams and ice accumulation, which can 944 prevent further progress to ships and cause hull damage (for example, the case where two cargo ships were stuck and 945 damaged in Frobisher Bay, https://www.cbc.ca/news/canada/north/ ice-damages-hull-of-sealift-ship-near-igaluit-1.1230034). 946 Depending on the ability of the ship (ice-strengthened or icebreaker), different sea ice conditions can be the limit of safe 947 operations. Given the ice can vary quickly, recent efforts have been made to include a dynamical ice edge in fully coupled 948 model for weather prediction (Day et al., 2022) and improve forecasts of the ice edge itself (Posev et al., 2015) A typical use 949 of sea ice short-term forecasts is to assess whether the ice edge is advancing or retreating (which would then feed into 950 decisions related to navigation on the short-term, such as whether or not it is safe for a ship to either stay in a given location 951 for deployments, or to navigate in a certain direction; for example, the use of VENUS for monitoring sea ice in Bering Strait, 952 Cirano et al., 2025). One of the main limitations of accessing information from a ship is a reliable internet connection, 953 meaning forecasts must be readily available and not hard to download. A number of users still rely on manual ice charts 954 drawn by experts.

955 Ship operators rely on operational forecast models to adhere to the Polar Code, which is the International Marine 956 Organisation's international code for ships operating in polar waters, in place since 1st January 2017 957 (https://www.imo.org/en/ourwork/safety/pages/polar-code.aspx); it is relevant for navigation (and, as part of this, design and 958 capabilities of ships wishing to work in polar waters) and operational procedures, search and rescue, and protection of 959 ecosystems. Mandatory measures cover safety and pollution prevention, and ships going into the polar regions require a 960 Polar Ship Certificate determining what conditions the ship is suited to 961 (https://www.dnv.com/maritime/polar/requirements.html). Forecasts can contribute to helping users abide by the Code, for 962 example by assessing whether ships will be able/authorized to operate in upcoming sea-ice conditions. The definition of 963 "environmental conditions" is evolving in the Polar Code and may in the future include variables that can be skillfully 964 forecast.

965 Ultimately, all ship-based operations in the Arctic region rely on navigation and sea ice information for navigation, either to 966 avoid or get close to the ice edge, and this is the most mature of the forecast applications. Tools exist to condense or combine 967 multiple forecast outputs and observations to provide near-real time and forecasted conditions in a user-friendly way. Two 968 such examples are IcySea (<u>https://driftnoise.com/icysea/</u>), which uses ice charts with a sea ice drift forecast, and Activities 969 (<u>https://arctivities.noveltis.fr/overview/</u>), which provides a risk index and anthropic noise levels. Such tools can be used to 970 support maritime users with varying needs.

## 971 Research Support

972 Forecasts of the Arctic Ocean can be used to inform new developments or deployments of equipment for scientific purposes. 973 One such example is the Sea Ice Drift forecast Experiment (SIDFEx<sup>22</sup>). Two of the main aims of the campaign were to 974 gather information on available sea ice drift forecasts in order to a) decide on an optimal starting position for the research 975 icebreaker Polarstern to commence a year-long study of conditions while frozen into the sea ice, and b) use the drift forecasts 976 to inform where to order high-resolution satellite images of the local domain around the ship for the coming days as they 977 become available. Using sea ice drift models to selectively download these images saved limited bandwidth and image fees.

978 Another example of the use of short-term forecasts is the use of the VENUS (VEssel Navigation Unit support System), a 979 forecasting platform which can use a variety of domains to provide forecasts for research ships on demand. This was 980 successfully deployed in a cruise in 2018 (Dethloff et al., 2019). The ice-strengthened ship MIRAI could only go a) where 981 ice thickness was less than 70cm and concentration less than 0.1, and b) where air temperature was greater than -15 degrees

108 109

<sup>107 &</sup>lt;sup>22</sup> <u>https://www.polarprediction.net/key-yopp-activities/sea-ice-prediction-and-verification/sea-ice-drift-forecast-experiment/</u>

C (Inoue, 2019). Scientists were deploying equipment near the marginal ice zone in order to investigate the predictability of conditions during autumn freezing; further, the ship needed to gather as much data as possible while being able to exit through the Bering Strait before ice blocked it for the winter (Dethloff et al., 2019). Using VENUS, which combines forecast from ECMWF, sea ice forecasts from ICEPOM (University of Tokyo) and passive microwave data helped to inform these. Such use of forecasts can also feed back into the development - for example, on the MIRAI cruise, the bandwidth was such that it was hard to download data; therefore 2D fields were more valuable (Inoue, 2019).

988

## 989 9.4 Climate Adaptation

990 The rapidly declining sea ice, environmental changes and potential economic opportunities of the Arctic region have 991 attracted a lot of interest, but with this comes a new state that is still being understood even as it evolves. Large uncertainties 992 in Arctic forecasts somewhat impede their use in climate adaptation, but the strategic and economic interest for the region as 993 well as presence of coastal communities has made it a very active field of research. For example, decadal predictions such as 994 those from the IPCC 6th Assessment Report (https://www.ipcc.ch/synthesis-report/) are used to predict future states, often by 995 selecting some variables in conjunction with past and present in-situ and satellite monitoring to make the predictions more 996 robust and downscaled to more local areas. Examples include frequency of marine heatwaves (He et al., 2024), and sea level 997 rise and coastal erosion (Tanguy et. Al. (2024)). In the Barents Sea, climate prediction models have also been used to predict 998 phytoplankton up to 5 years in advance (Frasner et al., 2024) and cod populations under evolving ocean physical properties 999 (Kjesbu et al., 2022). Such studies can provide new understanding which can contribute to decision-making and planning in vulnerable communities and occupations that are dependent on knowing the physical conditions or biological activity. 1000

1001 Another key tool in developing understanding of the changing Arctic is to use reanalyses or hindcasts to see how the present 1002 situation compares to earlier years. Many of the available short-term forecasts in the Arctic (Cirano et al., 2025) have an 1003 accompanying reanalysis or hindcast so that past seasonal evolution of relevant conditions. For some maritime users, 1004 seasonal predictions can supplement this information to aid voyage planning (Wagner et al. (2020), for both safety and 1005 ensuring adherence to the Polar Code (see section 9.3). An additional example is the Disko Bay model run by the Disko Ice 1006 and Ocean service (https://marine.copernicus.eu/services/use-cases/monitoring-ecosystem-within-disko-bay), which provides 1007 both forecasts and a hindcast of ocean conditions at the high resolution required for Greenlandic fiord environments, using 1008 output from a lower-resolution forecasting model as boundary conditions. Outputs from this fjord model have been provided 1009 to an ecosystem model; these applications contribute to monitoring efforts to ensure long-term sustainability of the Blue 1010 Economy in Greenland.

## 1011 **10. Education, stakeholder engagement and ocean literacy**

Education, stakeholder engagement and ocean literacy activities are essential components in supporting the full value chain from data production (operational forecast systems) to the provision of useful downstream applications. These activities are carried out in all regions and at various different stages along the value chain: from education outreach activities with learners, technical workshops, to community engagement and co-design workshops with stakeholder groups. They help to ensure that the downstream applications produced have real value and are measurably impactful. Below we provide some examples of the types of education and engagement activities that take place.

1018 Technical Workshops

1019 The Sub-Commision of the Intergovernmental Oceanographic Commission of the United Nations Educational, Scientific and 1020 Cultural Organization for the Western Pacific and adjacent seas (WESTPAC) develops and strengthens regional and 1021 Member States' capacity for ocean model development, data assimilation, model validation, and development of Ocean 1022 Forecasting System, through a series of national and regional trainings, scientific workshops, and professional exchanges 1023 among partner institutions (https://ioc-westpac.org/ofs/capacities/). The Regional Training and Research Center on Ocean 1024 Dynamics and Climate (RTRC-ODC) was officially established at the 8th WESTPAC Intergovernmental Session in 2010. 1025 The First Institute of Oceanography, State Oceanic Administration of China, organized the lecture series on ocean models 1026 (2011), ocean dynamics (2012), air-sea interaction and modeling (2013), climate models (2014), climate change (2015), 1027 ocean dynamics and multi-scales interaction (2016), development of coupled regional ocean models (2017), ocean forecast 1028 system (2018) and climate dynamics and air-sea interactions (2019). In the evaluation period of 2015-2019, 191 young 1029 scientists from 36 countries joined the lectures (https://ioc-westpac.org/rtrc/odc/).

## 1030 Ocean Literacy

1031 With ongoing Arctic Sea ice decline, scientific results from the region are more frequently appearing in national news and 1032 the general public are more aware of the Arctic environment and how it is changing. The freely accessible forecast maps 1033 from most services, with an interface that can select given variables and watch as they run forward in time, provide a useful 1034 tool to demonstrate how changeable, for example, the ice edge is in response to forcing even on the short term, which can be 1035 used to engage with wider audiences and educate about the Arctic as a dynamic system. For example, Coursera, a website 1036 offering a number of free online courses for studying in evenings, has a course entitled "Frozen in the Ice: Exploring the 1037 Arctic", based out of the University of Boulder, Colorado (https://www.coursera.org/learn/frozen-in-the-ice); the course 1038 allows participants to act as virtual participants on the MOSAiC Arctic research campaign, and one of the six modules is 1039 based around Arctic forecasting. Activities such as this allow the public to get closer to polar research, and many large 1040 research campaigns now include outreach as part of their programs.

#### 1041 Stakeholder Engagement and Co-Design

1042 With NOAAs Office of Response and Restoration, the Emergency Response Division (ERD) develops tools, guidelines, and 1043 small, field-oriented job aids to assist preparedness for response communities. In addition, NOAA provides standard 1044 techniques for observing oil, assessing shoreline impact, and evaluating and selecting cleanup technologies that have been 1045 widely accepted by response agencies.

South Africa's National Oceans and Coastal Information Management System (OCIMS) holds annual stakeholder
 engagement workshops that facilitates the co-design of the decision support tools. Between the workshops, dialogue between
 stakeholders and developers is maintained through active whattsapp groups.

While INCOIS provides extensive training to users for efficient utilization of their forecast products, they have noticed that NGOs, Universities, local government departments and localized user community networks are found to be very effective in ensuring that the information reaches the user in time. User-uptake is supported by their good relationship with local fishing communities who are involved with the safe-keeping of their observation platforms in exchange for timely warnings of maritime hazards. This relationship builds awareness as well as trust with coastal communities.

1054 Citizen Science

1055 New Zealand's Moana Project innovatively incorporates citizen science by partnering with commercial fishers to gather 1056 essential oceanographic data. Fishing vessels are equipped with the "Mangōpare" sensor system, which automatically 1057 collects and transmits subsurface temperature measurements in near real-time as the vessels go about their normal fishing 1058 activities. This transforms the fishing fleet into a vast, mobile observation network, expanding data coverage across a wider

- 114 115
  - 5

spatial range than traditional research methods. This mutually beneficial partnership provides scientists with valuable data, while fishers gain access to information that can enhance their own operations. By empowering local communities and increasing data accessibility, Moana fosters collaboration and contributes to a deeper understanding of the marine environment, ultimately supporting sustainable fisheries management and scientific research.

#### 1063 11. Summary

Operational oceanography supports the Blue Economy, providing the knowledge and tools for us to sustainably use our oceans for economic growth, better livelihoods, and job creation. Around the world, scientists and forecasters are developing cutting-edge tools that transform raw ocean data into practical solutions for a variety of challenges. These tools help us understand and protect our marine environments, manage resources, and ensure safety at sea.

1068 This report has provided some examples of Downstream Applications, based on operational forecast systems, for eight of the 1069 nine regional teams, identified by the OP DCC. It is by no means a comprehensive review, but it does provide an indication 1070 of the needs and services in each region as well as the relative maturity level of downstream applications. The regions with 1071 the most established and most numerous operational forecast systems (e.g. the Mediterranean and Black Sea; North East 1072 Atlantic; North America; parts of the Western Pacific and Asia and to some extent the Arctic) tend to also have the most 1073 mature downstream applications. The forecasting systems of the Indian Seas, South America and Africa can be thought of as 1074 'emerging' and by this we mean: new, rapidly growing and often under- or less-resourced. Despite this, the INCOIS system 1075 developed for the Indian Seas is a sophisticated system that incorporates real time observations and provides mature tools for 1076 stakeholders that support various offshore activities. Part of their success is related to their close engagement with their 1077 stakeholders. The African region is one of the least developed in terms of regionally optimized forecast systems, with only a 1078 few developed in various parts of the continent. However, they do have two fairly mature user-support platforms that are 1079 based primarily on earth observations and whose tools are co-designed with stakeholders. These dissemination platforms are 1080 ready to ingest tools based on regionally optimized forecasts.

1081 In this review, a sample of various downstream applications around the Globe reveals that while established and reliable 1082 forecast systems are a key factor in their abundance, a good relationship with stakeholders is critical for their uptake.

1083

1084

#### 1085 References

#### 1086 **References: Introduction**

Alvarez Fanjul, E., Ciliberti, S., Bahurel, P.: Implementing Operational Ocean Monitoring and Forecasting Systems. IOC UNESCO, GOOS-275. https://doi.org/10.48670/ETOOFS, 2022.

#### 1089 References Region 1: West Pacific and Marginal Seas of South and East Asia

1090 Cirano, M., Alvarez-Fanjul, E., Capet, A., Ciliberti, S., Clementi, E., Dewitte, B., Dinápoli, M., El Serafy, G., Hogan, P., Joseph,

1091 S., Miyazawa, Y., Montes, I., Narvaez, D., Regan, H., Simionato, C. G., Tanajura, C. A. S., Thupaki, P., Urbano-Latorre, C., and

- 1092 Veitch, J.: A description of existing Operational Ocean Forecasting Services around the Globe, State Planet Discuss. [preprint],
   1093 https://doi.org/10.5194/sp-2024-26, in review, 2025.
- 1094 Hirose, N., Kumaki, Y., Kaneda, A., Ayukawa, K., Okei, N., Ikeda, S., Igeta, Y., and Watanabe, T.: Numerical simulation of
- the abrupt occurrence of strong current in the southeastern Japan Sea. Continental Shelf Research, 143, 194-205.
   https://doi.org/10.1016/j.csr.2016.07.005, 2017.
- 1097 Nagao, M., Takasugi, Y., Suzuki A., Tanaka, Y., Sugishima, H., Matsui, T., Okamoto, N. .: Confirming the Validity of
- 1098 ADCP Velocity Measurements for Physical Environmental Assessment in Japan's Exploration Areas for Cobalt-Rich
- Ferromanganese Crusts. Proceedings of the Twenty-eighth International Ocean and Polar Engineering Conference, 136-142,2018.
- Nakada, S., Hirose, N., Senjyu, T., Fukudome, K.-I., Tsuji, T., and Okei, N.: Operational ocean prediction experiments for
  smart coastal fishing. Progress in Oceanography, 121, 125-140. <u>https://doi.org/10.1016/j.pocean.2013.10.008</u>, 2014.
- Qiao, F., Wang, G., Khokiattiwong, S. et al.: China published ocean forecasting system for the 21st-Century Maritime Silk
  Road on December 10, 2018. Acta Oceanol. Sin. 38, 1-3. https://doi.org/10.1007/s13131-019-1365-y, 2019.
- 1105 Ramdhani, A.: Development of Indonesia coastal inundation forecasting system (Ina CIFS). Available at
   1106 <u>https://oceanexpert.org/downloadFile/40789</u>, 2019 (last access: 27/07/2024).
- 1107RepublicofKorea/OceanPredict.:KHOA(KOOFS)/KIOST(KOOS).Availableat1108https://oceanpredict.org/docs/Documents/General%20documents/National%20system%20reports/2020-21/National-
- 1109 <u>Report-2020-21-RepofKorea.pdf</u>, 2020 (last access: 27/07/2024).
- 1110 Sato, Y, and Horiuchi, K.: Energy-saving voyage by utilizing the ocean current. IoS-OP Taiwan Seminar, Archives On-
- 1111 demand, 19. Available at https://www.shipdatacenter.com/en/ios-op\_seminar\_twod\_202208, 2022 (last access: 27/07/2024).
- 1112 Schiller, A., Brassington, G. B., Oke, P., Cahill, M., Divakaran, P., Entel, M., ... Zhong, A. (2019). Bluelink ocean
- 1113 forecasting Australia: 15 years of operational ocean service delivery with societal, economic and environmental
- 1114 benefits. Journal of Operational Oceanography, 13(1), 1–18. https://doi.org/10.1080/1755876X.2019.1685834
- 1115 Takikawa, T. et al.: Hydrographic observations by fisherman and coastal ocean model—Salinity variations around Iki Island
- during the cooling season. Journal of Advanced Marine Science and Technology Society, 25, 15-20. (in Japanese with English abstract and figure captions). https://doi.org/10.14928/amstec.25.1 15, 2019.
- 1118 Webb, A., Waseda, T. and Kiyomatsu, K.: A high-resolution, long-term wave resource assessment of Japan with wave-
- 1119 current effects. Renewable Energy, 161, 1341-1358. <u>https://doi.org/10.1016/j.renene.2020.05.030</u>, 2020.
- 1120 Qiao, F., Wang, G., Khokiattiwong, S., Akhir, M. F., Zhu, W., & Xiao, B. (2019). China published ocean forecasting system
- for the 21st-Century Maritime Silk Road on December 10, 2018. Acta Oceanologica Sinica= Hai Yang Hsueh Pao, 38(1), 1-3.

#### 1123 References Region 3: African Seas

- 1124 ABALOBI Impact Report 2018-2019. Available at 1125 https://drive.google.com/file/d/1wbi0PPDOr8oZS b0LMJs5PFv37tOAiv5/view (last access: 27/0772024). 1126 Cambaza, E., Mongo, E., Anapakala, E., Nhambire, R., Singo, J., Machava, E.: Outbreak of Cholera Due to Cvclone 1127 Kenneth in Northern Mozambigue. Int. J. Environ. Res. Public Health, 16(16), 2925. doi: 10.3390/ijerph16162925, 2019. 1128 Cirano, M., Alvarez-Faniul, E., Capet, A., Ciliberti, S., Clementi, E., Dewitte, B., Dinápoli, M., El Serafy, G., Hogan, P., Joseph, 1129 S., Miyazawa, Y., Montes, I., Narvaez, D., Regan, H., Simionato, C. G., Tanajura, C. A. S., Thupaki, P., Urbano-Latorre, C., and 1130 Veitch, J.: A description of existing Operational Ocean Forecasting Services around the Globe, State Planet Discuss. [preprint], 1131 https://doi.org/10.5194/sp-2024-26, in review, 2025. 1132 de Vos, M., Vichi, M., and Rautenbach, C.: Simulating the Coastal Ocean Circulation Near the Cape Peninsula Using a 1133 Coupled Numerical Model. Journal of Marine Science and Engineering, 9(4), 359. https://doi.org/10.3390/jmse9040359, 2021. 1134 1135 Groom, S., Sathyendranath, S., Ban, Y., Bernard, S., Brewin, R., Brotas, V., Brockmann, C., Chauhan, P., Choi, J.-K., 1136 Chuprin, A., Ciavatta, S., Cipollini, P., Donlon, C., Franz, B., He, X., Hirata, T., Jackson, T., Kampel, M., Krasemann, H., 1137 Lavender, S., Pardo-Martinez, S., Mélin, F., Platt, T., Santoleri, R., Skakala, J., Schaeffer, B., Smith, M., Steinmetz, F., 1138 Valente, A., and Wang, M.: Satellite Ocean Colour: Current Status and Future Perspective. Front. Mar. Sci., 6, 485. 1139 https://doi.org/10.3389/fmars.2019.00485, 2019. 1140 Hart-Davis, M.G., and Backeberg, B.C.: Towards a particle trajectory modelling approach in support of South African 1141 search 16(2), and rescue operations at sea. Journal of Operational Oceanography, 131-139. 1142 https://doi.org/10.1080/1755876X.2021.1911485, 2023. 1143 Heve, S.: Surface Circulation in the KwaZulu-Natal Bight and its Impact on the Connectivity of Marine Protected Areas. 1144 University of Cape Town, South Africa, MSc thesis. Available at 1145 https://open.uct.ac.za/bitstream/handle/11427/36460/thesis sci 2022 heve%20sonia.pdf?sequence=1&isAllowed=v. 2021 1146 (last access: 27/07/2024). 1147 Hoteit, I., et al.: Towards an End-to-End Analysis and Prediction System for Weather, Climate, and Marine Applications in 1148 the Red Sea, Bulletin of the American Meteorological Society, 102(1), E99–E122, https://doi.org/10.1175/BAMS-D-19-1149 0005.1, 2021. 1150 Mather, A., and Stretch D.: A perspective on Sea Level Rise and Coastal Storm Surge from Southern and Eastern Africa: a 1151 Case Study Near Durban, South Africa. Water, 4(1), 237-259. https://doi.org/10.3390/w4010237, 2012. 1152 Molua, E.L., Mendelsohn, R.O., Akamin, A.: Economic vulnerability to tropical storms on the southeastern coast of Africa.
- 1153 Jamba, 12(1), 676. doi: 10.4102/jamba.v12i1.676, 2020.
- 1154 Pfaff, M., Hart-Davis, M., Smith, M.E., and Veitch, J.: A new model-based coastal retention index (CORE) identifies bays as
- 1155 hotspots of retention, biological production and cumulative anthropogenic pressures. Estuarine, Coastal and Shelf Science,
- 1156 273, 107909. <u>https://doi.org/10.1016/j.ecss.2022.107909</u>, 2022.

- 1157 Ravela, S., Verly, C., Ludwig, L.C., Neumann, J.E., and Emanuel, K.A.: Assessing the Risk of Cyclone-Induced Storm
- 1158 Surge and Sea Level Rise in Mozambique. UNU-WIDER. Available at <u>https://www.wider.unu.edu/publication/assessing-</u>
- 1159 <u>risk-cyclone-induced-storm-surge-and-sea-level-rise-mozambique</u>, 2013 (last access: 27/07/2024).
- Seveso, D., Louis, Y.D., Montano, S., Galli, P., Saliu, F.: The Mauritius Oil Spill: What's Next? Pollutants, 1(1), 18-28.
  <a href="https://doi.org/10.3390/pollutants1010003">https://doi.org/10.3390/pollutants1010003</a>, 2021.
- Singh, M., and Schoenmakers, E.: Comparative Impact Analysis of Cyclone Ana in the Mozambique Channel Using
  Satellite Data. Applied Science, 13(7), 4519. https://doi.org/10.3390/app13074519, 2023.
- Smith, M. E., Lain, L. R., and Bernard, S.: An optimized chlorophyll a switching algorithm for MERIS and OLCI in
  phytoplankton-dominated waters. Remote Sensing of Environment, 215, 217-227. <u>https://doi.org/10.1016/j.rse.2018.06.002</u>,
  2018.
- Troch, C., Terblanche, L. and Roussouw, M.: Moored Ship Motion Forecast Tool for the Port of Ngqura. Coastal
  Engineering Proceedings,67. <u>http://dx.doi.org/10.9753/icce.v37.structures.67</u>, 2024 .
- 1169 Uba, F., Essandoh, E.O., Nyantakyi, E.K. and Anumah, P.: Hydrodynamic Model for Operational Forecasting in Coastal
- 1170 Waters of Ghana. Open Journal of Modelling and Simulation, 8, 48-59. <u>https://doi.org/10.4236/ojmsi.2020.82004</u>, 2020

#### 1171 References Region 4: Mediterranean and Black Sea

- Alvarez Fanjul, E., Ciliberti, S., Bahurel, P.: Implementing Operational Ocean Monitoring and Forecasting Systems. IOC UNESCO, GOOS-275. https://doi.org/10.48670/ETOOFS, 2022.
- 1174 Álvarez Fanjul, E., Sotillo, M.G., Perez Gomez, B., Valdecasas, M.G., Perez Rubio, S., Lorente, P., Dapena, A.R., Martinez
- 1175 Marco, I., Luna, Y., Padorno, E., Santos Atienza, I., Hernandez, G.D., Lopez Lara, J., Medina, R., Grifoll, M., Espino, M.,
- 1176 Mestres, M., Cerralbo, P., and Sanchez Arcilla, A.: Operational oceanography at the service of the ports. In "New Frontiers
- in Operational Oceanography", E. Chassignet, A. Pascual, J. Tintoré, and J. Verron, Eds., GODAE OceanView, 729-736.
  https://doi.org/10.17125/gov2018.ch27, 2018.
- 1179 Brovchenko, I., Kuschan, A., Maderich, V., Shliakhtun, M., Yuschenko, S., and Zheleznyak, M.: The modelling system for
- simulation of the oil spills in the Black Sea. Elsevier Oceanography Series, 69, 586-591. <u>https://doi.org/10.1016/S0422-</u>
  <u>9894(03)80095-8</u>, 2003.
- 1182 Bruschi, A., Lisi, I., De Angelis, R., Querin, S., Cossarini, G., Di Biagio, V., et al.: Indexes for the assessment of bacterial
- 1183 pollution in bathing waters from point sources: The northern Adriatic Sea CADEAU service. Journal of Environmental
- 1184 Management, 293, 112878. <u>https://doi.org/10.1016/j.jenvman.2021.112878</u>, 2021.
- Capet, A., Stanev, E.V., Beckers, J.M., Murray, J.W., and Grégoire, M.: Decline of the Black Sea oxygen inventory.
  Biogeosciences, 13, 1287-1297. https://doi.org/10.5194/bg-13-1287-2016, 2016.

- 1187 Ciliberti, S.A.. Grégoire, M.. Staneva, J.. Palazov, A.. Coppini, G.. Lecci, R.. Peneva, E.. et al.: Monitoring and Forecasting
- the Ocean State and Biogeochemical Processes in the Black Sea: Recent Developments in the Copernicus Marine Service. J.
  Mar. Sci. Eng., 9, 1146. https://doi.org/10.3390/imse9101146, 2022.
- 1190 Coppini G., Fanjul, E.A., Crosnier, L., Dabrowski, T., Daniel, P., Leitão, J.C., Liubartseva, S., Mannarini, G., and Nolan, G.:
- 1191 Chapter 11, Downstream applications: From data to products. In Implementing Operational Ocean Monitoring and 1192 Forecasting Systems, Eds: Alvarez Fanjul, E., Ciliberti, S., Bahurel, P., 2022, IOC-UNESCO, GOOS-275. 1193 https://doi.org/10.48670/ETOOFS, 2022.
- 1194 Coppini, G., Clementi, E., Cossarini, G., Salon, S., Korres, G., Ravdas, M., Lecci, R., Pistoia, J., Goglio, A. C., Drudi, M.,
- 1195 Grandi, A., Aydogdu, A., Escudier, R., Cipollone, A., Lyubartsev, V., Mariani, A., Cretì, S., Palermo, F., Scuro, M., Masina,
- 1196 S., Pinardi, N., Navarra, A., Delrosso, D., Teruzzi, A., Di Biagio, V., Bolzon, G., Feudale, L., Coidessa, G., Amadio, C.,
- Brosich, A., Miró, A., Alvarez, E., Lazzari, P., Solidoro, C., Oikonomou, C., and Zacharioudaki, A.: The Mediterranean
  forecasting system. Part I: evolution and performance. EGUsphere [preprint], <u>https://doi.org/10.5194/egusphere-2022-1337</u>,
  2023.
- 1200 Coppini, G., Jansen, E., Turrisi, G., Creti, S., Shchekinova, E. Y., Pinardi, N., Lecci, R., Carluccio, I., Kumkar, Y.V.,
- D'Anca, A., Mannarini, G., Martinelli, S., Marra, P., Capodiferro, T. and Gismondi, T.: A new search-and-rescue service in
  the Mediterranean Sea: a demonstration of the operational capability and an evaluation of its performance using real case
  scenarios. Nat. Hazards Earth Syst. Sci., 16, 2713-2727, <a href="https://doi.org/10.5194/nhess-16-2713-2016">https://doi.org/10.5194/nhess-16-2713-2016</a>, 2016.
- 1204 Cossarini, G., Querin, S., Solidoro, C., Sannino, G., Lazzari, P., Di Biagio, V., and Bolzon, G.: Development of 1205 BFMCOUPLER (v1.0), the coupling scheme that links the MITgcm and BFM models for ocean biogeochemistry 1206 simulations, Geosci. Model Dev., 10, 1423-1445. <u>https://doi.org/10.5194/gmd-10-1423-2017</u>, 2017
- 1207 Cucco, A., Ribotti, A., Olita, A., Fazioli, L., Sorgente, B., Sinerchia, M., Satta, A., Perilli, A., Borghini, M., Schroeder, K.,
  1208 and Sorgente, R.: Support for oil spill emergencies in the Bonifacio Strait, western Mediterranean. Ocean Sci., 8, 443-454,
  1209 <u>https://doi.org/10.5194/os-8-443-2012</u>, 2012.
- Daniel, P.: Operational forecasting of oil spill drift at Meteo-France. Spill Science & Technology Bulletin, 3(1-2), 53-64.
  https://doi.org/10.1016/S1353-2561(96)00030-8, 1996.
- Dayan, H., McAdam, R., Juza, M., Masina and S., Speich, S.: Marine heat waves in the Mediterranean Sea: An assessment from the surface to the subsurface to meet national needs. Frontiers in Marine Science, 10, 045138. https://doi.org/10.3389/fmars.2023.1045138, 2023.
- 1215 De Dominicis, M., Pinardi, N., Zodiatis, G., and Lardner, R.: MEDSLIK-II, a Lagrangian marine surface oil spill model for
- 1216 short-term forecasting Part 1: Theory. Geosci. Model Dev., 6, 1851-1869, <u>https://doi.org/10.5194/gmd-6-1851-2013</u>, 2013.
- 1217 Ferrarin, C., Penna, P., Penna, A., Spada, V., Ricci, F., Bilić, J., Krzelj, M., Ordulj, M., Sikoronja, M., Duračić, I.,
- 1218 Iagnemma, L., Bućan, M., Baldrighi, E., Grilli, F., Moro, F., Casabianca, S., Bolognini, L., Marini, M.: Modelling the 1219 guality of bathing waters in the Adriatic Sea. Water, 13(11), 1525, https://doi.org/10.3390/w13111525, 2021.
- 129 130

- 1220 Fiori, E., Zavatarelli, M., Pinardi, N., Mazziotti, C. and Ferrari, C.R.: Observed and simulated trophic index (TRIX) values
- 1221 for the Adriatic Sea basin. Nat. Hazards Earth Syst. Sci., 16, 2043-2054, <u>https://doi.org/10.5194/nhess-16-2043-2016</u>, 2016.
- 1222 García-León, M., Sotillo, M.G., Mestres, M., Espino, M., Alvarez Fanjul, E.: Improving Operational Ocean Models for the
- 1223 Spanish Port Authorities: Assessment of the SAMOA Coastal Forecasting Service Upgrades. J. Mar. Sci. Eng., 10(2), 149. 1224 https://doi.org/10.3390/imse10020149, 2022.
- 1225 Gkanasos, A., Schismenou, E., Tsiaras, K., Somarakis, S., Giannoulaki, M., Sofianos, S., and Triantafyllou, G.: A three
- dimensional, full life cycle, anchovy and sardine model for the North Aegean Sea (Eastern Mediterranean): Validation,
  sensitivity and climatic scenario simulations. Mediterranean Marine Science, 22(3). <u>https://doi.org/10.12681/mms.27407</u>,
  2021.
- Gonzalez-Fernandez, D., Hanke, G., Pogojeva, M., Machitadze, N., Kotelnikova, Y., Tretial, I., Savenko, O., Bilashvili, K.,
  Gelashvili, N., Fedorov, A., Kulagin, D., Terentiev, A., and Slobodnik, J.: Floating marine macro litter in the Black Sea:
  Toward baselines for large scale assessment. Environmental Pollution, 309, 119816.

1232 <u>https://doi.org/10.1016/j.envpol.2022.119816</u>, 2022.

- 1233 Hatzonikolakis, Y., Tsiaras, K., Theodorou, J.A., Petihakis, G., Sofianos, S., Triantafyllou, G.: Simulation of mussel Mytilus
- 1234 galloprovincialis growth with a dynamic energy budget model in Maliakos and Thermaikos Gulfs (Eastern Mediterranean).
- 1235 Aquaculture Environment Interactions, 9, 371-383. <u>https://doi.org/10.3354/aei00236</u>, 2017.
- 1236 Kaandorp, M., Dijkstra, H.A., and Van Sebille, E.: Closing the Mediterranean marine floating plastic mass budget: inverse 1237 & modeling of sources and sinks. Environmental Science Technology, 54(19), 11980-11989. 1238 https://pubs.acs.org/doi/10.1021/acs.est.0c01984, 2020.
- Berline, L., Zakardjian, B., Molcard, A., Ourmières, Y., and Guihou, K.: Modeling jellyfish Pelagia noctiluca transport and
  stranding in the Ligurian Sea. Marine Pollution Bulletin, 70(1-2), 90-99. <u>https://doi.org/10.1016/j.marpolbul.2013.02.016</u>,
  2013.
- Laurent, C., Querin, S., Solidoro, C., and Canu, D.M.: Modelling marine particle dynamics with LTRANS-Zlev:
  Licerimplementation and validation. Environmental Modelling and Software, 125, 104621.
  <u>https://doi.org/10.1016/j.envsoft.2020.104621</u>, 2020.
- 1245 Ličer, M., Estival, S., Reyes-Suarez, C., Deponte, D., and Fettich, A.: Lagrangian modelling of a person lost at sea during the
- Adriatic scirocco storm of 29 October 2018. Nat. Hazards Earth Syst. Sci., 20, 2335-2349. <u>https://doi.org/10.5194/nhess-20-</u>
  2335-2020, 2020.
- Liubartseva, S., Coppini, G., Lecci, R., and Clementi, E.: Tracking plastics in the Mediterranean: 2D Lagrangian model.
  Mar. Pollut. Bull., 129(1), 151-162. https://doi.org/10.1016/j.marpolbul.2018.02.019, 2018.
- 1250 Liubartseva, S., Federico, I., Coppini, G., and Lecci, R.: Stochastic oil spill modeling for environmental protection at the
- 1251 Port of Taranto (southern Italy). In Mar. Pollut. Bull. 171, 112744. <u>https://doi.org/10.1016/j.marpolbul.2021.112744</u>, 2021.
- 1252 Mannarini, G., and Carelli, L.: VISIR-1.b: ocean surface gravity waves and currents for energy-efficient navigation, Geosci.
- 1253 Model Dev., 12, 3449-3480. <u>https://doi.org/10.5194/gmd-12-3449-2019</u>, 2019.

- Mannarini, G., Carelli, L., Orović, J., Martinkus, C.P., and Coppini, G.: Towards Least-CO2 Ferry Routes in the Adriatic Sea. J. Mar. Sci. Eng., 9(2), 115. https://doi.org/10.3390/jmse9020115, 2021.
- Mannarini, G., Pinardi, N., Coppini, G., Oddo, P., and Iafrati, A.: VISIR-I: small vessels least-time nautical routes using wave forecasts. Geosci. Model Dev., 9, 1597-1625. https://doi.org/10.5194/gmd-9-1597-2016, 2016.
- 1258 Marambio, M., Canepa, A., Lòpez, L., Gauci, A.A., Gueroun, S.K.M., Zampardi, S., Boero, F., Yahia, O.K.-D., Yahia,
- 1259 M.N.D., Fuentes, V., et al.: Unfolding Jellyfish Bloom Dynamics along the Mediterranean Basin by Transnational Citizen 1260 Science Initiatives. Diversity, 13(6), 274. https://doi.org/10.3390/d13060274, 2021.
- 1261 Melaku Canu, D., Laurent, C., Morello, E.B., Querin, S., Scarcella, G., Vrgoc, N., Froglia, C., Angelini, S., and Solidoro, C.:
- 1262 Nephrops norvegicus in the Adriatic Sea: Connectivity modeling, essential fish habitats, and management area network.
- 1263 Fisheries Oceanography, 30(4), 349-365. https://doi.org/10.1111/fog.12522, 2020.
- Miladinova, S., Stips, A., Macias-Moy, D., and Garcia-Gorriz, E.: Tracing water-soluble, persistent substances in the Black
  Sea. Environmental Pollution, 308, 119708. https://doi.org/10.1016/j.envpol.2022.119708, 2020.
- 1266 Morell Villalonga, M., Espino Infantes, M., Grifoll Colls, M., and Mestres Ridge, M.: Environmental management system
- 1267 for the analysis of oil spill risk using probabilistic simulations. J. Mar. Sci. Eng., 8(4), 277. 1268 <u>https://doi.org/10.3390/jmse8040277</u>, 2020.
- 1269 Pollani, A., Triantafyllou, G., Petihakis, G., Knostantinos, N., Costas, D., and Koutitas, C.: The Poseidon Operational Tool 1270 for the Prediction of Floating Pollutant Transport. Marine Pollution Bulletin. 43(7-12). 270-278. 1271 https://doi.org/10.1016/S0025-326X(01)00080-7, 2001.
- 1272 Porporato, E.M.D., Pastres, R., and Brigolin, D.: Site suitability for finfish marine aquaculture in the Central Mediterranean
- 1273 Sea. Frontiers in Marine Science, 6, 772. <u>https://doi.org/10.3389/fmars.2019.00772</u>, 2020.
- Schroeder, K., and Chiggiato, J.: Oceanography of the Mediterranean Sea An Introductory Guide. Elsevier, 1st Edition.
   <a href="https://doi.org/10.1016/C2020-0-00371-3">https://doi.org/10.1016/C2020-0-00371-3</a>, 2022.
- Sorgente, R., La Guardia, D., Ribotti, A., Arrigo, M., Signa, A., et al.: An operational supporting system for oil spill
  emergencies addressed to the Italian coast guard. J. Mar. Sci. Eng., 8(12), 1035. <u>https://doi.org/10.3390/jmse8121035</u>, 2020.
- Sotillo, M.G., Cerralbo, P., Lorente, P., Grifoll, M., Espino, M., Sanchez-Arcilla, A., et al.: Coastal ocean forecasting in
  Spanish ports: the Samoa operational service. Journal of Operational Oceanography, 13(1), 37-54.
  https://doi.org/10.1080/1755876X.2019.1606765., 2019.
- 1281 Soto-Navarro, J., Jorda, G., Deudero, S., Alomar, C., Amores, A., and Compa, M.: 3D hotspots of marine litter in the
- 1282 Mediterranean: A modeling study. Marine Pollution Bulletin. 155, 111159. https://doi.org/10.1016/j.marpolbul.2020.111159
- 1283 Stanev, E.V., and Ricker, M. (2019). The Fate of Marine Litter in Semi-Enclosed Seas: A Case Study of the Black Sea.
- 1284 Frontiers in Marine Science, 6. <u>https://doi.org/10.3389/fmars.2019.00660</u>, 2020.
- 1285 Tamvakis, A., Tsirtsis, G., Karydis, M., Patsidis, K. and Kokkoris, G.D.: Drivers of harmful algal blooms in coastal areas of
- 1286 Eastern Mediterranean: a machine learning methodological approach. Mathematical Biosciences and Engineering, 18(5),
- 1287 6484-6505. DOI: 10.3934/mbe.2021322, 2021.
- 135 136

- 1288 Tsagaraki, T.M., Petihakis, G., Tsiaras, K., Triantafyllou, G., Tsapakis, M., Korres, G., Kakagiannis, G., Frangoulis, C. and
- 1289 Karakassis.: Beyond the cage: ecosystem modelling for impact evaluation in aquaculture. Ecological Modelling, 222(14),
  1290 2512-2523. https://doi.org/10.1016/j.ecolmodel.2010.11.027, 2011.
- 1291 Tsiaras, K., Hatzonikolakis, Y., Kalaroni, S., Pollani, A., Triantafyllou, G.: Modeling the pathways and accumulation 1292 patterns of micro-and macro-plastics in the Mediterranean. Frontiers in Marine Science. 8. 1293 https://doi.org/10.3389/fmars.2021.743117, 2021.
- 1294 Tsiaras K., Costa E., Morgana S., Gambardella C., Piazza V., Faimali M., et al.: Microplastics in the Mediterranean: 1295 Variability From Observations and Model Analysis. Frontiers Marine Science, 9. in 1296 https://doi.org/10.3389/fmars.2022.784937, 2022a.
- 1297
- Tsiaras, K., Tsapakis, M., Gkanassos, A., Kalantzi, I. Petihakis, G., and Triantafyllou, G., (2022b). Modelling the impact of
  finfish aquaculture waste on the environmental status in an Eastern Mediterranean Allocated Zone for Aquaculture.
  Continental Shelf Research, 234, 104647. <a href="https://doi.org/10.1016/j.csr.2022.104647">https://doi.org/10.1016/j.csr.2022.104647</a>, 2022b.
- 1301 Umgiesser, G., Ferrarin, C., Bajo, M., Bellafiore, D., Cucco, A., De Pascalis, F., Ghezzo, M., Mc Kiver, W., and Arpaia, L.:
- 1302 Hydrodynamic modelling in marginal and coastal seas The case of the Adriatic Sea as a permanent laboratory for 1303 numerical approach. Ocean Modelling, 179, 102123. <u>https://doi.org/10.1016/j.ocemod.2022.102123</u>, 2022.
- van Sebille, E., Griffies, S.M., Abernathey, R. et al.: Lagrangian ocean analysis: Fundamentals and practices. Ocean
  Modelling, 121, 49-75. <u>https://doi.org/10.1016/j.ocemod.2017.11.008</u>, 2018.
- 1306 Zodiatis, G., Coppini, G., Sepp Neves, A. A., Liubartseva, S., Peña, J., Nikolaidis, A., and Hadjistassou, C.: Coastal oil spill
- 1307 predictions for port's Offensive Security Certified Professional (OSCP) gualification, EGU General Assembly 2024, Vienna,
- 1308 Austria, 14-19 Apr 2024, EGU24-15262, <u>https://doi.org/10.5194/egusphere-egu24-15262</u>, 2024.
- Zodiatis, G., De Dominicis, M., Perivoliotis, L., Radhakrishnan, H., Georgoudiset, E., et al.: The Mediterranean decision
  support system for marine safety dedi cated to oil slicks predictions. Deep Sea Res. II, 133, 4-20.
  https://doi.org/10.1016/j.dsr2.2016.07.014, 2016.
- 1312 Zodiatis, G., Lardner, R., Spanoudaki, K., Sofianos, S., Radhakrishnan, H., Coppini, G., Liubartseva, S., Kampanis, N.,
- 1313 Krokos, G., Hoteit, I., Tintore', J., Eremina, T., Drago, A.: Chapter 5 Operational oil spill modelling assessments. Marine
- 1314 Hydrocarbon Spill Assessments, 145-197. https://doi.org/10.1016/B978-0-12-819354-9.00010-7, 2021

#### 1315 **References Region 5: North East Atlantic**

- 1316 Alvarez Fanjul, E., Ciliberti, S., Bahurel, P.: Implementing Operational Ocean Monitoring and Forecasting Systems. IOC-
- 1317 UNESCO, GOOS-275. https://doi.org/10.48670/ETOOFS, 2022.

- 1318 Bruschi, A., Lisi, I., De Angelis, R., Querin, S., Cossarini, G., Di Biagio, V., et al.: Indexes for the assessment of bacterial
- pollution in bathing waters from point sources: The northern Adriatic Sea CADEAU service. Journal of Environmental
  Management, 293, 112878. https://doi.org/10.1016/j.jenvman.2021.112878, 2021.
- 1321 Cusack, C., Dabrowski, T., Lyons, K., Berry, A., Westbrook, G., Nolan, G., Silke, J.: Harmful algal bloom forecast system
- 1322 for SW Ireland. Part II: Are operational oceanographic models useful in a HAB warning system. Harmful Algae, 53, 86-101.
- 1323 DOI: 10.1016/j.hal.2015.11.013, 2016.
- 1324 Cusack, C., Silke, J., Ruiz-Villarreal, M., Eikrem, W., Dale, T., Moejes, F., Maguire, J., Chamberlain, T., Dabrowski, T.,
- 1325 Gerritsen, H., Hvnes, P., Leadbetter, A., Lvons, K., O'Rourke, E., Smyth, D., Miguez, B. M., Marty, S., McFadden, Y. and 1326 Bloom O'Toole, D.: Harmful Algal Bulletins. AtlantOS D8.6. Available Deliverable at 1327 https://atlantos-h2020.eu/download/deliverables/AtlantOS D8.6.pdf, 2018 (last access: 27/07/2024).
- Díaz, H., and Guedes Soares, C.: Review of the current status, technology and future trends of offshore wind farms. Ocean
  Engineering, 209, 107381. https://doi.org/10.1016/j.oceaneng.2020.107381, 2020.
- 1330 El Serafy GY, Mészáros, L., Fernandez, V., Capet, A., She, J., Garcia Sotillo, M., Melet, A., Legrand, S., Mourre, B.,
- 1331 Campuzano, F., Federico, I., Guarnieri, A., Rubio, A., Dabrowski, T., Umgiesser, G., Staneva, J., Ursella, L., Pairaud, I.,
- 1332 Bruschi, A., Frigstad, H., Baetens, K., Creach, V., Charria, G., Alvarez Fanjul, E. .: EuroGOOS Roadmap for (Operational)
- 1333 Coastal (Downstream) Services. Frontiers in Marine Science, 10. <u>https://doi.org/10.3389/fmars.2023.1177615</u>, 2023.
- European Commission, Directorate-General for Maritime Affairs and Fisheries, Joint Research Centre: The EU blue economy report 2020, Publications Office of the European Union, 2020, <u>https://data.europa.eu/doi/10.2771/363293</u>, 2020.
- 1336 Macias, J.C., Avila Zaragozá, P., Karakassis, I., Sanchez-Jerez, P., Massa, F., Fezzardi, D., Yücel Gier, G., Franičević, V.,
- 1337 Borg, J.A., Chapela Pérez, R.M., Tomassetti, P., Angel, D.L., Marino, G., Nhhala, H., Hamza, H., Carmignac, C., and
- 1338 Fourdain, L.: Allocated zones for aquaculture: a guide for the establishment of coastal zones dedicated to aquaculture in the
- 1339 Mediterranean and the Black Sea. General Fisheries Commission for the Mediterranean. Studies and Reviews, 97. Available
- 1340 at https://openknowledge.fao.org/server/api/core/bitstreams/5c494a74-d1f9-4f9d-9b26-9ff8f3b0f634/content, 2019 (last
- 1341 access: 27/07/2024).
- 1342 Sotillo, M.G., Cerralbo, P., Lorente, P., Grifoll, M., Espino, M., Sanchez-Arcilla, A., et al.: Coastal ocean forecasting in 1343 Spanish ports: the Samoa operational service. J. Operat. Oceanogr. 13, 37e54. 1344 https://doi.org/10.1080/1755876X.2019.1606765, 2019.

#### 1345 **References Region 6: South and Central America**

Cabral, A., Bonetti, C.H.C., Garbossa, L.H.P., Pereira-Filho, J., Besen, K., Fonseca, A.L.: Water masses seasonality and
meteorological patterns drive the biogeochemical processes of a subtropical and urbanized watershed-bay-shelf continuum.
Science of The Total Environment, 749, 141553. <a href="https://doi.org/10.1016/j.scitotenv.2020.141553">https://doi.org/10.1016/j.scitotenv.2020.141553</a>, 2020.

- 1349 Franz, G., Garcia, C. A., Pereira, J., de Freitas Assad, L. P., Rollnic, M., Garbossa, L. H. P., ... & Polejack, A.: Coastal ocean
- observing and modeling systems in Brazil: initiatives and future perspectives. Frontiers in Marine Science, 8, 681619.
  https://doi.org/10.3389/fmars.2021.681619, 2021.
- Franz, G., Leitão, P., Santos, A. D., Juliano, M., and Neves, R.: From regional to local scale modelling on the south-eastern Brazilian shelf: case study of Paranaguá estuarine system. Brazilian Journal of Oceanography, 64(3), 277-294.
- 1354 https://doi.org/10.1590/S1679-875920161195806403, 2016.
- 1355 Garbossa, L.H.P., Franz, G., Neves, R., Lapa, K.R.: Desenvolvimento de modelo operacional hidrodinâmico e de qualidade
- de água da baía da Ilha de Santa Catarina. CONGRESSO BRASILEIRO DE AQUICULTURA E BIOLOGIA AQUÁTICA,
  10th, 2023, Florianópolis, 2023.
- 1358 Garbossa, L.H.P., Lapa, K.R., Franz, G.; Saraiva, A.S., Neves, R.: Resultados preliminares da previsão numérica de
- 1359 fitoplâncton para produção de moluscos bivalves na região da baía da Ilha de Santa Catarina. CONGRESSO BRASILEIRO
- 1360 DE AQUICULTURA E BIOLOGIA AQUÁTICA, 9th, 2021, Online.
- 1361 Lapa, K.R., Saraiva, A.S., Garbossa, L.H.P., Gomes, C.H.A.M., Melo, E.M.C., De Melo, C.M.R., Neves, R.: Aplicação do
- 1362 MOHID BIVALVES para espécie Crassostrea gigas cultivas nas baías da Ilha de Santa Catarina. CONGRESSO
- 1363 BRASILEIRO DE AQUICULTURA E BIOLOGIA AQUÁTICA, 9th, 2021, Online.
- 1364 Ribeiro, R.B., Leitão, J.C., Leitão, P.C., Puia, H.L. and Sampaio, A.F.P.: Integration of high-resolution metocean forecast
- 1365 and observing systems at Port of Santos. In "Proceedings of the IX PIANC-COPEDEC, Conference on Coastal and Port
- 1366 Engineering in Developing Countries", Rio de Janeiro, Brasil, 2016.

#### 1367 References Region 8: Arctic

- 1368 Day, J. J., Keeley, S., Arduini, G., Magnusson, L., Mogensen, K., Rodwell, M., Sandu, I., and Tietsche, S.: Benefits and
- challenges of dynamic sea ice for weather forecasts- Weather Clim. Dynam., 3, 713-731. <u>https://doi.org/10.5194/wcd-3-713-</u>
  2022, 2022.
- 1371 Dethloff, K., Handorf, D., Jaiser, R., and Rinke, A.: Kältere Winter durch abnehmendes arktisches Meereis. Physik in 1372 Unserer Zeit, 50(60), 290-297. https://doi.org/10.1002/piuz.201901547, 2019.
- 1373 Eijgelaar, E., Thaper, C., and Peeters, P.: Antarctic cruise tourism: The paradoxes of ambassadorship, "last chance tourism"
- and greenhouse gas emissions. Journal of Sustainable Tourism 18(3), 337-354. <u>https://doi.org/10.1080/09669581003653534</u>,
  2010.
- Fauchald, P., Arneberg, P., Debernard, J. B., Lind, S., Olsen, E. and Hausner, V. H.: Poleward shifts in marine fisheries under Arctic warming. Environmental Research Letters, 16(7), 074057. DOI:10.1088/1748-9326/ac1010, 2021.
- 1378 Fransner, F., Olsen, A., Årthun, M., Counillon, F., Tjiputra, J., Samuelsen, A., Keenlyside, N. 2023: Phytoplankton
- 1379 abundance in the Barents Sea is predictable up to five years in advance. Commun Earth Environ.
- 1380 https://doi.org/10.1038/s43247-023-00791-9

- Grigoryev, T., Verezemskaya, P., Krinitskiy, M., Anikin, N., Gavrikov, A., Trofimov, I., Balabin, N., Shpilman, A.,
  Eremchenko, A., Gulev, S., Burnaev, E., Vanovskiy, V.: Data-Driven Short-Term Daily Operational Sea Ice Regional
  Forecasting. Remote Sensing, 14(22), 5837. https://doi.org/10.3390/rs14225837, 2022.
- He, Y., Shu, Q., Wang, Q. *et al.* Arctic Amplification of marine heatwaves under global warming. *Nat Commun* 15, 8265
  (2024). <u>https://doi.org/10.1038/s41467-024-52760-1</u>
- Inoue, R., Lien, R.-C., Moum, J.N., Perez, R.C., and Gregg, M.C.: Variations of Equatorial Shear, Stratification, and
  Turbulence Within a Tropical Instability Wave Cycle. Journal of Geophysical Research: Oceans, 124, 1858-1875.
  <u>https://doi.org/10.1029/2018JC014480</u>, 2019.
- Kjesbu, O. S., Alix, M., Sandø, A. B., Strand, E., Wright, P. J., Johns, D. G., Thorsen, A., Marshall, C. T., Bakkeplass,
  K. G., Vikebø, F. B., Skuggedal Myksvoll, M., Ottersen, G., Allan, B. J. M., Fossheim, M., Stiansen, J. E., Huse, G., &
  Sundby, S. (2023). Latitudinally distinct stocks of Atlantic cod face fundamentally different biophysical challenges under
  on-going climate change. *Fish and Fisheries*, 24, 297–320. https://doi.org/10.1111/faf.12728
- 1393 Kristensen, N.M., Tedesco, P., Rabault, J., Aarnes, O.J., Saetra, Ø., Breivik, Ø.: NORA-Surge: A storm surge hindcast for 1394 the Norwegian Sea, the North Sea and the Barents Sea. Ocean Modelling, in press. 1395 https://doi.org/10.1016/j.ocemod.2024.102406, 2024.
- 1396 Larsen, J. N., and Fondahl, G.: Arctic human development report: regional processes and global linkages. Copenhagen:
- 1397 Nordic Council of Ministers. Available at <u>http://norden.diva-portal.org/smash/get/diva2:788965/FULLTEXT03.pdf</u>, 2014
   1398 (last access: 27/07/2024).
- 1399 Marchenko, N., Borch, O. J., Markov, S. V., & Andreassen, N.: Maritime activity in the high north The range of unwanted
- 1400 incidents and risk patterns. Proceedings International Conference on Port and Ocean Engineering under Arctic Conditions
- 1401 2015. URI: http://hdl.handle.net/11250/2392588, 2015.
- Neis, B., Finnis, J., Pelot, R., Shewmake, J.: Insights from the History of Fishing Safety: Preparing for Increased Fisheries
  and Shipping in the Canadian Arctic. In: Chircop, A., Goerlandt, F., Aporta, C., Pelot, R. (eds) Governance of Arctic
  Shipping. Springer Polar Sciences. Springer, Cham. <a href="https://doi.org/10.1007/978-3-030-44975-9">https://doi.org/10.1007/978-3-030-44975-9</a> 11, 2020.
- 1405 Nordam, T., Beegle-Krause, CJ., Skancke, J., Nepstad, R., and Reed, M.: Improving oil spill trajectory modelling in the 1406 Arctic. Marine Pollution Bulletin, 140, 65-74. https://doi.org/10.1016/j.marpolbul.2019.01.019, 2019.
- 1407 Posey, P. G., Metzger, E. J., Wallcraft, A. J., Hebert, D. A., Allard, R. A., Smedstad, O. M., Phelps, M. W., Fetterer, F.,
- Stewart, J. S., Meier, W. N., and Helfrich, S. R.: Improving Arctic sea ice edge forecasts by assimilating high horizontal
  resolution sea ice concentration data into the US Navy's ice forecast systems- The Cryosphere, 9, 1735-1745.
  https://doi.org/10.5194/tc-9-1735-2015, 2015.
- 1411 Sutherland, G., de Aguiar, V., Hole, L.-R., Rabault, J., Dabboor, M., and Breivik, Ø.: Estimating a mean transport velocity in
- the marginal ice zone using ice–ocean prediction systems. The Cryosphere, 16, 2103-2114. <u>https://doi.org/10.5194/tc-16-</u>
  2103-2022, 2022.

Tanguy, R., Bartsch, A., Nitze, I., Irrgang, A., Petzold, P., Widhalm, B., et al. (2024). Pan-Arctic assessment of coastal
settlements and infrastructure vulnerable to coastal erosion, sea-level rise, and permafrost thaw. *Earth's Future*, 12,
e2024EF005013. <u>https://doi.org/10.1029/2024EF005013</u>

1417 Wagner, P. M., Hughes, N., Bourbonnais, P., Stroeve, J., Rabenstein, L., Bhatt, U., ... Fleming, A. (2020). Sea-ice
1418 information and forecast needs for industry maritime stakeholders. *Polar Geography*, 43(2–3), 160–187.
1419 https://doi.org/10.1080/1088937X.2020.1766592

#### 1420 **Competing interests**

1421 The contact author has declared that none of the authors has any competing interests.

#### 1422 Data and/or code availability

1423 N/A

#### 1424 Authors contribution

1425 All authors contributed to the content and writing of this manuscript. JV led and collated the various contributions.

#### 1426 Acknowledgements

1427 The authors would like to gratefully acknowledge the contributions from the Ocean to Climate Seamless Forecasting Decade

1428 Programme (OSF).