# 1 Numerical Models for Monitoring and Forecasting Sea Ice: a short

## 2 description of present status

## 3 Laurent Bertino<sup>1</sup>, Patrick Heimbach<sup>2</sup>, Ed Blockley<sup>3</sup>, Einar Ólason<sup>1</sup>

<sup>4</sup> <sup>1</sup>Nansen Environmental and Remote Sensing Center, Bergen, Norway

5 <sup>2</sup>Oden Institute for Computational Engineering and Sciences, The University of Texas at Austin, Austin, TX, United States

<sup>3</sup>Met Office Hadley Centre, Exeter, United Kingdom

8 *Correspondence to*: Laurent Bertino (laurent.bertino@nersc.no)

10 Abstract. The severe changes in climate resulting in the polar oceans getting warmer – with drastic consequences to their 11 physical, biogeochemical and biological state - require forecasting systems that can accurately simulate and skilfully predict 12 the state of the ice cover and its temporal evolution. Sea-ice processes significantly impact ocean circulation, water mass 13 formation and modifications, and air-sea fluxes. They comprise vertical processes, mainly related to thermodynamics, and 14 horizontal ones, due to internal sea ice mechanics and motion. We provide an overview on how these processes can be modelled 15 and how operational systems are working, in combination with data assimilation techniques, to enhance accuracy and 16 reliability. We also emphasize the need for advancing research on improving such numerical techniques by highlighting 17 currents limits and ways forward.

## 18 **1 Introduction**

9

The main objective of an operational sea-ice forecasting system is to provide users with a reliable estimate of the state of the ice cover and its temporal evolution. To meet this goal, the system needs to be coupled to, or use data from, ocean and atmosphere forecasting systems. Some form of data assimilation is also required to provide the model with the best possible starting position, accounting for the chaotic nature of the atmosphere-ocean-ice system. Users of sea-ice forecasting systems can either be ship captains operating in the polar regions or intermediate service providers. With a changing climate and warming polar oceans, the number of stakeholders interested in operating in ice infested waters is growing.

Sea-ice processes have a profound importance for the ocean circulation and water mass modifications, so that ocean models of the polar regions are always coupled to a sea-ice model, both for operational forecasting and climate projection purposes. Sea ice models have their origin in the climate modeling community in the 1970's and were subsequently part of the ocean general circulation model. They have since then evolved to provide sea ice forecasts in their own right and have been made modular to avoid being bound to a given choice of physical ocean model (Blockley et al., 2020). Sea ice observations from 30 satellites are assimilated in the prediction systems (Buehner et al., 2017). This chapter gives a summary of the short-term (up

31 to 10 days) sea ice forecasting systems for the polar regions.

#### 32 **2** Overview of processes in sea ice

The physical processes simulated by sea-ice models are commonly split into two: vertical processes, related to thermodynamic growth and melt, and mechanical and dynamical processes influencing the horizontal movement of ice. This dynamicthermodynamic separation has practical advantages for computations.

#### 36 2.1 Thermodynamics

37 The ocean can freeze in different phases of sea ice, starting with frazil crystals and their conglomerates into a liquid mush 38 referred to as grease ice, then pancake ice in the presence of waves or slush when the waves flood the snow (Wadhams 2000). 39 Slush, grease, pancakes and ice may sound like a perfect birthday party, until you realize that there is also salt in the ice 40 (Feltham et al., 2006, de la Rosa et al., (2011), Jutras et al., (2016)). The latter will be rejected to the ocean through brine 41 channels, but usually after its multi-year birthday party (e.g., Notz and Worster, 2009). Once a layer of ice has formed on the 42 surface of the ocean, new ice is mostly formed from below as crystals moving upward from the ocean mixed layer affix to the 43 base of the ice in a process known as 'congelation growth'. Sea ice also freezes laterally within open leads and between ice 44 floes. Snow accumulates on top of the sea ice and forms an efficient thermal insulator as well as a white coating that reflects 45 solar radiation back to the atmosphere. A smaller amount of snow-ice comes from compacted snow above the ice. The insulating effect of snow inhibits both sea ice growth in early winter and sea ice melt in late winter (Bigdeli et al., 2020). 46

When summertime approaches, the snow melts first, and forms melt ponds at the surface of the ice. These dark ponds absorbmore solar radiation and enhance the summer melt.

The sea ice itself works as an insulating layer between the ocean and the atmosphere, with thick ice a better insulator than thin ice.

## 51 2.2 Mechanics

Sea ice deforms under the action of winds and currents. Their surface drag accumulated over hundreds of kilometers of sea ice results in formidable forces able to crack open the thickest ice or pile it up into pressure ridges, cracks, leads and ridges in what are called linear kinematic features of sea ice. First-year ice (FYI) can become about 1 meter thick while multi-year ice (MYI) is more often deformed via compressive stresses and can easily reach 2 meters or above. The convergence of ice is a major threat to navigation and only a few ice-strengthened vessels or icebreakers are designed to withstand such forces. The deformation of sea ice has been measured by drifting buoys and satellite data and scaling laws have revealed multi-fractal properties (Weiss and Marsan, 2004) and power-law behavior (Weiss et al., 2009). Waves formed in the open ocean will often reach the ice and attenuate within the ice pack, flexing and occasionally breaking the ice into smaller floes along the way. Smaller ice floes offer more reflecting edges and are more efficient at scattering waves. Waves scattering represents a negative feedback in the wave-ice interactions, among other nonlinear energy dissipation processes (Squire, 2020). This equilibrium results in a wave-broken marginal ice zone (MIZ) which is typically 100 km wide in the Arctic but can reach 1000 km in the Southern Ocean where waves are bigger and the ice thinner. Sea ice can also be submerged by waves, making the surface more saline. Wave breaking effects enhance the lateral melting of ice during summer, but also enhance its freezing during winter.

#### 66 2.3 Biogeochemistry

- There is life in sea ice, not only the occasional seal innocently sunbathing as the polar bear lurks around, but as a dense activity under the sea ice following the growth of red ice algae (Duarte et al., 2017). The availability of light below the ice and the size of brine channels determines the growth of algae and the peculiar ecosystem that depends on them (Arrigo, 2014). The algae will find nutrients in the sea ice, some will be trapped in the ice during freezing, providing a sheltered food store for microorganisms and then later ejected to the ocean through brine channels (Lund-Hansen et al. 2024).
- Sea ice carries sediments while drifting from the shallow shelf seas to the central Arctic, together with nutrients, various
   biological materials and occasionally pollutants (Krumpen et al. 2019).
- Sea ice acts as a lid preventing the exchange of greenhouse gases between ocean and atmosphere, but the sea ice also holds its own carbon pump accounting for 30% of the Carbon uptake in the Arctic (Richaud et al. 2023).

#### 76 **3 Numerical models**

77 Operational sea-ice models are based on complex community codes, simulating the dynamical properties (the constitutive law 78 or rheology) and the thermodynamics of sea ice. The most widespread rheological model of sea ice is the Viscous-Plastic 79 model, often met in the Elastic-Viscous-Plastic (EVP) form which is more efficient for massively parallel computing. One or 80 the other is implemented in the Community sea Ice CodE (CICE), the Sea Ice modelling Integrated Initiative (SI<sup>3</sup>), the Louvain-81 la-Neuve sea Ice Model (LIM), the MIT general circulation model (MITgcm), and GFDL's Sea Ice Simulator (SIS2). The 82 previous models all use an Eulerian model grid, but a recent code, the neXt generation Sea Ice Model (neXtSIM) has adopted an adaptive Lagrangian mesh, as well as a more recent Brittle-Bingham Maxwell rheology (Ólason et al., 2022) that exhibits 83 84 linear features of sea ice deformations apparent in Figure 1. All recent sea-ice models are multi-category models and thus 85 explicitly simulate an ice thickness distribution. They also include a sea-ice age tracer and can thus predict areas of FYI and 86 MYI. Their use in operational forecasts is indicated in Table 1.

The above ocean and sea-ice models are coupled via advanced software (OASIS, ESMF, CCSM) that make them modular, but some ocean models come with an integrated sea ice model, for example the NEMO, the MITgcm, the MOM, the HBM and the FESOM2 codes. The latter is using finite volume (Danilov et al., 2017).



Figure 1: Example of sea ice thickness analysis from the neXtSIM-F (left) system and the assimilated CS2SMOS data, visualisation from the Copernicus Marine Services (<u>http://marine.copernicus.eu</u>).

## 93 4 Data assimilation

90

The most important step to initialize a forecast is to assimilate the latest available observations into a numerical model. Some of the most important observations are available in near-real time with sea-ice concentration, thickness, and motions, but feeding them into the model is a delicate matter (Bertino and Holland, 2017; Buehner et al., 2017). Unobserved variables as well as the ocean properties below the ice must be estimated by multivariate update because of the complex processes both within the sea ice and between the ice and ocean. The irregular observational sampling also requires a flow-dependent spatial interpolation. Operational centers run numerical models and data assimilation codes on dedicated High-Performance Computers (HPC).

- 01 The data assimilation methods in operation are most often the 3D-variational (3DVAR) method (Tonani et al., 2015; Waters 02 et al., 2015; Mogensen et al., 2012; Hebert et al., 2015; Smith et al., 2016; Usui et al., 2006), assimilating sea-ice concentration 03 and more recently sea-ice thickness (Mignac et al. 2022). The 4DVAR method is not presently used in operational forecasts 04 but can provide long-term optimized model trajectories that are fully consistent with the model equations (Nguyen et al., 2021). 05 The Ensemble Kalman Filter (EnKF) is also used in the TOPAZ system to assimilate concentrations, thickness, and motion 06 vectors (Xie et al., 2017) and has been tested with neXtSIM (Cheng et al., 2023) although a cheaper nudging is used 07 operationally (Williams et al., 2021). The EnKF does not intrude in the model software and the resulting forecast system is 08 modular. Even though operational centers use the state of the art with respect to sea-ice data assimilation, they are still 09 inaccurate in locating the ice edge (about 40 km at analysis time, Carrières et al., 2017), even less accurate in locating the 10 boundary between FYI and MYI (200 km errors rather than 40 km).
- 11 Biases in sea ice area coverage arise from multiple sources, primarily from biased ocean and atmospheric boundary conditions,
- 12 but also intrinsic biases of the sea ice model itself. These biases interact with each other in complex ways (feedback loops or

- 13 cancellation of errors). Data assimilation methods rely on unbiasedness assumptions and do not remove biases entirely, often
- 14 transferring them to unobserved variables. Short of a complete observing network, there are ongoing efforts in improving sea
- 15 ice models that we believe can reduce biases, provided that incoming biases from new ocean and atmospheric models are also
- 16 reducing.
- 17

18Table 1: List of present-day short-term Global and Arctic forecast systems including specification of spatial resolution, sea ice model,19assimilation method, variables and website. Sea ice variables are SIC concentration, SIT thickness, SIUV motions, SIALB albedo,20SNOW snow depth, SIAGE ice age. Note that the output spatial resolution may differ from the native resolution. Baltic forecasting21systems are omitted for brevity. Ocean data assimilated are also omitted. \* Output interpolated to 9 km. \*\* VENUS is deployed on22demand.

		Syste	Resolutio		Assimilation		
	Coun	m	n at NP	Sea ice	(method and	Variables	
Area	try	name	(km)	Model	sea ice data)	distributed	Website
Arctic	P.R. China	ArcIO PS	18 km	MITgcm	LESTKF SIC, SIT	SIC, SIUV, SIT	http://www.oceanguide .org.cn/IceIndexHome/ ThicknessIce
Glob al	USA	RTOF S	3.5 km	CICE5	3DVAR SIC	SIC, SIT, SIUV	https://polar.ncep.noaa .gov/global/
Arctic	Norw ay	TOPA Z5	6.25 km	CICE5	EnKF SIC, SIUV, SIT	SIC, SIT, SIUV. SNOW, SIALB, SIAGE	https://marine.copernic us.eu/
Arctic	Norw ay	neXtS IM-F	3km (output)	neXtSI M	Nudging SIC	SIC, SIT, SIUV, SNOW, SIALB, SIAGE	https://marine.copernic us.eu/
Glob al	Franc e	МОі	3.5 km	LIM2	SEEK SIC	SIC, SIT, SIUV	https://marine.copernic us.eu/
Glob al	Cana da	GIOPS	12 km	CICE4	3DVAR SIC		<u>CONCEPTS</u> - <u>Science.gc.ca</u>
Arctic	Cana da	RIOPS	3.5 km	CICE4	3DVAR SIC		https://science.gc.ca/ei c/site/063.nsf/eng/h 97 620.html
Glob al	USA	GOFS 3.1	3.5 km	CICE4	3DVAR SIC	SIC, SIT, SIUV	https://www7320.nrlssc .navy.mil/GLBhycomci ce1-12

Glob al	Europ e	ECMW F	12 km*	LIM2	3DVAR SIC	SIC, SIT	https://www.ecmwf.int/ en/forecasts/datasets/s et-i
Arctic	Denm ark	DMI	10 km	CICE4	Nudging SIC		http://ocean.dmi.dk/mo dels/hycom.uk.php
Glob al	UK	Met Office couple d DA	12 km	CICE5	3DVAR SIC	SIC, SIT, SIUV	https://marine.copernic us.eu/
Arctic	Japan	VENU S**	2.5km	IcePOM	N/A	SIC, SIT	https://ads.nipr.ac.jp/ve nus.mirai/#/mirai

23

24 With improved observational data coverage, increased computational power, and improved representation of key physical 25 processes, rapid improvements in sea ice modelling and forecasting capabilities are expected in the coming decade. One 26 research thrust concerns modelling the marginal ice zone, most notably wave-ice interactions (e.g. Boutin et al., 2022) and 27 modelling sea ice as individual floes (e.g., Horvat et al., 2022). A second thrust is improvements in the sea-ice rheology used 28 for the pack ice (e.g. Ólason et al., 2022). Improved rheology will improve the ice drift and the location of the boundary 29 between FYI and MYI (e.g. Regan et al., 2023). Finally, machine learning approaches are flourishing, which seek to develop 30 fast, surrogate modelling and forecasting capabilities (e.g., Hoffman et al., 2023; Durand et al., 2024; Gregory et al., 2024). 31 Sea-ice exists at the boundary between the atmosphere and ocean, so sea-ice forecasts depend on accurate atmosphere, ocean, 32 and even wave forecasts. Improving those is, therefore, very important for improving sea-ice forecasts. We see fully coupled 33 atmosphere-ocean-wave-ice models with fully coupled data assimilation as a vital long-term goal for sea-ice forecasting 34 systems.

Even though every improvement of the atmosphere, ice or ocean models is welcome, they require time-consuming rounds of testing in forced and coupled models. In the meantime, post-processing techniques, now aided by machine learning, are a novelty in sea ice forecasting (Parleme et al. 2021, 2023) and reanalysis (Edel et al. 2025).

#### 38 **References**

Arrigo, K. R.: Sea Ice Ecosystems. Annual Review of Marine Science, 6(1), 439-467. <u>https://doi.org/10.1146/annurev-marine-</u>
 <u>010213-135103</u>, 2014.

41 Bertino, L., and Holland, M. M.: Coupled ice-ocean modeling and predictions. In The Sea, the Science of Ocean Prediction,

42 Part 2. Special Issue, Journal of Marine Resources, 75(6), 839-875. <u>https://doi.org/10.1357/002224017823524017</u>, 2017.

- 43 Bigdeli, A., Nguyen, A. T., Pillar, H. R., Ocana, V. and Heimbach, P.: Atmospheric Warming Drives Growth in Arctic Sea
- 44 Ice: A Key Role for Snow. Geophysical Research Letters, 47(20), 5204. <u>https://doi.org/10.1029/2020gl090236</u>, 2020.
- 45 Blockley, E., Vancoppenolle, M., Hunke, E., Bitz, C., Feltham, D., Lemieux, J., Losch, M., Maisonnave, E., Notz, D., Rampal,
- 46 P., Tietsche, S., Tremblay, B., Turner, A., Massonnet, F., Ólason, E., Roberts, A., Aksenov, Y., Fichefet, T., Garric, G., Iovino,
- 47 D., Madec, G., Rousset, C., Salas y Melia, D., and Schroeder, D.: The future of sea ice modeling: Where do we go from here?
- 48 Bull. Amer. Meteor. Soc., 101, E1302-E1309. https://doi.org/10.1175/BAMS-D-20-0073.1, 2020.
- 49 Boutin, G., Williams, T., Horvat, C., & Brodeau, L.: Modelling the Arctic wave-affected marginal ice zone: a comparison with
- 50 ICESat-2 observations. Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences,
- 51 380(2235). https://doi.org/10.1098/rsta.2021.0262, 2022
- 52 Buehner, M., Bertino, L., Caya, A., Heimbach, P., and Smith, G.: Sea ice data assimilation. In Sea Ice Analysis and Forecasting:
- Towards an Increased Reliance on Automated Prediction Systems (pp. 51-108). Cambridge University Press.
   https://doi.org/10.1017/9781108277600.005, 2017.
- 55 Carrieres, T., Casati, B., Caya, A., Posey, P., Metzger, E. J., Melsom, A., Sigmond, M., Kharin, V., and Dupont, F.: System
- 56 evaluation, in: Sea Ice Analysis and Forecasting, edited by: Carrieres T., Buehner M., Lemieux J. F., and Pedersen, L. T.,
- 57 Cambridge University Press. <u>https://doi.org/10.1017/9781108277600</u>, 2017.
- 58 Cheng, S., Chen, Y., Aydoğdu, A., Bertino, L., Carrassi, A., Rampal, P., and Jones, C. K. R. T.: Arctic sea ice data assimilation
- combining an ensemble Kalman filter with a novel Lagrangian sea ice model for the winter 2019–2020. The Cryosphere, 17,
  1735-1754. https://doi.org/10.5194/tc-17-1735-2023, 2023.
- 61 Danilov, S., Sidorenko, D., Wang, Q., and Jung, T.: The Finite-volumE Sea ice-Ocean Model (FESOM2), Geosci. Model
- 62 Dev., 10, 765-789. <u>https://doi.org/10.5194/gmd-10-765-2017</u>, 2017.
- De La Rosa, S., Maus, S., and Kern, S. (2011): Thermodynamic investigation of an evolving grease to pancake ice field. Annals
   of Glaciology, 52(57), 206-214. doi:10.3189/172756411795931787, 2011.
- 65 Duarte, P., Meyer, A., Olsen, L.M., Kauko, H.M.m Assmy, P., Rosel, A., Itkin, P., Hudson, S.R., Granskog, M.A., Gerland,
- 66 S., Sundfjord, A., Steen, H., Hop, H., Cohen, L., Peterson, A.K., Jeffrey, N., Elliott, S.M., Hunke, E.C., Tuner, A.K. .: Sea ice
- 67 thermohaline dynamics and biogeochemistry in the Arctic Ocean: Empirical and model results, Journal of Geophysical
- 68 Research: Biogeosciences, 122, 1632-1654, <u>https://doi.org/10.1002/2016JG003660</u>, 2017.
- Durand, C., Finn, T. S., Farchi, A., Bocquet, M., Boutin, G. and Ólason, E.: Data-driven surrogate modeling of high-resolution
   sea-ice thickness in the Arctic. The Cryosphere, 18(4), 1791–1815. https://doi.org/10.5194/tc-18-1791-2024, 2024.
- 71 Edel, L., Xie, J., Korosov, A., Brajard, J., and Bertino, L.: Reconstruction of Arctic sea ice thickness (1992–2010) based on a
- hybrid machine learning and data assimilation approach, The Cryosphere (in press), https://doi.org/10.5194/egusphere-2024-
- 73 1896, 2025.
- 74 Feltham, D. L., Untersteiner, N., Wettlaufer, J. S., and Worster, M. G.: Sea ice is a mushy layer. Geophysical Research Letters,
- 75 33(14), L14501. <u>https://doi.org/10.1029/2006GL026290</u>, 2006.

- 76 Hebert, D. A., Allard, R. A., Metzger, E. J., Posey, P. G., Preller, R. H., Wallcraft, A. J., Phelps, M. W., and Smedstad, O. M.:
- 77 Short-term sea ice forecasting: An assessment of ice concentration and ice drift forecasts using the U.S. Navy's Arctic Cap
- 78 Nowcast/Forecast System. J. Geophys. Res. Oceans, 120, 8327-8345. <u>https://doi.org/10.1002/2015JC011283</u>, 2015.
- 79 Gregory, W., Bushuk, M., Zhang, Y., Adcroft, A. & Zanna, L.: Machine Learning for Online Sea Ice Bias Correction Within
- 80 Global Ice-Ocean Simulations. Geophysical Research Letters, 51(3). https://doi.org/10.1029/2023gl106776, 2024.
- 81 Hoffman, L., Mazloff, M. R., Gille, S. T., Giglio, D., Bitz, C. M., Heimbach, P. and Matsuyoshi, K., 2023: Machine learning
- 82 for daily forecasts of Arctic sea-ice motion: an attribution assessment of model predictive skill. Artificial Intelligence for the
- 83 Earth Systems, 1–45. <u>https://doi.org/10.1175/aies-d-23-0004.1</u>, 2023.
- 84 Horvat, C. Floes, the marginal ice zone and coupled wave-sea-ice feedbacks. Philosophical Transactions of the Royal Society
- 85 A, 380(2235), 20210252. <u>https://doi.org/10.1098/rsta.2021.0252</u>, 2022.
- 86 Jutras, M., Vancoppenolle, M., Lourenço, A., Vivier, F., Carnat, G., Madec, G., ... Tison, J. L.: Thermodynamics of slush and
- snow-ice formation in the Antarctic sea-ice zone. Deep Sea Research Part II: Topical Studies in Oceanography, 131, 75–83.
  https://doi.org/10.1016/j.dsr2.2016.03.008, 2016.
- 89 Krumpen, T., Belter, H.J., Boetius, A. et al. Arctic warming interrupts the Transpolar Drift and affects long-range transport of
- 90 sea ice and ice-rafted matter. Sci Rep 9, 5459 (2019). https://doi.org/10.1038/s41598-019-41456-y
- Lund-Hansen, L.C., Gradinger, R., Hassett, B. *et al.* Sea ice as habitat for microalgae, bacteria, virus, fungi, meio- and
  macrofauna: A review of an extreme environment. *Polar Biol* (2024). <a href="https://doi.org/10.1007/s00300-024-03296-z">https://doi.org/10.1007/s00300-024-03296-z</a>
- 93 Mignac, D., Martin, M., Fiedler, E., Blockley, E. and Fournier, N., Improving the Met Office's Forecast Ocean Assimilation
- 94 Model (FOAM) with the assimilation of satellite-derived sea-ice thickness data from CryoSat-2 and SMOS in the Arctic. Q J
- 95 R Meteorol Soc, 1–24, https://doi.org/10.1002/qj.4252, 2022
- 96 Mogensen, K., Balmaseda, M. A., and Weaver, A. (2012) ..: The NEMOVAR ocean data assimilation system as implemented
- 97 in the ECMWF ocean analysis for System 4, ECMWF Technical Memorandum, 668, 1-59. <u>https://doi.org/10.21957/x5y9yrtm</u>,
  98 2012.
- 99 Nguyen, A. T., Pillar, H., Ocaña, V., Bigdeli, A., Smith, T. A., & Heimbach, P.: The Arctic Subpolar gyre sTate Estimate:
- 00 Description and assessment of a data-constrained, dynamically consistent ocean-sea ice estimate for 2002–2017. Journal of
- 01 Advances in Modeling Earth Systems, 13, e2020MS002398. <u>https://doi.org/10.1029/2020MS002398</u>, 2021.
- 02 Notz, D. & Worster, M. G.: Desalination processes of sea ice revisited. Journal of Geophysical Research: Oceans (1978–2012),
- 03 114(C5). <u>https://doi.org/10.1029/2008jc004885</u>, 2009.
- 04 Ólason, E., Boutin, G., Korosov, A., Rampal, P., Williams, T., Kimmritz, M., et al.: A new brittle rheology and numerical
- framework for large-scale sea-ice models. Journal of Advances in Modeling Earth Systems, 14, e2021MS002685.
   <a href="https://doi.org/10.1029/2021MS002685">https://doi.org/10.1029/2021MS002685</a>, 2022.
- 07 Palerme, C. and Müller, M.: Calibration of sea ice drift forecasts using random forest algorithms, The Cryosphere, 15, 3989–
- 08 4004, https://doi.org/10.5194/tc-15-3989-2021, 2021.

- 09 Palerme, C., Lavergne, T., Rusin, J., Melsom, A., Brajard, J., Kvanum, A. F., Macdonald Sørensen, A., Bertino, L., and Müller,
- 10 M.: Improving short-term sea ice concentration forecasts using deep learning, The Cryosphere, 18, 2161-2176,
- 11 https://doi.org/10.5194/tc-18-2161-2024, 2024.
- 12 Richaud, B., Fennel, K., Oliver, E. C. J., DeGrandpre, M. D., Bourgeois, T., Hu, X., and Lu, Y.: Underestimation of oceanic
- carbon uptake in the Arctic Ocean: ice melt as predictor of the sea ice carbon pump, The Cryosphere, 17, 2665–2680,
  https://doi.org/10.5194/tc-17-2665-2023, 2023.
- 15 Shih, Y., Mehlmann, C., Losch, M. and Stadler, G.: Robust and efficient primal-dual Newton-Krylov solvers for viscous-
- 16 plastic sea-ice models. Journal of Computational Physics, 474, 111802. <u>https://doi.org/10.1016/j.jcp.2022.111802</u>, 2023.
- 17 Smith, G. C., Roy, F., Reszka, M., Surcel Colan, D., He, Z., Deacu, D., Belanger, J.-M., Skachko, S., Liu, Y., Dupont, F.,
- 18 Lemieux, J.-F., Beaudoin, C., Tranchant, B., Drévillon, M., Garric, G., Testut, C.-E., Lellouche, J.-M., Pellerin, P., Ritchie,
- 19 H., Lu, Y., Davidson, F., Buehner, M., Caya, A. and Lajoie, M. (2016),.: Sea ice forecast verification in the Canadian Global
- 20 Ice Ocean Prediction System. Q.J.R. Meteorol. Soc., 142(695), 659-671. <u>https://doi.org/10.1002/qj.2555</u>, 2016.
- Squire, V.A.: Ocean Wave Interactions with Sea Ice: A Reappraisal. Annual Review of Fluid Mechanics 2020 52:1, 37-60.
   https://doi.org/10.1146/annurev-fluid-010719-060301, 2020.
- 23 Tonani M., Balmaseda, M., Bertino, L., Blockey, E., Brassington, G., Davidson, F., Drillet, Y., Hogan, P., Kurugano, T., Lee,
- 24 T., Mehra, A., Paranathara, F., Tanajura, C.A.S., and Wang, H.: Status and future of global and regional ocean prediction
- 25 systems, Journal of Operational Oceanography, 8(sup2), s201-s220. <u>http://doi.org/10.1080/1755876X.2015.1049892</u>, 2015.
- 26 Usui N, Ishizaki S, Fujii Y, Tsujino H, Yasuda T, Kamachi M. (2006) ..: Meteorological Research Institute multivariate ocean 27 variational estimate (MOVE) Some early results. Adv. system: Space Res., 37, 806-822. 28 https://doi.org/10.1016/j.asr.2005.09.022, 2006.
- 29 Wadhams, P. (2000). Ice in the Ocean (1st ed.). CRC Press. <u>https://doi.org/10.1201/9781482283082</u>
- 30 Waters, J., Lea, D.J., Martin, M.J., Mirouze, I., Weaver, A., and While, J. (2015),.: Implementing a variational data assimilation
- system in an operational 1/4 degree global ocean model. Q.J.R. Meteorol. Soc., 141, 333-349. <u>https://doi.org/10.1002/qj.2388</u>,
  2015.
- 33 Weiss, J., and Marsan, D. Scale properties of sea ice deformation and fracturing. Comptes Rendus Physique, 5, 735-751.
- 34 <u>https://doi.org/10.1016/j.crhy.2004.09.005</u>, 2004.
- 35 Weiss, J., Marsan, D., Rampal, P.: Space and Time Scaling Laws Induced by the Multiscale Fracturing of The Arctic Sea Ice
- 36 Cover. In: Borodich, F. (eds) IUTAM Symposium on Scaling in Solid Mechanics. Iutam Bookseries, vol 10. Springer,
- 37 Dordrecht. https://doi.org/10.1007/978-1-4020-9033-2 10, 2009.
- 38 Williams, T., Korosov, A., Rampal, P., Ólason, E.: Presentation and evaluation of the Arctic sea ice forecasting system
- 39 neXtSIM-F. The Cryosphere, 15, 3207-3227. <u>https://doi.org/10.5194/tc-15-3207-2021</u>, 2021.
- 40 Xie, J., Bertino, L., Counillon, F., Lisæter, K. A., and Sakov, P.: Quality assessment of the TOPAZ4 reanalysis in the Arctic
- 41 over the period 1991–2013. Ocean Science, 13(1), 123-144. <u>http://doi.org/10.5194/os-13-123-2017</u>, 2017.

### 42 Competing interests

43 The contact author has declared that none of the authors has any competing interests.

#### 44 Data and/or code availability

45 Data used in Figure 1 is freely available from https://marine.copernicus.eu

## 46 Authors contribution

47 LB prepared the manuscript with contribution from all co-authors.

## 48 Acknowledgements

- 49 LB acknowledges financial support from the European Union's Horizon Europe project ACCIBERG (grant agreement No.
- 50 101081568). EB was funded by the Met Office Advancing Arctic meteorological and oceanographic Capabilities & services
- 51 project, which is supported by the Department for Science, Innovation & Technology (DSIT).