

Towards Earth System Modeling: Coupled Ocean Forecasting

Ségolène Berthou¹, John Siddorn², Vivian Fraser-Leonhardt¹, Pierre-Yves Le Traon³, Ibrahim Hoteit⁴

¹MetOffice, Exeter, UK

²Data, Science and Technology, National Oceanography Centre, Southampton, UK

5 ³Mercator Ocean International, Toulouse, France

⁴Physical Science and Engineering Division, King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia

Correspondence to: Ségolène Berthou (segolene.berthou@metoffice.gov.uk)

10

Abstract. Forecasting across different earth system components has initially been achieved independently, but increasing computer power, increasing model accuracy, increasing connectivity between experts and increasing need for multi-hazard weather warning is changing the scene. Coupling methods, which involve exchanging information between discrete modelling systems, enable to gain accuracy and consistency across earth system components. The paper explains the principles of two-way coupling, where models run simultaneously and exchange information both ways. As individual models reach better accuracy, coupling becomes a key factor to improve forecasting capability because it reproduces the natural complexity of the environment: a wealth of literature shows the benefits of coupling. However, coupling is still limited in operational oceanography by its large demands on computational resources, by data assimilation techniques (currently not very well harmonized between the different models) and by administrative separation of forecasts across different earth system components. Overcoming these barriers will support ocean predictions towards a multi-hazard approach and a more accurate representation of the earth systems component interactions and improve collaborations between multi-disciplinary forecasting communities.

20

1 Introduction

25 Coupling can be loosely defined as the process of exchanging information between discrete modelling systems, generally of components of the earth system, to better represent exchange processes (Shapiro et al., 2010). The number of components of a coupled system, and indeed the level of coupling between the components, varies depending on the application. Coupled global climate models (GCMs) generally include the ocean, ice, atmosphere and land surface. Increasingly surface waves are included to represent the exchange between the ocean and the atmosphere better, especially for applications that require representation of natural hazards such as storms. For earth system models which need to include predictions of the biogenic components to predict carbon and other nutrient transfers the components are often extended to include ocean biogeochemistry and atmospheric chemistry (Mulcahy et al., 2023).

30

There are a number of solutions to how this coupling may be achieved, and which is preferred will depend both on the scientific importance of the exchanges and the timescales on which they occur and on technical limitations. In the “traditional” way of working the models are run independently with a flux of information from adjacent components of the earth system being calculated based on independent and non-interactive models. This implies that the winds, precipitation and air temperatures (“forcing”) used to drive the exchanges at the ocean’s surface do not respond to changes in the ocean conditions themselves. The forcing is not calculated on a timestep basis but over a period generally somewhere between an hour and a day. Forecasts run in this mode are termed forced or one-way coupled.

Coupled systems exist with varying complexity of exchanges between models. For example, a common approach for the coupling of hydrodynamics and sea ice is to run both systems at the same time and exchange information both ways. These are termed fully or two-way coupled systems. In these two-way coupled systems, the independent models often communicate with each other through an interface code (“coupler”) which allows the independent models to operate on different grids and with different timesteps (Larson et al. 2005, Valcke, 2013, Hanke et al. 2016). As the number of components interacting with each other increases the flexibility of including a coupler becomes increasingly attractive. A coupling software creates a computational interface between separate systems that allows the passing of information between them without undue intrusion into the code of the modelling systems. This approach is widely used (e.g. Lewis et al., 2018; Pianezze et al., 2022; Wahle et al., 2017) but other approaches exist. ECMWF (Wedi et al., 2015) have integrated their various modelling components into a single executable with the passing of information being done internally within the code rather than through a separate coupling software. Figure 1 illustrates the Regional Environmental Prediction system under development in the United Kingdom, with complex exchanges between five different models, using three different coupling approaches (Best et al., 2004, Valcke 2005, Bruggeman and Bolding, 2014).




Figure 1: Regional coupled system under development in the United Kingdom for the Regional Environmental Prediction project (Lewis et al., 2019), bringing together all the models run by the Met Office for short-term predictions and climate projections. Arrows represent exchanges between models – either as integrated coupling at the time-step (Best et al. 2004) (UM/JULES), 2D coupling through the OASIS coupler (Valckle et al. 2013) (UM / WaveWatch III / NEMO) or 3D coupling through the FABM coupler (Bruggeman and Bolding, 2014) (NEMO/ERSEM).

2 Why is coupling important for Ocean Prediction?

Atmosphere / ocean coupling is common practice at the seasonal and decadal timescales. At these scales, most of the memory is contained in the ocean and in coupled interactions, such as for the El Nino Southern Oscillation (ENSO). Indeed, both the ocean and atmosphere can propagate an anomaly in the other component to remote places. For example, oceanic equatorial waves generated by wind anomalies can propagate to the whole tropical Pacific and generate an El Nino event, and in turn the atmosphere may generate teleconnections from the tropics to the mid-latitudes through upper-level Rossby wave trains in the troposphere or planetary waves in the stratosphere, and influence the ocean back in remote ocean basins (Hardiman et al., 2019; Kim et al., 2012). These may take longer than 10-days to propagate and are therefore sources of seasonal and multi-annual forecast signals. For short-term marine prediction, coupling is emerging as a new potential for improving both atmospheric and oceanic predictions (Brassington et al., 2015).

A clear and extremely well documented weather situation when air-sea coupling is key for both atmosphere and ocean are tropical cyclone forecasts: the strength of tropical cyclones is decreased through large decreases in sea surface temperature (SST) caused by intense turbulent fluxes, deepening of the surface mixed layer by entrainment (Vellinga et al., 2020; Mogensen

et al., 2017; Castillo et al., 2022; Feng et al., 2019) and (if the cyclone translation speed is slow) by upwelling (Corale et al.,
75 2023; Yablonsky et al., 2009). In more general situations, coupling reduces the lifetime of mesoscale eddies and dampens
submesoscale currents through dampening of the wind stress curl and heat fluxes (Yang et al., 2019; Renault et al., 2016;
Renault et al., 2018; Dawe and Thompson, 2006). Coupling also sometimes involves a higher resolution atmosphere than
forcing, which then results in more turbulent eddy kinetic energy in the ocean (Storto et al., 2023). In the tropics, dynamical
80 waves in the atmosphere and ocean can influence each other. For example, Madden Julian Oscillation (MJO) atmospheric
events in the Indian Ocean can be modulated by coupling (Fu et al., 2017) or simply by the diurnal cycle of SST (Karlowska
et al., 2023). Convectively Coupled Kelvin waves also generate a strong signal in the Indian ocean (Azaneu et al., 2021).
At the coastal scale, coupling also becomes interesting since the assumptions of equilibrium between earth system components
often break down (e.g. wave state is not in equilibrium with winds in the sheltered North Sea - Grayek et al., 2023; Wiese et
al., 2019; Whale et al., 2017). Some examples in the literature include better near-surface currents and upwelling forecasting
85 with the inclusion of the Stokes-Coriolis drift by a wave model, which induce an extra term of advection in the direction of
wave group speed (Alari et al., 2016; Bruciaferri et al., 2021). Coupling also benefits wave modelling, for example where tidal
currents modulate wave and wind activity (Renault et al., 2022; Valiente et al., 2021). Coupling an ocean with waves can have
considerable impacts on SSTs, which can go in either direction, depending on the difference in momentum stress passed to the
ocean (more momentum input by the waves in the case of Lewis et al. (2019), resulting in a near-surface cooling, but less
90 momentum in Alari et al. (2016), resulting in warming), through modulation of the ocean stratification. Coupling a wave model
with an atmospheric model will tend to decrease wind speed over young seas and increase ocean momentum flux, especially
important during storms (Gentile et al., 2022; Bouin and Lebeaupin Brossier, 2020). In general, coupling will tend to dampen
air-sea fluxes because components will tend to adjust to one another, so this may decrease ocean spread at the start of ensemble
forecasts (Lea et al., 2022). However, the spread in SST will increase rapidly in regions which have a shallow surface mixed
95 layer, which respond quickly to atmospheric spread (Lea et al., 2022). Precipitation and river flows can also have a local
influence on near-surface temperatures and salinity in the ocean, especially during extreme precipitation events (Bouin et al.,
202; Sauvage et al., 2018). The ocean can finally act as memory between two intense atmospheric events (e.g strong winds,
and strong precipitation (Berthou et al., 2016; Lebeaupin Brossier et al., 2012) or in the case of marine heatwaves and extreme
temperature or precipitation event (Berthou et al., 2024; Martín et al., 2024), in which cases a coupled system is beneficial for
100 longer range forecasting. In regional atmospheric forecasts, using a predicted SST (either obtained through coupling or forcing)
is beneficial for variables such as near-surface temperature (Mahmood et al., 2021), fog (Fallmann et al., 2019) or snow
(Yamamoto et al., 2011).

However, it is worth noting that differences in near-surface parameterisations can also generate differences which are as large
or larger than coupling differences (Gentile et al., 2022), indicating the need for continuous research and investment in
105 observation systems of near-surface characteristics. Coupling is most successful when the water, heat and momentum budgets
are closed, which can be challenging when model parameterisations are designed in forced mode. Recent parameterisation
improvements taking into account coupled variables include wave coupling in the NEMO Turbulent Kinetic Energy scheme

(Couvelard et al., 2020), or current feedback taken into account in atmospheric turbulence (Renault et al., 2019), or finally the new Wave-Age dependent Stress Parameterisation (Bouin et al., 2024)). In some situations, increasing the complexity of air-sea exchanges can be beneficial, for example including sea spray effects on moisture and heat fluxes (Yang et al., 2019; Xu et al., 2021; Zhang et al., 2005; Bianco et al., 2011).

Coupling with land and river models are also attractive to provide river-flow forecasts, especially as the coupling interface gets more complex, to include back-water effects into rivers and coastal wetting and drying (Bianco et al., 2011). Finally, coupling with biogeochemistry and sediment transport models can provide interesting feedback on the ocean colour, with a feedback loop between thermal stratification and phytoplankton bloom, through the modulation of depth penetration of the solar heat flux (Skákala et al., 2022). Other feedbacks include chemistry and aerosols, where the atmosphere can then provide deposition fluxes (e.g. iron, nitrogen) to the ocean, and the phytoplankton sends back chemicals which can affect low-level cloud cover (Mulcahy et al., 2023).

The potential benefits of using a coupled framework are also reinforced by the move towards a multi-hazard approach to predictions. Natural hazards from multiple sources may combine or occur concurrently. Large waves, storm surges, high-wind speeds, and extreme precipitation are all hazards that are likely to co-occur, and influence each other through coupled feedback that can compound one another (for example through over-topping). Coupled systems that predict this feedback may enable an improvement in the range and consistency of actionable information to be provided through hazard warnings and guidance.

3 How extended is the use of coupled modelling for Ocean Prediction?

Many centres and research groups have developed monitoring and prediction tools independently for individual Earth components (e.g. atmosphere, ocean, land, waves, etc.). This is natural based on the historical context of their development and limitations on computing capabilities, but it has created an infrastructure within and across institutions that adds complexity to the task of unifying prediction systems. The major prediction centres are making progress towards an integrated approach by unifying software infrastructure for models and data assimilation capabilities as well as providing opportunities to increase interactions among the development teams of each system component. At the global scale, the use of atmosphere-ocean-sea-ice coupled model has increased rapidly in the past few years, usually starting with deterministic and then ensemble coupled capability: Allard et al., 2012, Komaromi et al., 2021 (Naval Research Laboratory), Mogensen et al. 2017 (European Centre of Medium Range Forecasting), Smith et al., 2018, Peterson et al., 2022 (Environment and Climate Change Canada), Guiavarc'h et al., 2019 (Met Office). In parallel, the perspective of seamless predictive capability (Ruti et al, 2020), especially important during impactful extreme cyclonic or convective events, means km-scale regional coupled systems are either operational (Durnford et al (2018) for the Great Lakes and Saint Lawrence river, Komaromi et al. 2021 for tropical cyclone regions) or are actively being developed in several centres or research group. Examples include western Europe (Sauvage et al., 2021), southwestern Indian ocean (Corale et al., 2023), the Northwest European shelf (Lewis et al., 2019), the northern Indian ocean (Castillo et al., 2022), the Red Sea (Rui et al., 2019, 2024). Finally, coupled river-ocean models, including two-

140 way coupling between river and ocean are used for operational forecasting of compound flooding during hurricanes in the Gulf of Mexico (Bao et al., 2024, using the COWAST system (Warner et al, 2010)).

The extent to the uptake of coupled modelling is still limited, however, by several barriers. First, it places extreme demands on computational resources: the cost of running an extra model is often prohibitive for agencies with limited forecasting remits (e.g. only ocean forecasting). However, recognising the benefits acknowledged above, these agencies are exploring

145 alternatives, such as coupling with a single-column mixed layer model, either in the atmosphere or the ocean (Voldoire et al., 2017; Lemarié et al., 2021). For the agencies with several remits (e.g. weather, marine, hydrology, air quality forecasting), coupled modelling is more attractive and has the potential to reduce the complexity of the modelling chains, as well as prevent large data transfers between platforms.

A second major barrier is data assimilation, which requires the processing of environmental observations. It is itself a

150 technically challenging problem which is made harder if you try and harmonise that across all the earth system components. Data assimilation requires the calculation of an innovation (difference between the modelled and observed value) and then appropriately adjusting the model parameter space to create a state estimate that is optimised to best reflect understanding of model and observation errors. In coupled systems there are correlations between parameters in the different systems that need to be respected: for example, sea surface and air surface temperature are closely correlated. This creates an additional scientific

155 and technical challenge that needs to be addressed in coupled forecasting systems (Penny and Hamill, 2017). The differing time scales inherent in ocean forecasting and atmospheric NWP is also problematic, though Lea et al. (2015) suggest using the shorter NWP based windows does allow for the retention of the longer oceanic time scales, as long as the memory inherited with cycling the system in time remains intact. Nevertheless, strongly coupled data assimilation means an observation in one model can be beneficial for both models (Fu et al. (2021), Phillipson et al, (2021)) and allows for coupled observation operators.

160 Indeed, remote sensed observations of the ocean (remote sensed SST, radiances, colour, ice freeboard) require filtering out an atmospheric signal, a task which could be dealt with by a coupled assimilation system instead of externally, which potentially introduces contradictory biases from other systems.

Weaker barriers include the need for different frequency of running forecasts: ocean forecasts often run daily with a single deterministic member but the atmospheric and the wave forecasts require sub-daily ensembles with several members. In

165 ensemble modelling, inflated spread schemes are often employed (e.g. in the SST), to generate a much larger spread than the ocean uncertainty, and must be modified in coupled systems (Lea et al., 2022). Nevertheless, the ocean and sea ice uncertainty needs thorough quantification against independent observational datasets for these schemes to be effective. Finally, simple bureaucratic barriers such as the constraint of a common forcing model in international projects can also prevent the adoption of coupled modelling.

Coupling models of different earth system components is a technical task which requires scientific software engineering expertise and high computing resources. Whilst common for seasonal and climate prediction, a handful of operational centres have achieved this for NWP timescales, most of them in the past five years. Coupling enables better treatment of air-sea interactions, especially important in the tropics, for intense events (tropical cyclones), for regions of strong SST gradients, eddies, tidal influence or complex coastlines. The cost is affordable for centres which have the responsibility for forecasting across different earth system components. In these cases, in addition to the benefits of coupled feedback, coupled forecasting allows forecast consistency, essential for impact-based forecasting of multi-hazard events. For other centres, cheaper solutions exist, such as only treating the boundary layer of the other earth system component, which is the most important part for coupling at short timescales.

Coupling models also increases knowledge exchange between researchers in different earth system components, which helps build our understanding of the earth system as a whole. Novel methods such as machine learning and artificial intelligence offer great hope in overcoming some of the barriers faced by traditional NWP. At a time of greater coupling between traditional numerical forecasting system, the use of ML/AI should cut across earth system components, and avoid the pitfalls of parameterisations designed with a single component in mind. This can only be achieved by a strong and organised coupling research community.

References

- Alari, V., Staneva, J., Breivik, Ø. et al.: Surface wave effects on water temperature in the Baltic Sea: simulations with the coupled NEMO-WAM model. *Ocean Dynamics*, 66, 917–930. <https://doi.org/10.1007/s10236-016-0963-x>, 2016.
- Allard, R.A., et al.: Validation test report for the coupled ocean/atmosphere mesoscale prediction system (COAMPS) version 5.0: Ocean/wave component validation. Naval Research Laboratory, p. 99, 2012.
- Azaneu, M., Matthews, A.J., and Baranowski, D.B.: Subsurface Oceanic Structure Associated With Atmospheric Convectively Coupled Equatorial Kelvin Waves in the Eastern Indian Ocean. *Journal of Geophysical Research: Oceans*, 126(7), e2021JC017171. <https://doi.org/10.1029/2021JC017171>, 2021.
- Berthou, S., Mailler, S., Drobinski, P. et al.: Lagged effects of the Mistral wind on heavy precipitation through ocean-atmosphere coupling in the region of Valencia (Spain). *Clim Dyn*, 51, 969-983. <https://doi.org/10.1007/s00382-016-3153-007/s00382-016-3153-0>, 2016.
- Berthou, S., Renshaw, R., Smyth, T. et al.: Exceptional atmospheric conditions in June 2023 generated a northwest European marine heatwave which contributed to breaking land temperature records. *Commun Earth Environ* 5, 287. <https://doi.org/10.1038/s43247-024-01413-8>, 2024.
- Best, M. J., Beljaars, A., Polcher, J., & Viterbo, P. (2004). A Proposed Structure for Coupling Tiled Surfaces with the Planetary Boundary Layer. *Journal of Hydrometeorology*, 5(6), 1271–1278. <https://doi.org/10.1175/JHM-382.1>

- Bianco, L., Bao, J.W., Fairall, C.W. et al.: Impact of Sea-Spray on the Atmospheric Surface Layer. *Boundary-Layer Meteorol* 140, 361-381. <https://doi.org/10.1007/s10546-011-9617-1>, 2011.
- 205 Bouin, M.-N. and Lebeaupin Brossier, C.: Impact of a medicane on the oceanic surface layer from a coupled, kilometre-scale simulation. *Ocean Sci.*, 16, 1125-1142. <https://doi.org/10.5194/os-16-1125-2020>, 2020.
- Bouin, M.-N. and Lebeaupin Brossier, C.: Surface processes in the 7 November 2014 medicane from air–sea coupled high-resolution numerical modelling, *Atmos. Chem. Phys.*, 20, 6861-6881. <https://doi.org/10.5194/acp-20-6861-2020>, 2020.
- Bouin, M.-N., Lebeaupin Brossier, C., Malardel, S., Voldoire, A., and Sauvage, C.: The wave-age-dependent stress parameterisation (WASP) for momentum and heat turbulent fluxes at sea in SURFEX v8.1. *Geosci. Model Dev.*, 17, 117-141. <https://doi.org/10.5194/gmd-17-117-2024>, 2024.
- 210 Brassington, G. B., Martin, M. J., Tolman, H. L., Akella, S., Balmeseda, M., Chambers, C. R. S., ... Todling, R.: Progress and challenges in short- to medium-range coupled prediction. *Journal of Operational Oceanography*, 8(sup2), s239-s258. <https://doi.org/10.1080/1755876X.2015.1049875>, 2015.
- Bruciaferri, D., Tonani, M., Lewis, H. W., Siddorn, J. R., Saulter, A., Castillo Sanchez, J. M., et al.: The impact of ocean-wave coupling on the upper ocean circulation during storm events. *Journal of Geophysical Research: Oceans*, 126, e2021JC017343. <https://doi.org/10.1029/2021JC017343>, 2021.
- 215 Bruggeman, J., & Bolding, K. (2014). A general framework for aquatic biogeochemical models. *Environmental Modelling & Software*, 61, 249–265. <https://doi.org/10.1016/j.envsoft.2014.04.002>
- Castillo, J. M., Lewis, H. W., Mishra, A., Mitra, A., Polton, J., Brereton, A., Saulter, A., Arnold, A., Berthou, S., Clark, D., Crook, J., Das, A., Edwards, J., Feng, X., Gupta, A., Joseph, S., Klingaman, N., Momin, I., Pequignet, C., Sanchez, C., Saxby, J., and Valdivieso da Costa, M.: The Regional Coupled Suite (RCS-IND1): application of a flexible regional coupled modelling framework to the Indian region at kilometre scale. *Geosci. Model Dev.*, 15, 4193-4223. <https://doi.org/10.5194/gmd-15-4193-2022>, 2022.
- 220 Corale, L., Malardel, S., Bielli, S., and Bouin, M.-N.: Evaluation of a Mesoscale Coupled Ocean-Atmosphere Configuration for Tropical Cyclone Forecasting in the South West Indian Ocean Basin. *Earth and Space Science*, 10(3), e2022EA002584. <https://doi.org/10.1029/2022EA002584>, 2023.
- Couvelard, X., Lemarié, F., Samson, G., Redelsperger, J.-L., Ardhuin, F., Benshila, R., and Madec, G.: Development of a two-way-coupled ocean–wave model: assessment on a global NEMO(v3.6)–WW3(v6.02) coupled configuration. *Geosci. Model Dev.*, 13, 3067-3090. <https://doi.org/10.5194/gmd-13-3067-2020>, 2020.
- 230 Dawe, J. T., and Thompson, L.: Effect of ocean surface currents on wind stress, heat flux, and wind power input to the ocean. *Geophysical Research Letters*, 33(9). <https://doi.org/10.1029/2006GL025784>, 2006.
- Fallmann, J., Lewis, H., Sanchez, J. C., and Lock, A.: Impact of high-resolution ocean–atmosphere coupling on fog formation over the North Sea. *Quarterly Journal of the Royal Meteorological Society*, 145(720), 1180-1201. <https://doi.org/10.1002/qj.3488>, 2019.

- 235 Durnford, D., Fortin, V., Smith, G. C., Archambault, B., Deacu, D., Dupont, F., Dyck, S., Martinez, Y., Klyszejko, E., MacKay, M., Liu, L., Pellerin, P., Pietroniro, A., Roy, F., Vu, V., Winter, B., Yu, W., Spence, C., Bruxer, J., & Dickhout, J. (2018). Toward an Operational Water Cycle Prediction System for the Great Lakes and St. Lawrence River. *Bulletin of the American Meteorological Society*, 99(3), 521–546. <https://doi.org/10.1175/BAMS-D-16-0155.1>
- Feng, X., Klingaman, N. P., and Hodges, K. I.: The effect of atmosphere–ocean coupling on the prediction of 2016 western
240 North Pacific tropical cyclones. *Quarterly Journal of the Royal Meteorological Society*, 145(723), 2425–2444. <https://doi.org/10.1002/qj.3571>, 2019.
- Fu, J.-X., Wang, W., Shinoda, T., Ren, H.-L., and Jia, X.: Toward Understanding the Diverse Impacts of Air-Sea Interactions on MJO Simulations. *Journal of Geophysical Research: Oceans*, 122(11), 8855–8875. <https://doi.org/10.1002/2017JC013187>, 2017.
- 245 Fu, D., Small, J., Kurian, J., Liu, Y., Kauffman, B., Gopal, A., Ramachandran, S., Shang, Z., Chang, P., Danabasoglu, G., Thayer-Calder, K., Vertenstein, M., Ma, X., Yao, H., Li, M., Xu, Z., Lin, X., Zhang, S., & Wu, L. (2021). Introducing the New Regional Community Earth System Model, R-CESM. *Bulletin of the American Meteorological Society*, 102(9), E1821–E1843. <https://doi.org/10.1175/BAMS-D-20-0024.1>
- Gentile, E. S., Gray, S. L., and Lewis, H. W.: The sensitivity of probabilistic convective-scale forecasts of an extratropical
250 cyclone to atmosphere–ocean–wave coupling. *Quarterly Journal of the Royal Meteorological Society*, 148(743), 685–710. <https://doi.org/10.1002/qj.4225>, 2022.
- Grayek, S., Wiese, A., Ho-Hagemann, H. T. M., and Staneva, J.: Added value of including waves into a coupled atmosphere–ocean model system within the North Sea area. *Frontiers in Marine Science*, 10. <https://doi.org/10.3389/fmars.2023.1104027>, 2023.
- 255 Guiavarc'h, C., Roberts-Jones, J., Harris, C., Lea, D. J., Ryan, A., and Ascione, I.: Assessment of ocean analysis and forecast from an atmosphere–ocean coupled data assimilation operational system, *Ocean Sci.*, 15, 1307–1326. <https://doi.org/10.5194/os-15-1307-2019>, 2019.
- Hanke, M., Redler, R., Holfeld, T., & Yastremsky, M. (2016). YAC 1.2.0: new aspects for coupling software in Earth system modelling. *Geoscientific Model Development*, 9(8), 2755–2769. <https://doi.org/10.5194/gmd-9-2755-2016>
- 260 Hardiman, S. C., Dunstone, N.J., Scaife, A.A., Smith, D.M., Ineson, S., Lim, J., and Fereday, D.: The Impact of Strong El Niño and La Niña Events on the North Atlantic. *Geophysical Research Letters*, 46(5), 2874–2883. <https://doi.org/10.1029/2018GL081776>, 2019.
- kákala, J., Bruggeman, J., Ford, D., Wakelin, S., Akpınar, A., Hull, T., Kaiser, J., Loveday, B.R., O’Dea, E., Williams, C.A.J., and Ciavatta, S.: The impact of ocean biogeochemistry on physics and its consequences for modelling shelf seas. *Ocean Modelling*, 172, 101976. <https://doi.org/10.1016/j.ocemod.2022.101976>, 2022.
- 265 Karłowska, E., Matthews, A. J., Webber, B. G. M., Graham, T., and Xavier, P.: The effect of diurnal warming of sea-surface temperatures on the propagation speed of the Madden–Julian oscillation. *Quarterly Journal of the Royal Meteorological Society*, 150(758), 334–354. <https://doi.org/10.1002/qj.4599>, 2023.

- Kim, H.M., Webster, P.J. & Curry, J.A.: Seasonal prediction skill of ECMWF System 4 and NCEP CFSv2 retrospective
 270 forecast for the Northern Hemisphere Winter. *Clim Dyn* 39, 2957-2973. <https://doi.org/10.1007/s00382-012-1364-6>, 2012.
- Komaromi, W. A., Reinecke, P. A., Doyle, J. D., and Moskaitis, J. R.: The Naval Research Laboratory's Coupled Ocean–
 Atmosphere Mesoscale Prediction System-Tropical Cyclone Ensemble (COAMPS-TC Ensemble). *Weather and Forecasting*,
 36(2), 499-517. <https://doi.org/10.1175/WAF-D-20-0038.1>, 2021.
- Larson, J., Jacob, R., & Ong, E. (2005). The Model Coupling Toolkit: A New Fortran90 Toolkit for Building Multiphysics
 275 Parallel Coupled Models. *The International Journal of High Performance Computing Applications*, 19(3), 277–292.
<https://doi.org/10.1177/1094342005056115>
- Lea, D. J., While, J., Martin, M. J., Weaver, A., Storto, A., and Chrust, M.: A new global ocean ensemble system at the Met
 Office: Assessing the impact of hybrid data assimilation and inflation settings. *Quarterly Journal of the Royal Meteorological
 Society*, 148(745), 1996-2030. <https://doi.org/10.1002/qj.4292>, 2022.
- 280 Lebeaupin Brossier, C., Drobinski, Béranger, P.K., Bastin, S. and Orain, F.: Ocean memory effect on the dynamics of coastal
 heavy precipitation preceded by a mistral event in the northwestern Mediterranean. *Quarterly Journal of the Royal
 Meteorological Society*, 139(675), 1583-1897. <https://doi.org/10.1002/qj.2049>, 2012.
- Lemarié, F., Samson, G., Redelsperger, J.-L., Giordani, H., Brivoal, T., and Madec, G.: A simplified atmospheric boundary
 layer model for an improved representation of air–sea interactions in eddying oceanic models: implementation and first
 285 evaluation in NEMO (4.0). *Geosci. Model Dev.*, 14, 543-572. <https://doi.org/10.5194/gmd-14-543-2021>, 2021.
- Lewis, H. W., Castillo Sanchez, J. M., Arnold, A., Fallmann, J., Saulter, A., Graham, J., Bush, M., Siddorn, J., Palmer, T.,
 Lock, A., Edwards, J., Bricheno, L., Martínez-de la Torre, A., and Clark, J.: The UKC3 regional coupled environmental
 prediction system. *Geosci. Model Dev.*, 12, 2357-2400. <https://doi.org/10.5194/gmd-12-2357-2019>, 2019.
- Lewis, H. W., Castillo Sanchez, J. M., Siddorn, J., King, R. R., Tonani, M., Saulter, A., Sykes, P., Pequignot, A.-C., Weedon,
 290 G. P., Palmer, T., Staneva, J., and Bricheno, L.: Can wave coupling improve operational regional ocean forecasts for the north-
 west European Shelf? *Ocean Sci.*, 15, 669-690- <https://doi.org/10.5194/os-15-669-2019>, 2019.
- Mahmood, S., Lewis, H., Arnold, A., Castillo, J., Sanchez, C., and Harris, C.: The impact of time-varying sea surface
 temperature on UK regional atmosphere forecasts. *Meteorological Applications*, 28(2), e1983.
<https://doi.org/10.1002/met.1983>, 2021.
- 295 Martín, M. L., Calvo-Sancho, C., Taszarek, M., et al.: Major Role of Marine Heatwave and Anthropogenic Climate Change
 on a Giant Hail Event in Spain. *Geophysical Research Letters*, 51(6), e2023GL107632.
<https://doi.org/10.1029/2023GL107632>, 2024.
- Mogensen, K.S., Magnusson, L., and Bidlot, J.-R.: Tropical cyclone sensitivity to ocean coupling in the ECMWF coupled
 model. *Journal of Geophysical Research: Ocean*, vol. 122, no. 5, pp. 4392-4412. <https://doi.org/10.1002/2017JC012753>, 2017.
- 300 Mulcahy, J. P., Jones, C. G., Rumbold, S. T., Kuhlbrodt, T., Dittus, A. J., Blockley, E. W., Yool, A., Walton, J., Hardacre, C.,
 Andrews, T., Bodas-Salcedo, A., Stringer, M., de Mora, L., Harris, P., Hill, R., Kelley, D., Robertson, E., and Tang, Y.:

- UKESM1.1: development and evaluation of an updated configuration of the UK Earth System Model, *Geosci. Model Dev.*, 16, 1569-1600. <https://doi.org/10.5194/gmd-16-1569-2023>, 2023.
- Penny, S. G., and Hamill, T. M.: Coupled Data Assimilation for Integrated Earth System Analysis and Prediction: Goals, Challenges, and Recommendations. *Bulletin of the American Meteorological Society*, 98(7), ES169–ES172. <https://www.jstor.org/stable/26243775>, 2017.
- Pianezze, J., Beuvier, J., Lebeaupin Brossier, C., Samson, G., Faure, G., and Garric, G.: Development of a forecast-oriented kilometre-resolution ocean–atmosphere coupled system for western Europe and sensitivity study for a severe weather situation. *Nat. Hazards Earth Syst. Sci.*, 22, 1301-1324. <https://doi.org/10.5194/nhess-22-1301-2022>, 2022.
- Renault, L., Lemarié, F., and Arsouze, T.: On the implementation and consequences of the oceanic currents feedback in ocean–atmosphere coupled models. *Ocean Modelling*, 141, 101423. <https://doi.org/10.1016/j.ocemod.2019.101423>, 2019.
- Renault, L., Marchesiello, P.: Ocean tides can drag the atmosphere and cause tidal winds over broad continental shelves. *Communications Earth & Environment*, 3, 70. <https://doi.org/10.1038/s43247-022-00403-y>, 2022.
- Renault, L., McWilliams, J.C., and Gula, J.: Dampening of Submesoscale Currents by Air-Sea Stress Coupling in the Californian Upwelling System. *Sci Rep* 8, 13388. <https://doi.org/10.1038/s41598-018-31602-3>, 2018.
- Renault, L., Molemaker, M.J., McWilliams, J.C., Shchepetkin, A.F., Lamarie’, F., Chelton, D., Illig, S., and Hall, A.: Modulation of Wind Work by Oceanic Current Interaction with the Atmosphere. *Journal of Physical Oceanography*, 46(6), 1685-1704. <https://doi.org/10.1175/JPO-D-15-0232.1>, 2016.
- R. Sun, A. Subramanian, A. Miller, M. Mazloff, I. Hoteit, and B. Cornuelle: SKRIPS v1.0: a regional coupled ocean–atmosphere modeling framework (MITgcm–WRF) using ESMF/NUOPC, description and preliminary results for the Red Sea. *Geoscientific Model Development*, gmd-2018-252, 2019.
- R. Sun, S. Sanikomm, A. Subramanian, M. Mazloff, B. Cornuelle, G. Gopalakrishnan, A. Miller and I. Hoteit: Enhanced regional ocean ensemble data assimilation through atmospheric coupling in the SKRIPS model. *Journal of Ocean Modeling*, 191, 102424, 2024.
- Ruti, P.M., et al.: Advancing Research for Seamless Earth System Prediction. *Bulletin of the American Meteorological Society*, 101(1), E23–E35. <https://doi.org/10.1175/BAMS-D-17-0302.1>, 2020.
- Sauvage, C., Lebeaupin Brossier, C., and Bouin, M.-N.: Towards kilometer-scale ocean–atmosphere–wave coupled forecast: a case study on a Mediterranean heavy precipitation event. *Atmos. Chem. Phys.*, 21, 11857-11887. <https://doi.org/10.5194/acp-21-11857-2021>, 2021.
- Sauvage, C., Lebeaupin Brossier, C., Ducrocq, V., Bouin, M.-N., Vincendon, B., Verdecchia, M., Taupier-Letage, I., Orain F.: Impact of the representation of the freshwater river input in the Western Mediterranean Sea. *Ocean Modelling*, 131, 115-131. <https://doi.org/10.1016/j.ocemod.2018.09.005>, 2018.
- Shapiro, M. A., Shukla, J., Brunet, G., Nobre, C., Belánd, M., Dole, R., ... Wallace, J. M.: An Earth-system prediction initiative for the twenty-first century. *Bulletin Of The American Meteorological Society*, 91, 1377-1388. <https://doi.org/10.1175/2010BAMS2944.1>, 2010.

Smith, G. C., Bélanger, J.-M., Roy, F., Pellerin, P., Ritchie, H., Onu, K., Roch, M., Zadra, A., Colan, D. S., Winter, B., Fontecilla, J.-S., & Deacu, D. (2018). Impact of Coupling with an Ice–Ocean Model on Global Medium-Range NWP Forecast Skill. *Monthly Weather Review*, 146(4), 1157–1180. <https://doi.org/10.1175/MWR-D-17-0157.1>

Storto, A., Hesham Essa, Y., de Toma, V., Anav, A., Sannino, G., Santoleri, R., and Yang, C.: MESMAR v1: a new regional coupled climate model for downscaling, predictability, and data assimilation studies in the Mediterranean region. *Geosci. Model Dev.*, 16, 4811–4833. <https://doi.org/10.5194/gmd-16-4811-2023>, 2023.

Valcke, S.: The OASIS3 coupler: a European climate modelling community software, *Geosci. Model Dev.*, 6, 373–388. <https://doi.org/10.5194/gmd-6-373-2013>, 2013.

Valiente, N. G., Saulter, A., Edwards, J. M., Lewis, H. W.; Castillo Sanchez, J. M. C.; Bruciaferri, D., Bunney, C., Siddorn, J.: The impact of wave model source terms and coupling strategies to rapidly developing waves across the north-west European shelf during extreme events. *Journal of Marine Science and Engineering*, 9 (4), 403. <https://doi.org/10.3390/jmse9040403>, 2021.

Vellinga, M., Copsey, D., Graham, T., Milton, S., and Johns, T.: Evaluating Benefits of Two-Way Ocean–Atmosphere Coupling for Global NWP Forecasts. *Weather and Forecasting*, 35(5), 2127–2144. <https://doi.org/10.1175/WAF-D-20-0035.1>, 2020.

Voldoire, A., Decharme, B., Pianezze, J., Lebeaupin Brossier, C., Sevault, F., Seyfried, L., Garnier, V., Bielli, S., Valcke, S., Alias, A., Accensi, M., Ardhuin, F., Bouin, M.-N., Ducrocq, V., Faroux, S., Giordani, H., Léger, F., Marsaleix, P., Rainaud, R., Redelsperger, J.-L., Richard, E., and Riette, S.: SURFEX v8.0 interface with OASIS3-MCT to couple atmosphere with hydrology, ocean, waves and sea-ice models, from coastal to global scales. *Geosci. Model Dev.*, 10, 4207–4227. <https://doi.org/10.5194/gmd-10-4207-2017>, 2017.

Wahle, K., Staneva, J., Koch, W., Fenoglio-Marc, L., Ho-Hagemann, H. T. M., and Stanev, E. V.: An atmosphere–wave regional coupled model: improving predictions of wave heights and surface winds in the southern North Sea, *Ocean Sci.*, 13, 289–301. <https://doi.org/10.5194/os-13-289-2017>, 2017.

Warner, J. C., Armstrong, B., He, R., & Zambon, J. B. (2010). Development of a Coupled Ocean–Atmosphere–Wave–Sediment Transport (COAWST) Modeling System. *Ocean Modelling*, 35(3), 230–244. <https://doi.org/https://doi.org/10.1016/j.ocemod.2010.07.010>

Wedi, N., et al.: The modelling infrastructure of the Integrated Forecasting System: Recent advances and future challenges. ECMWF Technical Memoranda, no. 760. ECMWF, Apr. 2015. Available at <https://www.ecmwf.int/en/elibrary/78758-modelling-infrastructure-integrated-forecasting-system-recent-advances-and-future>, 2015 (last access: 29/07/2024).

Wiese, A., Stanev, E., Koch, W., Behrens, A., Geyer, B., Staneva, J.: The Impact of the Two-Way Coupling between Wind Wave and Atmospheric Models on the Lower Atmosphere over the North Sea. *Atmosphere*, 10(7), 386. <https://doi.org/10.3390/atmos10070386>, 2019.

Xu, X., Voermans, J. J., Liu, Q., Moon, I.-J., Guan, C., and Babanin, A. V.: Impacts of the Wave-Dependent Sea Spray Parameterizations on Air–Sea–Wave Coupled Modeling under an Idealized Tropical Cyclone, *J Mar Sci Eng*, 9(12).
370 <https://doi.org/10.3390/jmse9121390>, 2021.

Yablonsky, R. M., and Ginis, I.: Limitation of One-Dimensional Ocean Models for Coupled Hurricane–Ocean Model Forecasts. *Monthly Weather Review*, 137(12), 4410–4419. <https://doi.org/10.1175/2009MWR2863.1>, 2009.

Yamamoto, M., Ohigashi, T., Tsuboki, K., and Hirose, N.: Cloud-resolving simulation of heavy snowfalls in Japan for late December 2005: application of ocean data assimilation to a snow disaster. *Nat. Hazards Earth Syst. Sci.*, 11, 2555–2565.
375 <https://doi.org/10.5194/nhess-11-2555-2011>, 2011.

Yang, H., Chang, P., Qiu, B., Zhang, Q., Wu, L., Chen, Z., Wang, H.: Mesoscale Air–Sea Interaction and Its Role in Eddy Energy Dissipation in the Kuroshio Extension. *Journal of Climate*, 32(24), 8659–8676. <https://doi.org/10.1175/JCLI-D-19-0155.1>, 2019.

Zhang, L., Zhang, X., Perrie, W., Guan, G., Dan, B., Sun, C., Wu, X., Liu, K., Li, D.: Impact of Sea Spray and Sea Surface
380 Roughness on the Upper Ocean Response to Super Typhoon Haitang (2005). *Journal of Physical Oceanography*, 51(6), 1929–1945. <https://doi.org/10.1175/JPO-D-20-0208.1>, 2011.

Competing interests

The contact author has declared that none of the authors has any competing interests.

Data and/or code availability

385 Not applicable.

Authors contribution

JS started a draft of this document, SB took over and completed the article, helped with a literature review completed by VFL. PYLT, IH reviewed the text.

Acknowledgements

390 The authors would like to acknowledge the two reviewers for their suggestions of improvement of the manuscript.