Connecting Ocean Observations with Prediction

Pierre-Yves Le Traon¹, Antonio Novellino², Andrew M Moore³

¹Mercator Ocean International, Toulouse, France ²ETT SpA, Genoa, Italy

5 ³Ocean Sciences Department, University of California Santa Cruz, Santa Cruz, California, USA

Correspondence to: Pierre-Yves Le Traon (pletraon@mercator-ocean.fr)

- 10 Abstract. Ocean prediction relies on the integration between models, satellite and in-situ observations through data assimilation techniques. Satellites offer nowadays high-resolution observations of essential ocean variables at the surface, widely adopted in combination with precise but sparse in-situ measurements that, from the surface to the deep ocean, can constrain large scale variability in models. Moreover, observations are a valuable source of information for validating and assessing model products, for improving them and for developing the next generation of machine learning algorithms aimed
- 15 at enhancing the accuracy and scope of ocean forecasts. The authors discuss the role of observations in operational ocean forecasting systems, describing the state-of-the-art of satellite and in-situ observing networks and defining the paths for addressing multi-scale monitoring and forecasting.

1 Introduction: the role of observations for ocean prediction

Ocean prediction relies on the integration between models, satellite and in-situ observations through data assimilation techniques (Bell et al., 2015). Data assimilation provides a 4D dynamical interpolation of observations by considering the complementarities between the different types of observations. High space (e.g. from 10 km at global scale to 1 km or less at regional and coastal scales) and time resolution (e.g. daily) ocean fields consistent with observations and model dynamics are thus derived and can be used to initialize ocean forecasts models. The development of machine learning techniques such as deep neural networks offer different and complementary pathways for ocean prediction. Machine learning techniques analyze and learn from patterns in past data or ocean reanalyses to make ocean predictions from current data. Several studies have already shown the potential of machine learning-based ocean forecast systems (e.g., Chen et al., 2023).

Whatever the techniques used to produce them, the quality of ocean analyses and forecasts observations at global and regional/coastal scales is directly dependent on the availability of high quality in situ and satellite observations with a sufficient

30 space and time resolution. These dependencies vary according to ocean dynamics. Data assimilation is, for example, mandatory and quite effective for constraining the mesoscale variability at global and regional scales. At coastal scales, it is

more challenging to constrain ocean dynamics where small-scale, high frequency and non-linear processes play an important role.

35 Observations are also essential to validate ocean analysis and prediction models (e.g., Gutknecht et al., 2019), to improve ocean models (required both for assessing model performances, for ocean prediction and for digital twins) (e.g., Wang et al., 2023) and for training machine learning algorithms.

For both data assimilation and validation aspects, data must be carefully validated and information on data errors must be 40 documented. Higher quality reprocessed data sets are required for reanalyses.

The monitoring of the impact of observations should be part of any ocean prediction activity. This is done through Observing System Evaluations (OSEs) and Observing System Simulation Experiments (OSSEs) (Fujii et al., 2019; Gasparin et al., 2019). OSEs allow the impact of an existing observing system to be assessed (by withholding observations). OSSEs help in the design

45 of new observing systems, evaluate their different configurations, and perform preparatory data assimilation work. Other complementary approaches for quantifying the impact of observations on ocean analysis and forecast systems also exist (Fujii et al., 2019; Drake et al., 2023).

We briefly review in the following sections the role of the different ocean observing systems for ocean prediction at global, regional and coastal scales. Sections 2 and 3 deal respectively with satellite and in-situ observations.

2 Satellite observations

Satellite observations have a major role and impact on ocean prediction (Le Traon, 2018). Satellites can provide real time and global observations of key ocean variables at high space and time resolution: sea level and geostrophic currents, sea surface temperature, ocean colour, sea ice, surface wave, surface winds (Figure 1). The spatial resolution depends on the nature of the

55 sensor and ranges from a few hundreds of meters (e.g. infrared and ocean colour sensors) to tens of kilometres (e.g. microwave

sensors). The time resolution or revisit time ranges from one hour or less for geostationary satellites up to a few days or longer for polar-orbiting satellites.



Sea level, ocean currents



Sea Ice (concentration, drift, thickness)



Sea Surface Temperature



Ocean Colour, primary production



Waves and Winds



Sea Surface Salinity

Figure 1: The unique contribution of satellite oceanography for ocean prediction

60

Ocean modelling and data assimilation systems have a high dependency on the status of the altimeter constellation (Le Traon et al., 2017). Satellite altimeters provide all-weather observations of sea level which is an integral of the ocean interior and provides a strong constraint on ocean state estimation at the mesoscale. At least four altimeters are required, and a precise knowledge of the Mean Dynamic Topography (MDT) is also a strong requirement for assimilation into operational ocean for a strong constraint on ocean state estimation at the Description of the strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into operational ocean for a strong requirement for assimilation into oper

65 forecasting systems (Le Traon et al., 2017; Hamon et al., 2019).

Sea surface temperature (SST) is a key variable for all ocean prediction systems. SST data can be used to correct for errors in forcing fields (heat fluxes, wind) and to constrain the mesoscale variability of the upper ocean. High resolution SST data from a combination of infra-red (polar orbiting and geostationary) (e.g. S3 SLSTR, VIIRS, GOES, MTG) and microwave sensors

70 (e.g. AMSR-2) are thus essential to constrain ocean prediction systems.

Satellite sea ice concentration and, more recently, sea ice thickness data (SMOS and Cryosat) are routinely assimilated in sea ice models. The assimilation of sea ice drift remains challenging due to the short memory of sea ice drift and sea ice models deficiencies (Sakov et al., 2012). Numerous impact studies have been carried out for sea ice data assimilation, in particular for sea ice thickness that form Crucest but on well for this ice thickness form SMOS and hoth establishes to establish the complete the second sea ice data assimilation.

75 for sea ice thickness products from Cryosat but as well for thin ice thickness from SMOS and both satellites together (Xie et al., 2018).

Sea Surface Salinity observations (SMOS, Aquarius, SMAP) from space (Reul et al., 2020) provide valuable information (Martin et al., 2019; Tranchant et al., 2019) for ocean prediction. Satellite SSS data assimilation can now constrain the model forecasts without introducing incoherent information compared to the other assimilated observations.

80

85

Satellite significant wave height observations are routinely assimilated in global and regional wave models, and their impact is very well demonstrated. Wave spectra provided by Sentinel-1 SAR instruments and, more recently, with the more precise CFOSAT SWIM instrument can, in addition, significantly improve the quality of wave forecasts (Aouf et al., 2021; Hauser et al., 2023).

Ocean color missions (e.g. S3 OLCI, VIIRS) provide essential 'green ocean' observations for a wide range of applications (e.g., water quality, eutrophication, Harmful Algal Blooms). Higher resolution and specialized ocean color products (e.g., case-II water algorithms) are particularly needed for coastal areas. Ocean-color data are being used to assess the performance of model

90 simulations of chlorophyll-a (Chl-a) fields (Gutknecht et al., 2019) and to improve simulations through data assimilation (Ford et al., 2018; Fennel et al., 2019). However, the assimilation of ocean color data is arguably less widespread than that of physical variables. The potential for ocean color data to improve biogeochemical (BGC) models remains significant, though many challenges persist (e.g., error characterization, observation operators such as bio-optical models, and the integration of ocean color data with in-situ measurements like BGC Argo).

95

100

While wind observations from multiple scatterometers are essential for improving the forcing fields required for ocean prediction, the primary pathway for utilizing scatterometer data is through assimilation in numerical weather prediction (NWP) systems. However, NWP data assimilation systems do not incorporate all the information available from scatterometers, particularly at smaller spatial scales (Belmonte Rivas and Stoffelen, 2019). Therefore, using these observations to directly constrain ocean models may be more beneficial.

3 In-situ observations

In-situ observing systems play a fundamental role to provide measurements of the ocean water column and to complement satellite observations. The combination of high-resolution satellite data with sparse and precise in-situ observations of the ocean interior is the only means to provide a high-resolution 3D description and forecast of the ocean state. In situ temperature

105 and salinity data are crucial to constrain large scale variability in models (Gasparin et al., 2023). In situ observations of highfrequency and high-resolution ocean processes in the coastal zone are also essential to validate coastal ocean prediction systems. Ocean prediction uses surface observations, vertical profiles and time series coming from different types of instruments (e.g.,

110 floats, drifters, moorings, marine mammals, gliders, tide gauges, research vessels, ships of opportunity, ferryboxes, saildrones, HF radars) and different parameters (temperature, salinity, currents, sea level, wave, chlorophyll, oxygen, nutrients, pH, fugacity of CO₂) (Figure 2).



115 Figure 2: In-situ networks from the Global Ocean Observing System (GOOS)

Some available observations, such as surface drifters, Thermosalinograph (TSG), Acoustic Doppler Current Profiler (ADCP), are not always assimilated. However, non-assimilated observations are essential for the independent validation of analyses and forecasts, as well as for evaluating model and system improvements.

120

The global Argo array plays a fundamental role for ocean prediction (Le Traon, 2013). Impact studies have confirmed and quantified the major impact of Argo on ocean analysis and forecasting systems (e.g. Turpin et al., 2016). The evolution of Argo into OneArgo, which includes deep and BGC components, already shows very promising results to improve ocean prediction systems (Gasparin et al., 2020; Cossarini et al., 2019; Wang et al., 2021; Mignot et al., 2023).

The most important other source of global observations is the surface drifter network, which provides data on surface currents, sea surface temperature, and, for some drifters, sea surface salinity. Additionally, met-ocean and deep-ocean mooring arrays (temperature, salinity, velocity, and biogeochemical parameters) (OceanSITES, including the TAO/PIRATA/TRITON tropical arrays) provide essential data to validate and constrain models. These are complemented by the Voluntary Observing Ship (VOS) network, which provides SST/SSS data as well as surface carbon measurements.

130

135

There is a growing need to increase in situ data coverage in shelf and coastal areas. Other data sources, such as HF radars, ferryboxes, gliders, tide gauges, and coastal monitoring stations, are regularly used to validate and constrain ocean prediction models. Unmanned surface vehicles (USVs), like saildrones, are also being used with increasing frequency. The assimilation of HF radar data in regional coastal models is an area of active development (Hernandez-Lasheras et al., 2021; Drake et al., 2023), and the assimilation of glider observations with sufficiently dense spatial and temporal sampling at regional and coastal scales has also proven highly effective (Pasmans et al., 2019; Levin et al., 2021; Drake et al., 2023). The development of low-cost technologies and citizen science can also support expanding coverage, particularly in coastal areas.

3 Most important near future challenges

- 140 Ensuring the continuity of existing ocean observing systems is a necessary, but not sufficient, requirement for ocean prediction. Higher spatial and temporal resolution is required to match the increasing model resolution and improve the ability of ocean prediction systems to monitor and forecast smaller scales, including in coastal areas. In this regard, the development of operational swath altimetry (e.g., Morrow et al.2019; Benkiran et al., 2022), following the outstanding results of the SWOT mission (Fu et al., 2024), is one of the most critical requirements for the evolution of the satellite observing system. For in-situ
- 145 observations, critical gaps remain in coastal areas, shelf seas, and polar regions. On a global scale, the lack of biogeochemical observations limits our ability to monitor and forecast the 'green ocean,' making the development of OneArgo a high priority. Data standardization, quality assurance and quality control are also essential to ensure that ocean prediction systems make the best possible use of observations.

References

 Aouf, L., Hauser, D., Chapron, B., Toffoli, A., Tourain, C., Peureux, C.: New Directional Wave Satellite Observations: Towards Improved Wave Forecasts and Climate Description in the Southern Ocean. Geophysical Research Letter, 48(5), e2020GL091187. <u>https://doi.org/10.1029/2020GL091187</u>, 2021.
 Bell, M. J., Le Traon, P.-Y., Smith, N. R., Dombrowsky, E., and Wilmer-Becker, K.: An introduction to GODAE oceanview.

Journal of Operational Oceanography, 8, 2-11. <u>https://doi.org/10.1080/1755876X.2015.1022041</u>, 2015.

- Belmonte Rivas, M. and Stoffelen, A.: Characterizing ERA-Interim and ERA5 surface wind biases using ASCAT, Ocean Sci., 15, 831-852. <u>https://doi.org/10.5194/os-15-831-2019</u>, 2019.
 Benkiran, M., Le Traon, P.-Y., and Dibarboure, G.: Contribution of a constellation of two wide-swath altimetry missions to global ocean analysis and forecasting, Ocean Sci., 18, 609–625, https://doi.org/10.5194/os-18-609-2022, 2022.
 Chen, J., Ashton, I. G. C., Steele, E. C. C., and Pillai, A. C.: A Real-Time Spatiotemporal Machine Learning Framework for
- 160 the Prediction of Nearshore Wave Conditions. Artificial Intelligence for the Earth Systems, 2(1), e220033. https://doi.org/10.1175/AIES-D-22-0033.1, 2023.

Cossarini, G., Mariotti, L., Feudale, L., Mignot, A., Salon, S., Taillandier, V., Teruzzi, A., D'Ortenzio F.: Towards operational 3D-Var assimilation of chlorophyll Biogeochemical-Argo float data into a biogeochemical model of the Mediterranean Sea. Ocean Modelling, 133, 112-128. <u>https://doi.org/10.1016/j.ocemod.2018.11.005</u>, 2019.

165 Drake, P., Edwards, C. A., Arango, H. G., Wilkin, J., TajalliBakhsh, T., Powell, B., and Moore, A. M.: Forecast Sensitivitybased Observation Impact (FSOI) in an analysis–forecast system of the California Current Circulation. Ocean Modelling, 182, 102159. <u>https://doi.org/10.1016/j.ocemod.2022.102159</u>, 2023.

Fennel, K., Gehlen, M., Brasseur, P., Brown, C. W., Ciavatta, S., Cossarini, G., Crise, A., Edwards, C. A., Ford, D., Friedrichs,M. A. M., Gregoire, M., Jones, E., Kim, H.-C., Lamouroux, J., Murtugudde, R., Perruche, C., and the GODAE OceanView

170 Marine Ecosystem Analysis and Prediction Task Team: Advancing Marine Biogeochemical and Ecosystem Reanalyses and Forecasts as Tools for Monitoring and Managing Ecosystem Health. Frontiers in Marine Science, 6, 89. <u>https://doi.org/10.3389/fmars.2019.00089</u>, 2019.

Ford, D., Kay, S., Mcewan, R., and Totterdell, I. A.: Marine biogeochemical modelling and data assimilation for operational forecasting, reanalysis, and climate research. In "New Frontiers in Operational Oceanography", edited by E. Chassignet, A.

Pascual, J. Tintoré, and J. Verron (Tallahassee, FL: Florida State University), 625-652. <u>https://doi.org/10.17125/gov2018.ch22</u>, 2018.

Fu, L.-L., Pavelsky, T., Cretaux, J.-F., Morrow, R., Farrar, J. T., Vaze, P., et al.: The Surface Water and Ocean Topography Mission: A breakthrough in radar remote sensing of the ocean and land surface water. Geophysical Research Letters, 51, e2023GL107652. <u>https://doi.org/10.1029/2023GL107652</u>, 2024.

- 180 Fujii, Y., Rémy, E., Zuo, H., Oke, P., Halliwell, G., ... Usui, N.: Observing System Evaluation Based on Ocean Data Assimilation and Prediction Systems: On-Going Challenges and a Future Vision for Designing and Supporting Ocean Observational Networks. Frontiers in Marine Science, 6, 417. <u>https://doi.org/10.3389/fmars.2019.00417</u>, 2019. Gasparin, F., Guinehut, S., Mao, C. et al.: Requirements for an Integrated in situ Atlantic Ocean Observing System From
- Coordinated Observing System Simulation Experiments. Frontiers in Marine Science, 6, 83. https://doi.org/10.3389/fmars.2019.00083, 2019.
 - Gasparin, F., Hamon, M., Rémy, E., Le Traon, P. Y.: How deep argo will improve the deep ocean in an ocean reanalysis. Journal of Climate, 33(1), 77-94. <u>https://doi.org/10.1175/JCLI-D-19-0208.1</u>, 2020.

Gasparin, F., Lellouche, J.-M., Cravatte, S.E., Ruggiero, G., Rohith, B., Le Traon, P.Y., and Rémy, E.:: On the control of spatial and temporal oceanic scales by existing and future observing systems: An observing system simulation experiment

 approach. Frontiers in Marine Science, 10. <u>https://doi.org/10.3389/fmars.2023.1021650</u>, 2023.
 Gutknecht, E., Reffray, G., Mignot, A., Dabrowski, T., and Sotillo, M. G.: Modelling the marine ecosystem of Iberia–Biscay– Ireland (IBI) European waters for CMEMS operational applications. Ocean Sci., 15, 1489-1516. <u>https://doi.org/10.5194/os-15-1489-2019</u>, 2019.

Hamon, M., Greiner, E. Le Traon, P.-Y., and Remy, E ..: Impact of Multiple Altimeter Data and Mean Dynamic Topography

195 in a Global Analysis and Forecasting System. Journal of Atmospheric and Oceanic Technology, 36, 1255-1266. <u>https://doi.org/10.1175/JTECH-D-18-0236.1</u>, 2019.

Hauser, D., Abdalla, S., Ardhuin, F. et al.: Satellite Remote Sensing of Surface Winds, Waves, and Currents: Where are we Now? Surv. Geophys., 44, 1357-1446. <u>https://doi.org/10.1007/s10712-023-09771-2</u>, 2023.

Hernandez-Lasheras, J., Mourre, B., Orfila, A., Santana, A., Reyes, E., and Tintoré, J. (2021). Evaluating high-frequency radar
data assimilation impact in coastal ocean operational modelling. Ocean Sci., 17, 1157-1175. <u>https://doi.org/10.5194/os-17-1157-2021</u>, 2021.

Le Traon, P. Y.: From satellite altimetry to Argo and operational oceanography: three revolutions in oceanography. Ocean Sci., 9, 901-915. <u>https://doi.org/10.5194/os-9-901-2013</u>, 2013.

Le Traon, P.Y.: Satellites and operational oceanography. In "New Frontiers in Operational Oceanography", E. Chassignet, A. 205 Pascual, J. Tintoré, and J. Verron, Eds., GODAE OceanView, 161-190. DOI: 10.17125/gov2018.ch07

Le Traon, P.Y., Dibarboure, G., Jacobs, G., Martin, M., Remy, E. and Schiller, A.: Use of satellite altimetry for operational oceanography. In "Satellite Altimetry Over Oceans and Land Surfaces", CRC Press, Editors: Detlef Stammer and Anny Cazenave. <u>https://doi.org/10.1201/9781315151779</u>, 2017.

Levin, J., Arango, H.G., Laughlin, B., Hunter, E., Wilkin, J. and Moore, A.M.: Observation impacts on the Mid-Atlantic Bight

210 front and cross-shelf transport in 4D-Var ocean state estimates: Part II—The Pioneer Array. Ocean Modelling, 157, https://doi.org/10.1016/j.ocemod.2020.101731, 2021.

Martin M. J., King, R. R., While, J., and Aguiar, A.: Assimilating satellite sea surface salinity data from SMOS, Aquarius and SMAP into a global ocean forecasting system, Quarterly Journal of Royal Meteorological Society, 145, 705-726. https://doi.org/10.1002/qj.3461, 2019.

215 Mignot, A., Claustre, H., Cossarini, G., D'Ortenzio, F., Gutknecht, E., Lamouroux, J., Lazzari, P., Perruche, C., Salon, S., Sauzède, R., Taillandier, V., and Teruzzi, A.: Using machine learning and Biogeochemical-Argo (BGC-Argo) floats to assess biogeochemical models and optimize observing system design. Biogeosciences, 20, 1405-1422. <u>https://doi.org/10.5194/bg-20-1405-2023</u>, 2023.

Morrow R., et al: Global Observations of Fine-Scale Ocean Surface Topography with the Surface Water and Ocean 220 Topography (SWOT) Mission. Front. Mar. Sci. 6:232. doi: 10.3389/fmars.2019.00232, 2019.

Pasmans, I., Kurapov, A. L., Barth, J. A., Ignatov, A., Kosro, P. M., and Shearman, R. K.: Why gliders appreciate good company: Glider assimilation in the Oregon-Washington coastal ocean 4DVAR system with and without surface observations. Journal of Geophysical Research: Oceans, 124(1), 750-772. <u>https://doi.org/10.1029/2018JC014230</u>, 2019. Reul, N., Grodsky, S.A., Arias, M., Boutin, J., Catany, R., Chapron, B., D'Amico, F., Dinnat, E., Donlon, C., Fore, A. Fournier,

- S., Guimbard, S., Hasson, A., Kolodziejczyk, N., Lagerloef, G., Lee, T., Le Vine, D.M., Lindstrom, E., Maes, C., Mecklenburg, S., Meissner, T., Olmedo, E., Sabia, R., Tenerelli, J., Thouvenin-Masson, C., Turiel, A., Vergely, J.L., Vinogradova, N., Wentz. F., Yueh, S.: Sea surface salinity estimates from spaceborne L-band radiometers: An overview of the first decade of observation (2010–2019). Remote Sensing of Environment, 242, 111769. <u>https://doi.org/10.1016/j.rse.2020.111769</u>, 2019.
 Roemmich D, et al.: On the Future of Argo: A Global, Full-Depth, Multi-Disciplinary Array. Front. Mar. Sci. 6:439. doi:
- 230 10.3389/fmars.2019.00439, 2019.

Sakov, P., Counillon, F., Bertino, L., Lisæter, K. A., Oke, P. R., and Korablev, A.: TOPAZ4: an ocean-sea ice data assimilation system for the North Atlantic and Arctic. Ocean Sci., 8, 633-656. <u>https://doi.org/10.5194/os-8-633-2012</u>, 2012. Tranchant, B., Remy, E., Greiner, E., and Legalloudec, O.: Data assimilation of Soil Moisture and Ocean Salinity (SMOS)

observations into the Mercator Ocean operational system: focus on the El Niño 2015 event. Ocean Science, 15, 543-563. https://doi.org/10.5194/os-15-543-2019, 2019.

Turpin, V., Remy, E., and Le Traon, P. Y.: How essential are Argo observations to constrain a global ocean data assimilation system? Ocean Sci., 12, 257-274. <u>https://doi.org/10.5194/os-12-257-2016</u>, 2016.

Wang, B., and Fennel, K.: An Assessment of Vertical Carbon Flux Parameterizations Using Backscatter Data From BGC Argo. Geophys. Res. Lett., 50(3). <u>https://doi.org/10.1029/2022GL101220</u>, 2023.

Wang, B., Fennel, K., and Yu, L.: Can assimilation of satellite observations improve subsurface biological properties in a numerical model? A case study for the Gulf of Mexico. Ocean Sci., 17, 1141-1156. <u>https://doi.org/10.5194/os-17-1141-2021</u>, 2021.

Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from CryoSat-2 and SMOS in the Arctic reanalysis. The Cryosphere, 12, 3671-3691. https://doi.org/10.5194/tc-12-3671-2018, 2018.

245 Competing interests

The contact author has declared that none of the authors has any competing interests.

Data and/or code availability

Not applicable for this paper

Authors contribution

250 P.Y. Le Traon wrote the paper. A. Novellino and A. Moore provided feedback on the paper content.