

1 Solving Coastal Dynamics: Introduction to High Resolution Ocean

2 Forecasting Services

3 Joanna Staneva¹, Angelique Melet², Jennifer Veitch³, Pascal Matte⁴

4
5 ¹Institute of Coastal Systems - Analysis and ~~Modeling~~Modelling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

6 ²Mercator Ocean International, Toulouse, France

7 ³Egagasini Node, South African Environmental Observation Network (SAEON), Cape Town, South Africa

8 ⁴Meteorological Research Division, Environment and Climate Change Canada, Québec, QC, Canada

9
10 *Correspondence to:* Joanna Staneva (joanna.staneva@hereon.de)
11

Abstract. Coastal services are fundamental for society, with approximately 60% of the world's population living within 60 km of the coast. Thus, predicting ocean variables with high accuracy is a challenge that requires numerical models able to simulate from mesoscale to submesoscale processes, to capture shallow water dynamics influenced by wetting-drying and resolve the ocean variables in very high-resolution spatial domains. This paper ~~introduces key~~introduces key aspects of coastal modelling, such as vertical structure of the mixed layer depth, parameterization of bottom roughness and the dissipation of kinetic energy in coastal areas. It stresses the need for models to account ~~for the~~for the nonlinear interactions between tidal currents, wind waves, and small-scale weather patterns, emphasizing their significance in refining coastal predictions. In addition, observational advancements, such as high-frequency (HF) radar and satellite missions like SWOT, provide unique opportunities to observe coastal dynamics. This integration enhances our ability to model physical and dynamical peculiarities in coastal waters, estuaries, and ports. Coastal models not only benefit from such high-resolution observations but also contribute to evolving observational systems, creating feedback loops that refine monitoring and prediction capabilities. ~~Modeling~~Modelling strategies are also examined, including downscaling and upscaling approaches, and numerical challenges like implementing robust data assimilation schemes to refine estimations of coastal ocean states are addressed. Emerging techniques, such as advanced turbulence closure models and dynamic vegetation drag parameterization, are highlighted for their role in enhancing the realism of modeled coastal processes. Furthermore, the integration of atmospheric forcing, tidal asymmetries, and estuarine dynamics underlines the necessity for models that span the complexities of the coastal continuum. It also demonstrates the critical importance of accurately ~~modeling~~modelling coastal and estuarine systems to capture interactions between mesoscale and submesoscale processes, their connections to broader oceanic systems, and their implications for sustainable coastal management and climate resilience. This work underscores the potential of advancing coastal forecasting systems through interdisciplinary innovation, paving the way for enhanced scientific understanding and practical applications.

1 Introduction

High resolution observation and modelling are needed so that marine services can be compliant with small-scale processes in the ocean, particularly in coastal areas where these processes have a significant impact on dynamics and biogeochemistry (Figure 1). The importance of high resolution in coastal services is underscored by the coastal ocean's significance to humanity, not least because about 60% of the world's population lives within 60 km of the coast (Rao et al., 2008). These areas are highly dynamic, subject to both direct and indirect anthropogenic ~~pressures~~impacts, respectively, such as eutrophication, overfishing, offshore wind farm development, dredging, and pollution, ~~and natural drivers, including sea level rise,~~ global warming, sea level rise and changes in meteorological and hydrological conditions. These combined influences frequently trigger regime

43 shifts, coastal erosion, flooding, and the introduction of invasive species, underscoring the vulnerability and complexity of
44 these systems.

45 Accurately predicting ocean variables in coastal environments is challenging due to the need to resolve mesoscale to
46 submesoscale dynamics and their interactions with atmospheric and hydrological processes. The inherent variability of these
47 systems requires models that can account for a wide range of phenomena, including tidal asymmetries, wetting-drying cycles,
48 nonstationary river and atmospheric forcing, and nonlinear feedback mechanisms between tidal currents and wind waves
49 (Staneva et al., 2017). These processes influence mixing, ocean circulation, and the accuracy of sea surface temperature
50 predictions. Thus, high-resolution models are indispensable for capturing the fine-scale interactions that drive coastal dynamics
51 and shape biogeochemical responses.

52 Observational data play a pivotal role in advancing coastal ~~modeling~~modelling. High-frequency (HF) radar and novel high
53 resolution satellite missions offer unprecedented opportunities to observe and understand coastal processes with fine spatial
54 and temporal resolution (De Mey-Frémaux et al., 2019). These data sources are integral to improving the representation of
55 physical and biogeochemical variability in the models, bridging the gap between observations and predictive frameworks. By
56 integrating data from remote sensing and in situ platforms, coupled with advanced data assimilation techniques, models can
57 better capture the complexity of estuarine and nearshore processes.

58 Changes occurring in the coastal ocean are attributed to both direct human impacts and climate change. Human Anthropogenic
59 impacts encompass factors such as eutrophication, overfishing, offshore wind farm construction, dredging, and pollution.
60 Natural changes in the coastal ocean result from sea-level rise, global warming, and alterations in meteorological and
61 hydrological conditions such as precipitation, evaporation, wind patterns, and river run-off. These natural and human-induced
62 changes can lead to significant regime shifts, including alterations in biogeochemistry, increased coastal erosion, heightened
63 flooding risks, and the proliferation of invasive species, among other impacts.

64 Science-based services in the coastal ocean are essential for ensuring efficient management, sustainable use of coastal systems,
65 and the development of strategies that are adaptable to the changing climate, including sea-level rise. These efforts, for
66 example, align with the marine strategy framework directive in the European context (Hyder et al., 2015).

67 The aim of this paper is to introduce high-resolution ocean forecasting services that address the challenges of coastal dynamics
68 by improving predictions of physical and biogeochemical processes. It focuses on the integration of advanced
69 ~~modeling~~modelling techniques and modern observational tools to enhance understanding of small-scale dynamics and their
70 connections to larger ocean systems. The paper first describes the spatial scales and processes that high-resolution models
71 address, focusing on local, regional, and transitional zones. It then explores advanced observational tools, such as satellite
72 missions and HF radars, and their role in improving coastal forecasts. Following this, the discussion highlights numerical
73 ~~modeling~~modelling techniques, including turbulence ~~modeling~~modelling and bottom drag parameterization, which are
74 essential for capturing small-scale coastal dynamics. It also examines the role of data assimilation techniques and observing
75 system experiments in improving prediction accuracy and guiding the design of observation networks. Finally, the paper
76 concludes with a summary of findings, identifies current challenges, and outlines future directions for advancing coastal

forecasting systems. By addressing these topics, the paper aims to support the development of more robust and adaptable tools for coastal forecasting, which are critical for sustainable management and improving resilience to environmental changes.




Figure 1: Schematic representation of the coastal zone, hazards (e.g. HAB (harmful alga bloom), metocean and biogeochemical variables, as well as observations and applications (adapted from Melet et al., 2020).

2 Typical spatial scales and processes solved by high-resolution services

High-resolution services in the coastal ocean operate at various spatial scales depending on the specific applications and objectives. These scales can range from local to regional levels, aiming to capture fine-scale processes and variations. Here are some typical spatial scales for high-resolution services:

1. Local Scale: At the local scale, high-resolution services focus on small coastal areas, such as individual bays, estuaries, or nearshore zones. These services aim to provide detailed information and predictions for specific locations of interest. Spatial resolutions in this range can be on the order of meters to a few kilometers, allowing for precise observations and ~~modeling~~modelling of localized processes.

2. Coastal Scale: High-resolution services at the coastal scale cover larger coastal regions, spanning multiple bays, estuaries, and coastal zones. These services provide a broader view of the coastal environment and its dynamics. Spatial resolutions in this range typically range from ~~a few kilometers to tens of kilometers~~ meters to a kilometer, enabling the capture of coastal- to regional-scale variations and interactions.

3. Transition Zones: Transition zones refer to areas where coastal and open ocean processes interact. These zones often exhibit complex dynamics and are of particular interest for high-resolution services. Spatial resolutions in transition zones can vary

98 depending on the specific characteristics and objectives, but they generally aim to capture the intricate interactions between
99 coastal and open ocean processes.

100 A collection of 11 recent studies on operational coastal services utilizing high-resolution models offers significant insights into
101 the relevant spatial scales, objectives, and applications, thereby strengthening the analysis in this context (Sotillo, 2022). Eddies
102 or isolated vortices, meandering currents or fronts and filaments are characteristic features of oceanic mesoscale processes.
103 These processes typically exhibit spatial scales ranging from 10 to 500 kilometers, depending on geographic latitude and
104 stratification, and time scales ranging from several days to approximately 100 days. Submesoscale processes in the ocean, on
105 the other hand, are characterized by smaller scales, typically ranging from 1 to 10 kilometers (McWilliams, 2016). These scales
106 are smaller than the Rossby radius of deformation. Submesoscale processes also have shorter temporal scales, usually lasting
107 only a few hours, and their relative vorticity is greater than the Coriolis parameter f . In contrast, for mesoscale motion, the
108 relative vorticity is comparable to f . Overall, studying and observing submesoscale processes require advanced techniques and
109 methods to overcome their small scale and rapid variability, but their understanding is crucial for comprehending the intricate
110 dynamics of the ocean.

111 The surface and bottom mixed layers in the open ocean occupy just a tiny part of the ocean volume because these layers are
112 much thinner than the almost viscousless ocean interior. However, in the coastal zone, drag parameterizations become
113 increasingly important in shallow water, and even more so where the impact of vegetation is significant. Furthermore, a large
114 part of kinetic energy in the ocean is dissipated in the coastal zone, which necessitates an adequate modelling of this important
115 small-scale process, vital for the global energy balance (Munk and Wunsch, 1998). To accurately represent the coastal
116 dynamics and the fine structure of these layers, models need to resolve the vertical structure of the mixed layers. This
117 requirement necessitates the use of turbulence closure models, which account for the effects of turbulence and mixing in these
118 regions. Additionally, models for coastal processes need to consider the impact of bottom drag. The parameterization of bottom
119 roughness, often based on the grain size distribution, allows for the inclusion of bottom drag effects. In cases where vegetation
120 is present, drag parameterizations become even more important. A significant portion of the kinetic energy in the ocean is
121 dissipated in the coastal zone. Therefore, it is crucial to adequately model these small-scale processes in order to maintain a
122 balanced representation of the global energy dynamics. Understanding and accurately simulating the dissipation of kinetic
123 energy in coastal areas contribute to a comprehensive understanding of the ocean's energy budget.

124 In shallow water, the variability of surface elevation caused by tides and storms becomes comparable to the water depth itself.
125 In some coastal areas, shallow-water tides play a significant role in the overall tidal dynamics. To improve the accuracy of
126 tidal predictions in shelf regions, it is necessary to consider higher harmonics and assess the ability of ocean models to fully
127 resolve the tidal spectrum.

128 Some important processes, such as the nonlinear feedback between strong tidal currents and wind waves, cannot be ignored in
129 the coastal zone (Staneva et al., 2016a, 2016b, 2017). Wave-current coupling tends to decrease strong winds through wave-
130 dependent surface roughness (Wahle et al., 2017), affects mixing and ocean circulation, and improves predictions for sea
131 surface temperature. Further examples of the value of the incorporation of coupling in the numerical models in the coastal

ocean are given by De Mey-Frémaux et al. (2019). These scientific developments of operational oceanography are in pace with the trend in the Earth System modelling to seamlessly couple different environmental prediction components of atmosphere, waves, hydrology, and ice.

The small spatial scales characteristic of coastal and estuarine systems ~~require~~requires coastal models to consider ageostrophic (deviating from the Earth's rotation) and three-dimensional dynamics, primarily driven by boundary-layer processes (Fringer et al., 2019). Understanding these small-scale processes is crucial, particularly the interactions between mesoscale and sub-mesoscale dynamics and their connection to larger-scale processes. It is essential to improve the representation of exchanges between the coastal and open ocean, as well as their coupling with estuaries and catchment areas, in order to capture the complexity of coastal systems. Accounting for high-resolution atmospheric forcing ~~into~~in the coastal models is essential for accurately capturing local meteorological dynamics, including wind patterns, temperature gradients, and precipitation rates. Such detailed atmospheric data drive fundamental processes like heat and momentum fluxes, profoundly influencing coastal hydrodynamics, sediment transport, and ecosystem responses. The implementation of a novel high-resolution atmospheric forcing, combined with the refinement of bulk formulae for surface flux computations, significantly enhances the performance of various high-resolution modelling systems for port environments (García-León et al.). Coastal models need to accurately account for frictional balances, taking into consideration the effects of friction on the movement of water. They must also address wetting and drying processes, as well as hydrological forcing, to capture the transitions between shallow environments and larger regional scales. By incorporating these factors, models can provide a more realistic representation of coastal dynamics. In addition, the grid characteristics used in coastal models should be carefully selected to accurately represent the dominant spatial scales present in the coastal environment. Choosing grid resolutions that capture the essential features of the coastal system is crucial for obtaining reliable and meaningful results.

In the coastal ocean, characteristic time scales are significantly shorter compared to the global ocean. These time scales, typically around 1 day, are determined by various processes, including tides, inertial motion, diurnal cycles, and synoptic weather patterns. The fast-paced dynamics of the coastal ocean require models to accurately capture these shorter time scales. In estuaries, the periodicity becomes more complex due to strong tidal asymmetries and the presence of secondary circulation patterns. The interactions between tidal forcing, river flow and estuarine geometry result in intricate and variable periodic patterns. (as shown in Campuzano et al. 2022 for the Western Iberian Buoyant Plume, Sotillo et al. 2021 for the whole European Atlantic façade, Pein et al. 2021 for the Elbe Estuary). The periodicity observed in coastal seas is mainly influenced by external forcing signals, such as atmospheric conditions or remote ocean signals. These external signals propagate ~~into~~in the coastal models through the specification of lateral boundary conditions, which is a crucial aspect of ~~modeling~~modelling in coastal areas. Unlike global models that can operate ~~with~~without open boundaries, coastal models require careful consideration of these boundary conditions to accurately represent the interactions between the coastal and open ocean.

The predictability limit of models depends on the geophysical processes. For synoptic processes in the open ocean, this limit is on the order of weeks to months. For the coastal ocean, it is on the order of hours to days. The loss of predictability, associated with nonlinear processes, is exemplified by the growth of errors in predictive models. Assimilation of data containing spatial

and temporal scales below the predictability limit is needed to address this issue. Simulations at grid resolutions that would sufficiently resolve the coastal submesoscale would require horizontal grid resolutions of approximately 1-10 meters in estuaries and 0.1-1 kilometer in coastal shelf domains. However, achieving such high resolutions poses significant computational challenges and resource demands.

By employing high-resolution services with appropriate spatial scales, scientists and stakeholders can gain a more detailed and accurate understanding of coastal processes, improve forecasting capabilities, and support effective coastal management and decision-making.

3 State-of-the-art data and tools for coastal forecasting

3.1 Required observations

Observing systems are spatiotemporally sparse in coastal regions compared to the small scales of ecosystem variability found there. A crucial challenge in observations is addressing the variety of important spatial and temporal scales within the coastal continuum, which encompasses the seamless transition from the deep ocean to estuaries through the shelf. In order to achieve this, observations should sample the multiscale, two-way interactions of estuarine, nearshore, and shelf processes with open ocean processes. Additionally, they need to account for the different pace of circulation drivers, such as fast atmospheric and tidal processes, as well as the slower general ocean circulation and climate forcing. It is also important to accurately sample the gradients of biological production, ranging from mesotrophic estuaries to oligotrophic oceans. Given the current situation, observational practices and strategies need to be strongly coupled with numerical ~~modeling~~modelling to effectively extract the information contained in the data and advance the quality of coastal services.

Most global and regional prediction products use a combination of satellite observations and in situ observations. Traditionally, in situ observations constituted the major data source for coastal ocean monitoring. During the end of the past century, satellite observations contributed significantly to the understanding of spatial variabilities. Novel instruments, such as the acoustic Doppler current profiler (ADCP), which measures current profiles throughout the water column, enhanced our understanding of current shear and bottom stress. Nowadays, high-resolution numerical simulations in the coastal ocean are keeping pace with high-resolution observations. A similar trend is observed in coastal waters, estuaries, and ports, which are rich in different activities and interests: fishing, recreational activities, search and rescue, protection of habitats, storm forecasts, maritime industries, as well as routine maintenance operations (De Mey-Frémaux et al., 2019).

The coastal ocean observations only are not sufficient to fully support the present-day need for high-quality ocean forecasting and monitoring because measurements may represent very localized and short scale dynamics, and it is not straightforward to know how fully they describe the complex coastal system. Therefore, recent practices employ the synergy between observations and numerical ~~modeling~~modelling, which ensures valuable research advancements and practical implementations (Kourafalou et al., 2015a, 2015b). The core components of operational oceanographic systems consist of a multi-platform observation network, a data management system, a data assimilative prediction system, and a dissemination/accessibility

system (Kourafalou et al., 2015a; De Mey-Frémaux et al., 2019; Davidson et al., 2019). By combining observations and models through data assimilation methods, ranging from coastal to global and from in situ to satellite-based, we can assess ocean conditions and create reliable forecasts. This integration adds value to coastal observations and enables a wide range of applications (De Mey-Frémaux et al., 2019; Ponte et al., 2019), as well as providing decision-making support. For a comprehensive review of ocean monitoring and forecasting activities in both the open and coastal oceans, please refer to Siddorn et al. (2016).

High-frequency radars (HFR) offer unique spatial resolution by providing reliable directional wave information and gridded data of surface currents in almost real time. The use of HFR networks has become an essential element of coastal ocean observing systems, contributing to high-level coastal services (Stanev et al., 2016; Rubio et al., 2017); Reyes et al., 2022). The outputs from prediction systems extend the utility of HFR observations beyond the immediate observation area (Stanev et al., 2015), enabling adequate estimates even where no direct observations have been made. This demonstrates how models connect observations, synthesize them, and assist in the design of observational networks. In turn, observations can guide the development of coastal models (De Mey-Frémaux et al., 2019).

Alongside ADCP data, HFR data are used for skill assessment of operational wave and circulation models (Lorente et al., 2016). Another valuable source of fine-resolution data in the coastal region is provided by color data from satellites. In terms of sea level observations, some challenges associated with the use of altimeter data in the coastal zone are expected to be overcome through the use of wide-swath Surface Water and Ocean Topography (SWOT) technology. SWOT provides-is a landmark satellite mission that delivers two-dimensional sea surface height observations of sea surface topography at kilometer-high resolution in across a 120 km swath. The expectation is-It represents a major step forward in resolving mesoscale and submesoscale features critical to coastal dynamics. Recent Observing System Simulation Experiments (OSSEs) have demonstrated that wide-swath altimetry will significantly enhance the quality of coastal-substantially enhances ocean forecasts through high spatial coverage and resolution. These new data are anticipated to (1) sample the forecasting capabilities. For instance, a constellation of two-way interactions between nearshore, estuarine, SWOT-like wide-swath altimeters provides a ~14% reduction in sea surface height forecast error compared to a 12-nadir altimeter constellation and shelf processes also improves estimates of surface currents and open-ocean processes, and (2) resolve both fast processes (atmospheric, hydrologic, tidal) and slower processes (general ocean circulation and climate evolution). High resolution models play a critical role as integrators Lagrangian trajectories (Benkiran et al., 2024). These results highlight the importance of these coastal continuum SWOT-type observations for resolving small-scale coastal variability and improving model-data integration. Further advances in coastal observations are enabled by autonomous platforms such as Slocum gliders. These gliders can carry a wide array of physical and biogeochemical sensors and perform repeated transects, thus providing high-resolution observations of dynamic features such as eddies, frontal systems, and upwelling events. Their operational flexibility and ability to collect subsurface data make them valuable for both sustained monitoring and adaptive sampling strategies (Rudnick, 2016; Testor et al., 2019). In parallel, satellite technologies continue to evolve. Moreover, the Japanese geostationary meteorological satellite Himawari-8 provides high-frequency (every 10 minutes) and high-resolution (up to 500 m) visible and infrared

imagery. These capabilities allow for near-real-time monitoring of sea surface temperature (SST), making it possible to track rapidly evolving coastal phenomena such as diurnal warming, river plumes, and thermal fronts (Kurihara et al., 2016). These complementary in situ and remote sensing platforms represent essential components of integrated coastal observing systems, supporting the growing demand for accurate forecasts, early warnings, and data-driven decision-making tools.

3.2 Numerical models

Addressing specific processes in the coastal ocean and accurately ~~modeling~~modelling the transition between regional and coastal scales cannot be achieved solely by adjusting the model resolution. Certain processes, such as shallow-water tides, which are often overlooked in global and regional forecasting, play a dominant role in coastal ocean dynamics. The previous sections have highlighted the importance of a tailored approach in observational practices and numerical models for the coastal ocean. For further information on other popular coastal models, refer to the comprehensive discussion by Fringer et al. (2019).

Table 1: Circulation models in alphabetical order, which can be used for coastal and regional studies and/or provision of services.

Model	Citationan	C: Coastal, R: Regional, G: Global	Finte-volume (FV) or Finite-element (FE)
ADCIRC	Luettich et al. (1992) Westerink et al. (1994)	C	FE
COAWST	Warner et al. (2008, 2010)	C/R	FV
<u>COMPAS</u>	<u>Herzfeld et al. (2020)</u>	<u>C/R</u>	<u>FV</u>
CROCO	Marchesiello et al. (2021)	<u>C/R</u>	FV
Delft3D	Delft3D-Flow User Manual (2024)	C	FV
FVCOM	Chen et al. (2003)	C/R/G	FV
GETM	Burchard and Bolding (2002)	C	FV
MITgcm	Marshall et al. (1997)	C/R/G	FV
<u>MPAS</u>	<u>Ringler et al. (2013)</u>	<u>R/G</u>	<u>FV</u>
<u>NEMO</u>	<u>Madec et al., (2016)</u>	<u>C/R/G</u>	<u>FV</u>
<u>POMS</u>	<u>Blumberg and Mellor (1987), Mellor (2004)</u>	<u>C/R</u>	<u>FV</u>

ROMS	Shchepetkin and McWilliams (2005)	R	FV
SCHISM	Zhang et al. (2016)	C/R/G	FV/FE
SELFE	Zhang and Baptista, 2008	C	FV/FE
SHYFEM	Umgiesser et al. (2004)	C	FE
SUNTANS	Fringer et al. (2006)	C	FV
TRIM/UnTRIM	Casulli (1999), Casulli and Zanolli (2002, 2005)	C	FV

3.3 Fine resolution nested models, downscaling and upscaling

High-resolution coastal services must properly resolve interactions between various coastal processes, including nearshore, estuarine, shelf, drying, and flooding dynamics. Achieving this requires a resolution of approximately 10-100 meters. Simultaneously, it is essential to capture open ocean processes at a resolution of around 1 kilometer or coarser. Common approaches employed in addressing this challenge include downscaling and multi-nesting techniques (e.g., Debreu et al., 2012; Kourafalou et al., 2015b; Trotta et al., 2017) as well as the use of unstructured-grid models (e.g., Zhang et al., 2016a, 2016b; Federico et al., 2017; Stanev et al., 2017; Ferrarin et al., 2018; Maicu et al., 2018). Another important aspect to consider is upscaling (Schulz-Stellenfleth and Stanev, 2016), which becomes relevant when addressing the two-way interaction between coastal and open-ocean systems.

Most coastal models are one-way nested, relying heavily on forcing data from larger-scale models as the coastal system is primarily influenced by the atmosphere, the hydrology and the open ocean. Enhancing the horizontal resolution of the North Sea operational model from 7 to 1.5 kilometers (Tonani et al., 2019) has shown improvements in off-shelf regions, but biases persist over the shelf area, indicating the need for further enhancements in surface forcing, vertical mixing, and light attenuation.

An important consideration in downscaling and coastal modelling is the treatment of open boundary conditions (OBCs), which play a critical role in determining model fidelity near the boundaries. OBCs are typically derived from larger-scale models but often require case-specific tuning to ensure dynamic consistency and minimize reflection or spurious signals. The choice and configuration of OBCs—such as Flather-type, radiation conditions, or relaxation zones—can significantly affect the transport and energy balance within the coastal model domain. Given the diversity of physical processes and geometries encountered in coastal environments (Marchesiello et al., 2001). Models equipped with a wide suite of configurable boundary condition types offer a practical advantage, particularly in multi-scale coupled frameworks. Ensuring consistency across nested domains while

preserving physical realism remains an ongoing challenge, motivating continued development and intercomparison of OBC strategies in operational and research settings.

While the downscaling of information from coarser global or regional models to high-resolution coastal models is well-established, the reverse process of upscaling is more challenging and continues to be a subject of research. Two-way nested models allow assimilated information from coastal observations, typically not assimilated by larger-scale forecasting systems, to propagate beyond the coastal region while maintaining dynamic consistency. This upscaling capability has the potential to benefit regional models. Coastal observations have demonstrated their potential to improve boundary forcing or surface wind forcing in regional models.

The coupling of a coarse-resolution regional model with a fine-resolution coastal model using a two-way nesting approach has been studied in the context of the straits connecting the North and Baltic Seas. The intricate topography and narrow cross-sections of the straits result in the dominance of small-scale motions, which play a vital role in the exchange between the two seas and significantly influence Baltic Sea stratification. The two-way nesting method, ~~developed by exchanging design to~~ exchange information between the child model in the straits and the parent model in the seas, incorporates elements of data assimilation and allows for different vertical discretizations in each model. The Adaptive Grid Refinement in FORTRAN (AGRIF), originally developed by Debreu et al. (2008; 2012), has found wide application as a library for seamless spatial and temporal refinement over rectangular regions in the NEMO ~~modeling~~modelling framework (<https://forge.ipsl.jussieu.fr/nemo/wiki/WorkingGroups/AGRIF>).

Recent advancements in two-way nesting frameworks have demonstrated their effectiveness in improving multi-scale model accuracy. The implementation of a general two-way nesting framework has enhanced the exchange of physical properties between nested grids while preserving numerical stability and computational efficiency. Additionally, the integration of two-way nesting in a global ocean model has significantly improved surface tidal accuracy, refining regional tidal dynamics without compromising large-scale coherence (Herzfeld & Rizwi, 2019; Jeon et al., 2019). Further applications of AGRIF have demonstrated improvements in hydrodynamic simulations and the estimation of environmental indicators in coastal systems, underscoring its potential to refine fine-scale hydrodynamics while ensuring consistency with larger-scale ocean processes (Petton et al., 2023).

The organization of these multi-model studies is identified by the coastal modelling community as a need. Firstly, to tackle common assessments of the wide range of overlapping (global/basin/regional and local) models that are available for users in some coastal zones. Secondly, these multi-model validation exercises, comparing the performance of global/regional “core” model forecasts (i.e. from services such as the Copernicus Marine one) and coastal model solutions, nested into the formers, are useful to identify the potential added value (and the limitations) of performed coastal downscaling with respect to the “parent” core operational solutions, in which high-resolution coastal models are nested.

Frishfelds et al. (2025) highlight the benefits of on-demand coastal modeling employing two-way nesting, emphasizing its capacity to dynamically refine coastal processes while maintaining consistency with larger-scale ocean simulations. This

approach enhances the accuracy and reliability of high-resolution forecasting systems, facilitating improved representation of fine-scale coastal dynamics.

In that sense, these multi-model intercomparison exercises are key elements for many initiatives, such as the Horizon Europe Project FOCCUS (Forecasting and Observing the Open-to-Coastal Ocean for Copernicus Users, <https://foccus-project.eu/>) Project, that have in their core the enhancing of existing coastal downscaling capabilities, developing innovative coastal forecasting products based on a seamless numerical forecasting from regional models of the Copernicus Marine Service covering the EU regional seas, to Member States coastal forecasting systems (authors can add here any other pertinent reference from literature). Espino et al. (2022) emphasized the significance of extending Copernicus Marine Environmental Monitoring Service (CMEMS) products to coastal regions, highlighting the integration of high-resolution models and observational data to improve coastal forecasting capabilities. Their work underscores the importance of tailoring operational ocean models to better capture nearshore dynamics, ensuring more accurate and actionable predictions for end-users. Furthermore, multi-model studies focused on extreme event simulations provide valuable insights into the performance of operational forecasting systems. For instance, Sotillo et al. (2021) examined the record-breaking Western Mediterranean Storm Gloria by evaluating five different model systems, including Copernicus Marine Service products (global, regional Mediterranean, and Atlantic IBI solutions) alongside two coastal nested models. Such studies play a crucial role in assessing model accuracy, leveraging local HF radar observations, and informing future improvements to regional and coastal forecasting services. Furthermore, and from an end-user perspective, multi-model study cases focused on extreme event simulations, such as the one performed by Sotillo et al. (2021) focused on the record-breaking Western Mediterranean Storm Gloria, allow to identify strengths and limitations of model solutions delivered by operational forecast services available in zones affected by extreme events; for instance, in the referred study case, 5 model systems were considered (including systems both from the Copernicus Marine service -with usages of the Global and the regional Mediterranean and Atlantic IBI solutions- and 2 coastal services nested into the regional solutions). This kind of multi-model study cases certainly help to enhance product quality assessments (in this Gloria Storm case, making extensive use of the local HF radar capabilities), increasing the knowledge about the model systems in operations, and outlining future model service upgrades (both in the regional and coastal services) aimed at achieving a better coastal forecasting of extreme events.

3.4 Unstructured-Grid Models for Cross-Scale Coastal Dynamics

The use of unstructured-grid models is crucial for cross-scale ~~modeling~~modelling and effectively addressing the interactions between estuaries and the open ocean. One key aspect is the accurate representation of freshwater transformation from rivers, which is often oversimplified in ocean models by specifying river runoff as a point source. Unstructured-grid models, ~~with their ability to employ higher~~while often employing lower-order spatial discretizations, ~~demonstrate good skill due to interpolation complexities on irregular meshes, provide enhanced flexibility in capturing~~resolution placement and transition zones. This allows them to effectively capture subtidal, tidal, and intermittent processes in coastal and estuarine environments,

providingsupporting a more realistic representation of estuarine processesdynamics and improved interfacecoupling with estuarine models.

Compared to curvilinear and Cartesian grids, unstructured grids excel in resolving complex bathymetric features without significant grid stretching. Since bathymetry plays a fundamental role in governing the dynamics of estuaries and the near coastal zone, unstructured grid models offer greater accuracy and computational efficiency in numerical forecasting. Their flexibility also enables more effective resolution of multiscale dynamic features. Fine spatial resolution in unstructured-grid models allows for the resolution of secondary (transversal) circulation in estuaries and straits, thereby improving mixing and enhancing the representation of long-channel changes in stratification, as demonstrated by Haid et al. (2020). Zhang et al. (2016) have emphasized the role of cross-scale modeling in capturing multi-scale hydrodynamic interactions, particularly in tidal straits, where unstructured-grid models enhance the representation of exchange flows and stratification dynamics. As Ilicak et al. (2021) have shown, these advancements contribute to more precise simulations of estuarine and strait dynamics. Recent research has further elucidated the mechanisms governing secondary circulation in tidal inlets. Chen et al. (2023) demonstrated that subtidal secondary circulation can arise due to the covariance between eddy viscosity and velocity shear, even in predominantly well-mixed tidal environments. This finding highlights the necessity of incorporating high-resolution turbulence parameterizations within unstructured-grid models to accurately capture submesoscale and cross-channel processes, thereby improving the fidelity of numerical simulations in complex coastal and estuarine systems.

However, the construction of grids and ensuring reproducibility in unstructured-grid modelingmodelling still present challenges. Grid generation is not always fully automated, and subjective decisions are often made based on the specific research problem, applications, and intended services. The development of more objective grid construction methods and reproducibility standards is an ongoing concern in unstructured-grid modeling (Candy and Pietrzak, 2018)-modelling (Candy and Pietrzak, 2018). One significant advancement is the introduction of the JIGSAW mesh generator (Engwirda, 2017), which enables the creation of high-quality unstructured grids designed to satisfy specific numerical requirements. JIGSAW produces centroidal Voronoi tessellations with well-centred, orthogonal cell geometries that are particularly suitable for mimetic finite-volume schemes. JIGSAW incorporates mesh optimisation strategies tailored to geophysical fluid dynamics and has been increasingly adopted in ocean modelling applications.

The generation of unstructured meshes is a critical component in configuring coastal and estuarine ocean models, as it directly influences numerical accuracy, computational efficiency, and the ability to represent complex shoreline and bathymetric features. Tools such as OceanMesh2D offer MATLAB-based workflows for high-quality, two-dimensional unstructured mesh generation, facilitating user control over mesh density and coastal geometry resolution (Roberts et al., 2019). Similarly, OPENCoastS provides an open-access, automated service that streamlines the setup of coastal forecast systems, integrating mesh generation, model configuration, and forecast production (Oliveira et al., 2019, 2021). The OCSMesh software developed by NOAA represents another important advancement. It enables data-driven, automated unstructured mesh generation tailored for coastal ocean modeling, offering a robust framework to ensure mesh quality, reproducibility, and interoperability with

NOAA modeling systems (Mani et al., 2021). Together, these developments represent the ongoing progress toward objective, reproducible, and user-oriented mesh generation in support of high-resolution coastal ocean modelling.

3.5 Observing System Simulation Experiments, Observing System Experiments and Data Assimilation

Data assimilation in coastal regions presents challenges due to the presence of multiple scales and competing forcings from open boundaries, rivers, and the atmosphere, which are often imperfectly known (Moore and Martin, 2019). However, studies Data assimilation is particularly challenging in tidal environments (especially for meso- and macro-tidal environments; and not so in micro-tidal coastal zones (De Mey et al., 2017, Stanev et al, 201, Holt et al, 2012). Studies by Oke et al. (2002), Wilkin et al. (2005), Shulman and Paduan (2009), Stanev et al. (2015, 2016), and others have demonstrated the value of assimilating HF radar observations to improve the estimation of the coastal ocean state.

Observing System Simulation Experiments (OSSE) and Observing System Experiments (OSE) are widely used techniques for assessing and optimizing ocean observational systems. OSSEs involve numerical simulations that test the potential impact of hypothetical observations on forecast models before actual observations are made, enabling improved planning and cost-effective observational strategies. In contrast, OSEs assess the impact of existing observations by systematically removing certain datasets from assimilation systems and evaluating the resulting degradation in model performance. OSSE and OSE have the capability to incorporate diverse observing systems, including satellite-based observations, HF radars, buoys with low-cost sensors, autonomous vehicles, and more. These approaches, when coupled with an assimilative system, can provide guidance on optimizing the observing network (Oke and Sakov, 2012; Fujii et al. These approaches provide valuable insights for refining data assimilation techniques and guiding the development of future observational networks. For further details, we refer readers to Oke and Sakov (2012) and Fujii et al. (2019), who provide comprehensive discussions on the methodologies and applications of OSSEs and OSEs in operational oceanography. an in-depth review of OSSE methodologies and insights into how OSSE and OSE methodologies contribute to improving ocean forecasting, designing observational systems, and refining numerical models is given in Zeng et al (2020). These approaches, 2019). They can help identify gaps in existing coastal observing networks, assess operational failure scenarios, and evaluate the potential of future observation types. Pein et al. (2016) used an OSE-type approach to investigate the impact of salinity measurements in the Ems Estuary on the reconstruction of the salinity field, identifying observation locations that are more suitable for model-data synthesis. This type of analysis can contribute to the design and optimization of both existing and future observational arrays, especially in coastal regions where fine resolution is required.

3.5 Riverine forcing and its role in coastal ocean Modeling

Rivers play a critical role in shaping coastal circulation and stratification by delivering freshwater, nutrients, and sediments that influence estuarine and shelf dynamics. The treatment of riverine inputs in ocean models remains a key source of uncertainty, especially when estuarine plume dynamics and mixing processes are unresolved. In many coarse-resolution

systems, river discharge is prescribed via simplified surface or salinity fluxes, which may misrepresent the spatial structure and strength of river plumes (Sun et al., 2017; Verri et al., 2020). To address this, high-resolution and regional-scale models increasingly incorporate momentum-carrying river inflows or artificial estuarine channels (Herzfeld, 2015; Sobrinho et al., 2021). For instance, Nguyen et al. (2024) demonstrated how high-resolution modeling in the German Bight captures the hydrodynamic and biogeochemical responses to extreme river discharge events, showing significant implications for salinity, stratification, and nutrient dispersion during floods. These findings underscore the importance of resolving riverine inflow variability and extreme events in coastal ocean prediction systems.

Recent work has also focused on operational strategies for river forcing (Matte et al., 2024), including real-time discharge data integration (e.g., from GloFAS; Harrigan et al., 2020), and estuary box models that approximate sub-grid plume behavior (Sun et al., 2017). These approaches aim to enhance predictive capabilities while maintaining computational feasibility in global-to-coastal modeling chains. Choosing the appropriate river input strategy is therefore application-dependent and strongly influenced by spatial resolution and target phenomena.

3.6 Integration of AI in Coastal Modeling and Forecasting

The integration of artificial intelligence (AI) and machine learning (ML) techniques in ocean and coastal forecasting has rapidly evolved, providing novel methodologies for improving predictive accuracy, computational efficiency, and data assimilation in operational models. Recent advances in AI-based approaches for parameterizing subgrid-scale processes, hybrid modelling techniques, and ensemble forecasting highlight the transformative potential of these methods in coastal modelling (Heimbach et al., 2024).

Machine learning applications in coastal ocean modeling primarily focus on two domains: (1) enhancing conventional physical models by integrating ML-based parameterizations and error corrections, and (2) fully data-driven approaches that employ neural networks as surrogate models (Zanna & Bolton, 2020; Bolton & Zanna, 2019). The former leverages ML techniques to optimize numerical model performance by improving subgrid parameterizations, bias correction, and data assimilation strategies, while the latter explores the potential of deep learning algorithms such as Fourier Neural Operators (FNOs) and Transformer-based architectures for high-resolution ocean forecasting (Bire et al., 2023; Wang et al., 2024).

Data assimilation, a critical component of operational forecasting, benefits from AI-enhanced methodologies that improve state estimation and predictive skill. AI-driven data assimilation frameworks, such as the combination of deep learning with variational assimilation (4D-VarNet) (Fablet et al., 2022), have demonstrated superior performance in coastal and regional models. Hybrid approaches incorporating AI techniques into numerical models have been applied to refine coastal simulations, allowing for better representation of multi-scale interactions (Brajard et al., 2021). Furthermore, convolutional neural networks (CNNs) have been successfully used for downscaling sea surface height and currents in coastal areas, addressing challenges related to observational gaps and improving model resolution (Yuan et al., 2024).

Coastal high-resolution models often suffer from errors stemming from inaccuracies in numerics, forcing (e.g., open boundaries, meteorological inputs), and unresolved physical processes. AI-based methods have been increasingly applied to

address these challenges, particularly in the realm of subgrid-scale parameterization. AI-enabled parameterizations of mesoscale and submesoscale processes using deep learning techniques, such as residual networks and generative adversarial networks (GANs), have shown promising results in reducing bias in numerical simulations (Gregory et al., 2023; Brajard et al., 2021). Additionally, hybrid methods combining physics-based models with ML correction schemes have demonstrated improved predictive skill for regional and coastal ocean models (Perezhogin et al., 2023).

The use of ML for extreme event prediction has gained increasing attention in the context of operational coastal forecasting. AI models trained on historical storm data and high-resolution numerical simulations have been utilized to enhance storm surge predictions and improve early warning systems (Xie et al., 2023). Transformer-based models, originally developed for atmospheric forecasting, have been adapted for ocean applications, achieving competitive skill in eddy-resolving ocean simulations (Wang et al., 2024).

The integration of AI in ensemble forecasting further contributes to uncertainty quantification, providing probabilistic predictions for extreme coastal events. Bayesian inference techniques, combined with ML-based ensemble prediction, offer a framework for optimizing multi-model ensembles and reducing systematic errors in operational forecasts (Bouallègue et al., 2024; Penny et al., 2022). The synergy between ML-driven emulators and traditional ensemble forecasting techniques has the potential to enhance coastal hazard predictions, particularly in regions prone to high-impact events.

Despite the advancements in AI for coastal modeling, several challenges remain. The interpretability and robustness of ML-based solutions need further improvement, particularly for operational applications requiring high levels of reliability (Bonavita, 2023). Additionally, integrating ML models with real-time observational data streams, including remote sensing and high-frequency radar (HFR) networks, remains an ongoing area of research (Reichstein et al., 2019). The extension of ML-based ocean forecasting to seasonal and interannual time scales also poses challenges related to long-term stability and physical consistency (Beucler et al., 2024).

4 Summary and outlook

The critical importance of high-resolution coastal ~~modeling~~~~modelling~~ is demonstrated in addressing the complexities of dynamic coastal systems. Coastal areas are shaped by the interplay of mesoscale and submesoscale processes, strong tidal currents, atmospheric and hydrologic forcing, and significant anthropogenic pressures. Advanced techniques, including turbulence closure models for capturing vertical mixing and parameterizations of bottom roughness and vegetation drag for representing energy dissipation, are essential for accurately ~~modeling~~~~modelling~~ these systems. The nonlinear interactions between tidal currents and wind waves emerge as a particularly influential factor, affecting ocean circulation and improving the accuracy of sea surface temperature predictions.

It is shown that the integration of high-resolution observational data, such as HF radar for surface currents and the SWOT satellite mission for sea surface topography, has the potential of substantially enhancing the resolution and reliability of coastal models. These data facilitate a detailed characterization of processes in transition zones spanning estuaries, nearshore areas,

and the open ocean. Improved coupling between regional and local models has advanced the representation of boundary conditions and enabled simulations of small-scale dynamics, essential for capturing the complexity of the coastal continuum. The application of data assimilation techniques addresses the rapid variability inherent in coastal processes, highlighting the challenges and limitations of predictability in these highly dynamic environments. Strategies to extend the accuracy of short-term and localized forecasts are provided, leveraging multiscale data integration to refine predictions. The ability to simulate interactions between atmospheric conditions, hydrological inputs, and oceanographic processes strengthens the foundation for more accurate [modelingmodelling](#). This contribution underscores the importance of bridging observational and [modelingmodelling](#) gaps to achieve a comprehensive understanding of coastal systems. It highlights the necessity of integrating small-scale dynamics with broader processes to better inform sustainable coastal management practices. By aligning advanced techniques with high-resolution data, this work offers a pathway for more robust representations of coastal ocean dynamics and supports informed decision-making in the face of growing environmental and societal challenges.

Several directions for advancing coastal ocean modelling to address evolving environmental and societal challenges are highlighted. Future efforts should focus on integrating emerging observational technologies, such as high-resolution satellites (e.g., SWOT), autonomous platforms like gliders and drones, and hyperspectral imaging. These tools, combined with machine learning techniques for data analysis, can bridge gaps in spatial and temporal data coverage, providing a richer understanding of coastal dynamics.

Developing coupled [modelingmodelling](#) systems that seamlessly integrate atmospheric, hydrological, and oceanographic processes will be essential for capturing the complexities of the land-ocean continuum. Incorporating river runoff, estuarine dynamics, and nearshore processes into such systems will significantly enhance the scope and accuracy of predictions. Addressing computational challenges associated with high-resolution [modelingmodelling](#) is equally critical; this includes leveraging high-performance computing, cloud-based processing, and optimizing numerical schemes to achieve efficient and precise simulations.

Improving data assimilation techniques through ensemble approaches and probabilistic forecasting is another priority. These methods will better integrate multiscale observational data, reduce uncertainties, and enhance the reliability of predictions in dynamic environments. Concurrently, there is a pressing need to explore the impacts of climate change on coastal systems, including sea-level rise, increased storm intensity, and shifting precipitation patterns. Understanding these impacts will guide the development of adaptive strategies and strengthen resilience in vulnerable coastal zones.

The future of coastal [modelingmodelling](#) also depends on fostering interdisciplinary collaboration, engaging expertise from oceanography, meteorology, hydrology, and ecology. By aligning scientific research with societal needs and practical applications, collaborative frameworks can ensure the relevance and effectiveness of [modelingmodelling](#) efforts. Additionally, applying artificial intelligence to optimize model parameterization, grid design, and predictive analyses will unlock new capabilities for simulating small-scale processes like sediment transport and ecosystem responses.

Finally, enhancing global and regional coordination for coastal monitoring and [modelingmodelling](#) will be vital. Strengthening networks to ensure consistency in data and [modelingmodelling](#) approaches can foster international collaboration, facilitating

494 the exchange of best practices and resources. These collective advancements promise to deepen our understanding of coastal
495 systems and provide robust tools to manage and protect these critical areas sustainably in the face of ongoing and future
496 challenges.
497
498
499
500
501
502

- Benkiran, M., Le Traon, P.-Y., Rémy, E., & Drillet, Y. (2024). Impact of two high-resolution altimetry mission concepts on ocean forecasting. *Frontiers in Marine Science*, 11, 1465065. <https://doi.org/10.3389/fmars.2024.1465065>
- Blumberg, A. F., & Mellor, G. L. (1987). Title: A description of a three-dimensional coastal ocean circulation model. Book Chapter: In *Three-dimensional coastal ocean models* (pp. 1-16). Publisher: American Geophysical Union. DOI: 10.1029/CO004p0001
- Campuzano, F., Santos, F., Simionesei, L., Oliveira, A. R., Olmedo, E., Turiel, A., Fernandes, R., Brito, D., Alba, M., Novellino, A., & Neves, R. (2022). Framework for Improving Land Boundary Conditions in Ocean Regional Products. *Journal of Marine Science and Engineering*, 10(7), 852. <https://doi.org/10.3390/jmse10070852>
- Candy, A.S., Pietrzak, J.D. (2018). Shingle 2.0: Generalising self-consistent and automated domain discretisation for multi-scale geophysical models. *Geosci. Model Dev.*, 11, 213-234. <https://doi.org/10.5194/gmd-11-213-2018>.
- Casulli, V. (1999), A semi-implicit finite difference method for nonhydrostatic free surface flows. *Internat. J. Numer. Methods Fluids*, 30(4), 425-440. [https://doi.org/10.1002/\(SICI\)1097-0363\(19990630\)30:4<425::AID-FLD847>3.0.CO;2-D](https://doi.org/10.1002/(SICI)1097-0363(19990630)30:4<425::AID-FLD847>3.0.CO;2-D).
- Casulli, V., Zanolli, P. (2002). Semi-implicit numerical modelling of non-hydrostatic free-surface flows for environmental problems. *Math. Comput. Model.*, 36, (9-10), 1131-1149. [https://doi.org/10.1016/S0895-7177\(02\)00264-9](https://doi.org/10.1016/S0895-7177(02)00264-9).
- Casulli, V., Zanolli, P. (2005). High resolution methods for multidimensional advection diffusion problems in free-surface hydrodynamics. *Ocean. Model.*, 10 (1-2), 137-151. <https://doi.org/10.1016/j.ocemod.2004.06.007>.
- Chan-Hoo Jeon, Maarten C. Buijsman, Alan J. Wallcraft, Jay F. Shriver, Brian K. Arbic, James G. Richman, Patrick J. Hogan, Improving surface tidal accuracy through two-way nesting in a global ocean model, *Ocean Modelling*, Volume 137, 2019, Pages 98-113, ISSN 1463-5003, <https://doi.org/10.1016/j.ocemod.2019.03.007>.
- Chen, W., Jacob, B., Valle-Levinson, A., Stanev, E., Staneva, J., & Badewien, T.H. (2023): Subtidal secondary circulation induced by eddy viscosity-velocity shear covariance in a predominantly well-mixed tidal inlet. *Front. Mar. Sci.*, 10:1105626. doi:10.3389/fmars.2023.1105626
- Chen, C., Liu, H., Beardsley, R.C. (2003). An unstructured grid, finite-volume, three dimensional, primitive equations ocean model: Application to coastal ocean and estuaries. *J. Atmos. Ocean. Technol.*, 20, 159-186. [https://doi.org/10.1175/1520-0426\(2003\)020<0159:AUGFVT>2.0.CO;2](https://doi.org/10.1175/1520-0426(2003)020<0159:AUGFVT>2.0.CO;2).
- Davidson, F., Alvera-Azcárate, A., Barth, A., Brassington, G.B., Chassignet, E.P., Clementi, E., De Mey-Frémaux, P., Divakaran, P., Harris, C., Hernandez, F., Hogan, P., Hole, L.R., Holt, J., Liu, G., Lu, Y., Lorente, P., Maksymczuk, J., Martin, M., Mehra, A., Melsom, A., Mo, H., Moore, A., Oddo, P., Pascual, A., Pequignet, A.-C., Kourafalou, V., Ryan, A., Siddorn, J., Smith, G., Spindler, D., Spindler, T., Stanev, E.V., Staneva, J., Storto, A., Tanajura, C., Vinayachandran, P.N., Wan, L., Wang, H., Zhang, Y., Zhu, X., and Zu, Z. (2019). Synergies in Operational Oceanography: The Intrinsic Need for Sustained Ocean Observations. *Front. Mar. Sci.* 6:450. <https://doi.org/10.3389/fmars.2019.00450>.

535 Debreu, L., Vouland, C., Blayo, E. (2008). AGRIF: Adaptive grid refinement in Fortran. *Comput. Geosci.*, 34(1), 8-13.
536 <https://doi.org/10.1016/j.cageo.2007.01.009>.

537 Debreu, L., Marchesiello, P., Penven, P., and Cambon, G. (2012). Two-way nesting in split-explicit ocean models: Algorithms,
538 implementation and validation. *Ocean Modelling*, 49-50, 1-21. <https://doi.org/10.1016/j.ocemod.2012.03.003>

539 Delft3D-Flow User Manual. Available at https://content.oss.deltares.nl/delft3d4/Delft3D-FLOW_User_Manual.pdf (last
540 access: 25/07/2024).

541 [De Mey, P., Oke, P. R., & Cummings, J. A. \(2017\). Assimilation of ocean and coastal observations: Report from the](#)
542 [International Workshop.Ocean Science, 13\(3\), 441-469. DOI: 10.5194/os-13-441-2017](#)

543 De Mey-Frémaux, P., Ayoub, N., Barth, A., Brewin, R., Charria, G., Campuzano, F., Ciavatta, S., Cirano, M., Edwards, C.A.,
544 Federico, I., Gao, S., Garcia Hermosa, I., Garcia Sotillo, M., Hewitt, H., Hole, L.R., Holt, J., King, R., Kourafalou, V., Lu, Y.,
545 Moure, B., Pascual, A., Staneva, J., Stanev, E.V., Wang, H. and Zhu, X.(2019). Model-Observations Synergy in the Coastal
546 Ocean. *Front. Mar. Sci.*, 6:436. doi: <https://doi.org/10.3389/fmars.2019.00436>

547 [Engwirda, D. \(2017\). JIGSAW: A mesh generator for geophysical modelling. Geoscientific Model Development, 10\(6\), 2117–](#)
548 [2140. https://doi.org/10.5194/gmd-10-2117-2017](#)

549 Federico, I., Pinardi, N., Coppini, G., Oddo, P., Lecci, R., and Mossa, M. (2017). Coastal ocean forecasting with an
550 unstructured grid model in the southern Adriatic and northern Ionian seas. *Nat. Hazards Earth Syst. Sci.*, 17, 45-59.
551 <https://doi.org/10.5194/nhess-17-45-2017>.

552 Ferrarin, C., Bellafore, D., Sannino, G., Bajo, M., and Umgiesser, G. (2018). Tidal dynamics in the inter-connected
553 Mediterranean, Marmara, Black and Azov seas. *Prog. Oceanogr.*, 161, 102-115. <https://doi.org/10.1016/j.pocean.2018.02.006>.

554 [FOCCUS \(Forecasting and Observing the Open-to-Coastal Ocean for Copernicus Users\) Project. https://foccus-project.eu/\).](#)
555 [Grant agreement ID: 101133911, DOI: 10.3030/101133911](#)

556 Fringer, O.B., Dawson, C. N., He, R., Ralston, D. K., Zhang, Y. J.(2019). The future of coastal and estuarine
557 ~~modeling~~modelling: Findings from a workshop. *Ocean Modelling*, 143, 101458,
558 <https://doi.org/10.1016/j.ocemod.2019.101458>

559 Fringer, O.B., Gerritsen, M., Street, R.L. (2006). An unstructured-grid, finite-volume, nonhydrostatic, parallel coastal ocean
560 simulator. *Ocean Model.*, 14 (3-4), 139-173. <https://doi.org/10.1016/j.ocemod.2006.03.006>.

561 [Frishfelds, V., She, J. & Murawski, J \(2025\). On-demand coastal modelling with two-way nesting. Ocean Dynamics 75, 23.](#)
562 [https://doi.org/10.1007/s10236-025-01670-x](#)

563 Fujii, Y., Remy, E., Zuo, H., Oke, P. R., Halliwell, G. R., Gasparin, F., et al. (2019). Observing system evaluation based on
564 ocean data assimilation and prediction systems: on-going challenges and future vision for designing/supporting ocean
565 observational networks. *Front. Mar. Sci.*, 6:417. <https://doi.org/10.3389/fmars.2019.00417>.

566 [García-León, M., Sotillo, M. G., Mestres, M., Espino, M., & Fanjul, E. Á. \(2022\). Improving Operational Ocean Models for](#)
567 [the Spanish Port Authorities: Assessment of the SAMOA Coastal Forecasting Service Upgrades. Journal of Marine Science](#)
568 [and Engineering, 10\(2\), 149. https://doi.org/10.3390/jmse10020149.](#)

569 Haid, V., Stanev, E.V., Pein, J., Staneva, J., and Chen, W. (2020). Secondary circulation in shallow ocean straits: Observations
570 and numerical ~~modeling~~modelling of the Danish Straits. *Ocean Modelling*, 148,
571 <https://doi.org/10.1016/j.ocemod.2020.101585>.

572 Harrigan, S., Zsoter, E., Alfieri, L., Prudhomme, C., Salamon, P., Wetterhall, F., Barnard, C., Cloke, H., and Pappenberger,
573 F.: GloFAS-ERA5 operational global river discharge reanalysis 1979–present, *Earth Syst. Sci. Data*, 12, 2043–2060,
574 <https://doi.org/10.5194/essd-12-2043-2020>, 2020.

575 Herzfeld, M. (2015). Methods for freshwater riverine input into regional ocean models. *Ocean Modelling*, 90, 1–15.
576 <https://doi.org/10.1016/j.ocemod.2015.04.001>

577 Herzfeld, M., Engwirda, D., & Rizwi, F. (2020). A coastal unstructured model using Voronoi meshes and C-grid staggering.
578 *Ocean Modelling*, 148, 101599. DOI: 10.1016/j.ocemod.2020.101599

579 Herzfeld M. and F. Rizwi, (2019), A two-way nesting framework for ocean models, *Environmental Modelling & Software*,
580 Volume 117, 2019, Pages 200-213, ISSN 1364-8152, <https://doi.org/10.1016/j.envsoft.2019.03.015>.

581 Holt, J., Allen, J. I., Proctor, R., & Gilbert, F. (2012). Error propagation in a North Sea tidal model: Implications for data
582 assimilation. *Continental Shelf Research*, 30(17), 2063-2071. DOI: 10.1016/j.csr.2010.10.003.

583 Hyder, K., Rossberg, A. G., Allen, J. I., Austen, M. C., Barciela, R. M., Bannister, H. J., et al. (2015). Making
584 ~~modeling~~modelling count - increasing the contribution of shelf-seas community and ecosystem models to policy development
585 and management. *Mar. Policy*, 62, 291-302. <https://doi.org/10.1016/j.marpol.2015.07.015>.

586 Ilicak, M., Federico, I., Barletta, I., Mutlu, S., Karan, H., Ciliberti, S. A., Clementi, E., Coppini, G., & Pinardi, N. (2021).
587 Modeling of the Turkish Strait System Using a High Resolution Unstructured Grid Ocean Circulation Model. *Journal of Marine*
588 *Science and Engineering*, 9(7), 769. <https://doi.org/10.3390/jmse9070769>

589 Kourafalou, V. H., De Mey, P., Le Henaff, M., Charria, G., Edwards, C. A., He, R., et al. (2015a). Coastal Ocean Forecasting:
590 system integration and evaluation. *J. Operat. Oceanogr.*, 8, S127–S146. <https://doi.org/10.1080/1755876X.2015.1022336>.

591 Kourafalou, V. H., De Mey, P., Staneva, J., Ayoub, N., Barth, A., Chao, Y., et al. (2015b). Coastal ocean forecasting: science
592 foundation and user benefits. *J. Operat. Oceanogr.*, 8, 147-167. <https://doi.org/10.1080/1755876X.2015.1022348>.

593 Kurihara, Y., Murakami, H., & Kachi, M. (2016). Sea surface temperature from the new Japanese geostationary meteorological
594 Himawari-8 satellite. *Geophysical Research Letters*, 43(3), 1234–1240. <https://doi.org/10.1002/2015GL067159>

595 Lorente, P., Piedracoba, S., Sotillo, M. G., Aznar, R., Amo-Baladrón, A., Pascual, A., et al. (2016). Characterizing the surface
596 circulation in Ebro Delta (NW Mediterranean) with HF radar and modeled current data. *J. Mar. Syst.*, 163, 61-79.
597 <https://doi.org/10.1016/j.jmarsys.2016.07.001>.

598 Luettich, Jr., R.A., Westerink, J.J., Scheffner, N.W. (1992). ADCIRC: an Advanced Three Dimensional Circulation Model for
599 Shelves, Coasts and Estuaries. Report 1: Theory and methodology of ADCIRC-2DDI and ADCIRC-3DL. Dredging Research
600 Program Technical Report DRP-92-6, U.S. Army Engineers Waterways Experiment Station, Vicksburg, MS, 137 pp. URI:
601 <http://hdl.handle.net/11681/4618>.

602 Madec, G. & the NEMO Team (2016). Title: NEMO ocean engine. Institution: Institut Pierre-Simon Laplace (IPSL), France.

Version: NEMO v3.6 DOI: <https://doi.org/10.5281/zenodo.1464816>

Madec, Gurvan, and The NEMO System Team. NEMO Ocean Engine Reference Manual. Zenodo, 2024. <https://doi.org/10.5281/zenodo.6334656>.

Maicu, F., De Pascalis, F., Ferrarin, C., and Umgiesser, G. (2018). Hydrodynamics of the Po River-Delta-Sea system. *J. Phys. Res. Oceans*, 123, 6349-6372. <https://doi.org/10.1029/2017JC013601>.

Matte, P., Wilkin, J., & Staneva, J. (2024). *The Role of Rivers in Ocean Forecasting*. Preprint. <https://doi.org/10.5194/sp-2024-9>

Marchesiello P., Auclair, F., Debreu, L, McWilliams, J.C., Almar, R., Benshila, R., Dumas, F. (2021). Tridimensional nonhydrostatic transient rip currents in a wave-resolving model. *Ocean Modelling*, 163, 101816. <https://doi.org/10.1016/j.ocemod.2021.101816>.

Marchesiello, P., McWilliams, J. C., & Shchepetkin, A. (2001). Open boundary conditions for long-term integration of regional oceanic models. *Ocean Modelling*, 3(1–2), 1–20. [https://doi.org/10.1016/S1463-5003\(00\)00013-5](https://doi.org/10.1016/S1463-5003(00)00013-5)

Mani, S., Calzada, J. R., Moghimi, S., Melton, C., & Signell, R. P. (2021). OCSMesh: a data-driven automated unstructured mesh generation software for coastal ocean modeling. NOAA Technical Memorandum NOS CS; 47. U.S. Department of Commerce. <https://repository.library.noaa.gov/view/noaa/33879>

Marshall, J., Adcroft, A., Hill, C., Perelman, L., Heisey, C. (1997). A finite volume, incompressible Navier–Stokes model for studies of the ocean on parallel computers. *J. Geophys. Res.* 102(C3), 5753-5766. <https://doi.org/10.1029/96JC02775>.

Mellor, G. L. (2004). *Users guide for a three-dimensional, primitive equation, numerical ocean model*. Institution: Princeton University, Program in Atmospheric and Oceanic Sciences. URL: http://jes.apl.washington.edu/modsims_two/usersguide0604.pdf

Mengual, B., Le Hir, P., Cayocca, F., Garlan, T. (2017). Modelling fine sediment dynamics: Towards a common erosion law for fine sand, mud and mixtures. *Water*, 9(8), 564. <https://doi.org/10.3390/w9080564>.

McWilliams, J.C. (2016). Submesoscale Currents in the Ocean. *Proceedings of the Royal Society A. Mathematical, Physical and Engineering Sciences*, 472(2189), 20160117. <http://doi.org/10.1098/rspa.2016.0117>.

Moore, A. M., and Martin, M. J. (2019). Synthesis of ocean observations using data assimilation: toward a more complete picture of the State of the Ocean. *Front. Mar. Sci.* 6:90. <https://doi.org/10.3389/fmars.2019.00090>.

Mourre, B., & Chiggiato, J. (2014). A comparison of ensemble-based data assimilation methods for the North Sea. *Journal of Marine Systems*, 129, 190-203. DOI: 10.1016/j.jmarsys.2013.05.009

Munk, W., Wunsch, C. (1998). Abyssal recipes II, Energetics of tidal and wind mixing. *Deep-Sea Res., Part I*, 45, 1977-2010. [https://doi.org/10.1016/S0967-0637\(98\)00070-3](https://doi.org/10.1016/S0967-0637(98)00070-3).

NOAA (2020). 2020 National Ocean Service Science Report. National Oceanic and Atmospheric Administration. Retrieved from: <https://repository.library.noaa.gov/view/noaa/33879>

Nguyen, T. T., Staneva, J., Grayek, S., Bonaduce, A., Hagemann, S., Pham, N. T., Kumar, R., & Rakovec, O. (2024). Impacts of extreme river discharge on coastal dynamics and environment: Insights from high-resolution modeling in the German Bight. *Regional Studies in Marine Science*, 66, 103476. <https://doi.org/10.1016/j.rsma.2024.103476>

Oke, P. R., Allen, J. S., Miller, R. N., Egbert, G. D., and Kosro, P. M. (2002). Assimilation of surface velocity data into a primitive equation coastal ocean model. *J. Geophys. Res.* 107:3122. <https://doi.org/10.1029/2000JC000511>.

Oke, P. R., and Sakov, P. (2012). Assessing the footprint of a regional ocean observing system. *J. Mar. Syst.*, 105, 30-51. <https://doi.org/10.1016/j.jmarsys.2012.05.009>.

Oliveira, A., Fortunato, A. B., Rogeiro, J., Teixeira, J., Azevedo, A., Lavaud, L., Bertin, X., Gomes, J., David, M., Pina, J., Rodrigues, M., & Lopes, P. (2019). OPENCoastS: An open-access service for the automatic generation of coastal forecast systems. *Environmental Modelling & Software*, 124, 104585. <https://doi.org/10.1016/j.envsoft.2019.104585>

A. Oliveira, A.B. Fortunato, M. Rodrigues, A. Azevedo, J. Rogeiro, S. Bernardo, L. Lavaud, X. Bertin, A. Nahon, G. Jesus, M. Rocha, P. Lopes, 2021. Forecasting contrasting coastal and estuarine hydrodynamics with OPENCoastS, *Environmental Modelling & Software*, Volume 143, 105132, ISSN 1364-8152, <https://doi.org/10.1016/j.envsoft.2021.105132>.

Pein, J.U., Grayek, S., Schulz-Stelleneth, J., Stanev, E.V. (2016). On the impact of salinity observations on state estimates in Ems Estuary. *Ocean Dynamics*, 66, 243-262. <https://doi.org/10.1007/s10236-015-0920-0>.

Pein, J., Staneva, J., Daewel, U., & Schrum, C. (2021): Channel curvature improves water quality and nutrient filtering in an artificially deepened mesotidal idealized estuary. *Continental Shelf Research*, Volume 231, 104582, [doi:10.1016/j.csr.2021.104582](https://doi.org/10.1016/j.csr.2021.104582)

Ponte, R. M., Carson, M., Cirano, M., Domingues, C., Jevrejeva, S., Marcos, M., et al. (2019). Towards comprehensive observing and modeling systems for monitoring and predicting regional to coastal sea level. *Front. Mar. Sci.*, 6. <https://doi.org/10.3389/fmars.2019.00437>.

Rao, Y.R., Murthy, C.R., Sinha, P.C (2008). The coastal ocean. In: Murthy, C.R., Sinha, P.C., Rao, Y.R. (Eds.), *Modelling and Monitoring of Coastal Marine Processes*. Springer (Dordrecht) & Capital Publishing Company (New Delhi), pp. 3-10. <https://doi.org/10.1007/978-1-4020-8327-3>

Reyes, E., Aguiar, E., Bendoni, M., Berta, M., Brandini, C., Cáceres-Euse, A., Capodici, F., Cardin, V., Cianelli, D., Ciraolo, G., Corgnati, L., Dadić, V., Doronzo, B., Drago, A., Dumas, D., Falco, P., Fattorini, M., Fernandes, M. J., Gauci, A., Gómez, R., Griffa, A., Guérin, C.-A., Hernández-Carrasco, I., Hernández-Lasheras, J., Ličer, M., Lorente, P., Magaldi, M. G., Mantovani, C., Mihanović, H., Molcard, A., Mourre, B., Révelard, A., Reyes-Suárez, C., Saviano, S., Sciascia, R., Taddei, S., Tintoré, J., Toledo, Y., Uttieri, M., Vilibić, I., Zambianchi, E., and Orfila, A.: Coastal high-frequency radars in the Mediterranean – Part 2: Applications in support of science priorities and societal needs, *Ocean Sci.*, 18, 797–837, <https://doi.org/10.5194/os-18-797-2022>, 2022.

Ringler, T., Petersen, M., Higdon, R. L., Jacobsen, D., Jones, P. W., & Maltrud, M. (2013). A multi-resolution approach to global ocean modeling. *Ocean Modelling*, 69, 211–232. DOI: 10.1016/j.ocemod.2013.04.010

Rudnick, D. L. (2016). *Ocean*

research enabled by underwater gliders. *Annual Review of Marine Science*, 8, 519–541. <https://doi.org/10.1146/annurev-marine-122414-033913>

Roberts, K. J., Pringle, W. J., Mattocks, C. A., & Westerink, J. J. (2019). OceanMesh2D 1.0: MATLAB-based software for two-dimensional unstructured mesh generation in coastal ocean modeling. *Geoscientific Model Development*, 12(5), 1847–1868. <https://doi.org/10.5194/gmd-12-1847-2019>

Tonani, M., Sykes, P., King, R. R., McConnell, N., Péquignot, A.-C., O'Dea, E., Graham, J. A., Polton, J., and Siddorn, J. (2019). The impact of a new high-resolution ocean model on the Met Office North-West European Shelf forecasting system. *Ocean Sci.*, 15, 1133–1158, <https://doi.org/10.5194/os-15-1133-2019>.

Rubio, A. J., Mader, L., Corgnati, C., Mantovani, A., Griffa, A., Novellino, C., et al. (2017). HF radar activity in European coastal seas: next steps towards a pan-european hf radar network. *Front. Mar. Sci.*, 4:8. <https://doi.org/10.3389/fmars.2017.00008>.

van Sebille, E., England, M. H., and Froyland, G. (2012). Origin, dynamics and evolution of ocean garbage patches from observed surface drifters. *Environ. Res. Lett.*, 7, 1026-1034. <https://dx.doi.org/10.1088/1748-9326/7/4/044040>

Shulman, I., and Paduan, J. D. (2009). Assimilation of HF radar-derived radials and total currents in the Monterey Bay area. *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56, 149-160. <https://doi.org/10.1016/j.dsr2.2008.08.004>.

Schulz-Stellenfleh, J., and Stanev, E. V. (2016). Analysis of the upscaling problem - a case study for the barotropic dynamics in the North Sea and the German Bight. *Ocean Model.*, 100, 109-124. <https://doi.org/10.1016/j.ocemod.2016.02.002>.

Siddorn, J. R., Good, S. A., Harris, C. M., Lewis, H. W., Maksymczuk, J., Martin, M. J., and Saulter, A. (2016). Research priorities in support of ocean monitoring and forecasting at the Met Office. *Ocean Sci.*, 12, 217–231, <https://doi.org/10.5194/os-12-217-2016>

Shchepetkin, A., McWilliams, J.C. (2005). The regional oceanic ~~modeling~~modelling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model. *Ocean Model.* 9(4), 347-404. <https://doi.org/10.1016/j.ocemod.2004.08.002>.

Sobrinho, J., de Pablo, H., Campuzano, F., & Neves, R. (2021). Coupling Rivers and Estuaries with an Ocean Model. *Water*, 13(16), 2284, DOI: 10.3390/w13162284.

Stanev, E. V., Al-Nadhairi, R., Valle-Levinson, A., & Badewien, T. H. (2011). Tidal dynamics and residence time in the German Bight: Linking transport and variability. *Journal of Geophysical Research: Oceans*, 116(C8). DOI: 10.1029/2010JC006835

Stanev, E. V., Grashorn, S., and Zhang, Y. J. (2017). Cascading ocean basins: numerical simulations of the circulation and interbasin exchange in the Azov-Black-Marmara-Mediterranean Seas system. *Ocean Dyn.*, 67, 1003-1025. <https://doi.org/10.1007/s10236-017-1071-2>

Stanev, E., Schulz-Stellenfleh, J., Staneva, J., Grayek, S., Grashorn, S., Behrens, A., et al. (2016). Ocean forecasting for the German Bight: from regional to coastal scales. *Ocean Sci.*, 12, 1105-1136. <https://doi.org/10.5194/os-12-1105-2016>,

701 Stanev, E. V., Ziemer, F., Schulz-Stellenfleth, J., Seemann, J., Staneva, J., and Gurgel, K. W. (2015). Blending surface currents
702 from HF radar observations and numerical modelling: tidal hindcasts and forecasts. *J. Atmos. Oceanic Technol.* 32, 256–281.
703 <https://doi.org/10.5194/os-12-1105-2016>.

704 Staneva, J., Alari, V., Breivik, O., Bidlot, J.-R., and Mogensen, K. (2017). Effects of wave-induced forcing on a circulation
705 model of the North Sea. *Ocean Dyn.*, 67, 81-101. <https://doi.org/10.1007/s10236-016-1009-0>.

706 Staneva, J., Wahle, K., Günther, H., and Stanev, E. (2016a). Coupling of wave and circulation models in coastal-ocean
707 predicting systems: a case study for the German Bight. *Ocean Sci.*, 12, 797-806. <https://doi.org/10.5194/os-12-797-2016>.

708 Staneva, J., Wahle, K., Koch, W., Behrens, A., Fenoglio-Marc, L., and Stanev, E. (2016b). Coastal flooding: impact of waves
709 on storm surge during extremes – a case study for the German Bight. *Nat. Hazards Earth Syst. Sci.* 16, 2373–2389.
710 <https://doi.org/10.5194/nhess-16-2373-2016>.

711 [Sotillo, M. G., Campuzano, F., Guihou, K., Lorente, P., Olmedo, E., Matulka, A., Santos, F., Amo-Baladrón, M. A., &
712 Novellino, A. \(2021\). River Freshwater Contribution in Operational Ocean Models along the European Atlantic Façade: Impact
713 of a New River Discharge Forcing Data on the CMEMS IBI Regional Model Solution. *Journal of Marine Science and
714 Engineering*, 9\(4\), 401. <https://doi.org/10.3390/jmse9040401>](#)

715 [Sotillo, M. G. \(2022\). Ocean Modelling in Support of Operational Ocean and Coastal Services. *Journal of Marine Science and
716 Engineering*, 10\(10\), 1482. <https://doi.org/10.3390/jmse10101482>](#)

717 [Sotillo MG, Mourre B, Mestres M, Lorente P, Aznar R, García-León M, Liste M, Santana A, Espino M and Álvarez E \(2021\)
718 Evaluation of the Operational CMEMS and Coastal Downstream Ocean Forecasting Services During the Storm Gloria
719 \(January 2020\). *Front. Mar. Sci.* 8:644525. doi: 10.3389/fmars.2021.644525](#)

720 [Sun, Q., Whitney, M.M., Bryan, F.O., & Tseng, Y. \(2017\). A box model for representing estuarine physical processes in Earth
721 system models. *Ocean Modelling*, 112, 139–153. <https://doi.org/10.1016/j.ocemod.2017.03.004>](#)

722 [Testor, P., de Young, B., Rudnick, D. L., et al. \(2019\). OceanGliders: A component of the integrated GOOS. *Frontiers in
723 Marine Science*, 6, 422. <https://doi.org/10.3389/fmars.2019.00422>](#)

724 Trotta, F., Pinardi, N., Fenu, E., Grandi, A., and Lyubartsev, V. (2017). Multinest high-resolution model of submesoscale
725 circulation features in the Gulf of Taranto. *Ocean Dyn.*, 67, 1609-1625. <https://doi.org/10.1007/s10236-017-1110-z>.

726 Umgiesser, G., Canu, D.M., Cucco, A., Solidoro, C. (2004). A finite element model for the Venice Lagoon. Development, set
727 up, calibration and validation. *Journal of Marine Systems* 51(1-4), 123–145. <https://doi.org/10.1016/j.jmarsys.2004.05.009>

728 [Sotillo, M. G., Campuzano, F., Guihou, K., Lorente, P., Olmedo, E., Matulka, A., Santos, F., Amo-Baladrón, M. A., &
729 Novellino, A. \(2021\). River Freshwater Contribution in Operational Ocean Models along the European Atlantic Façade: Impact
730 of a New River Discharge Forcing Data on the CMEMS IBI Regional Model Solution. *Journal of Marine Science and
731 Engineering*, 9\(4\), 401. <https://doi.org/10.3390/jmse9040401>](#)

732 [Verri, G., et al. \(2020\). Box model approaches to estuarine dynamics in coarse-resolution ocean models. *Ocean Modelling*,
733 148, 101587. <https://doi.org/10.1016/j.ocemod.2020.101587>](#)

Wahle, K., Staneva, J, Koch, W., Fenoglio-Marc, L., Ho-Hagemann, H. T. M., and Stanev, E.V. (2017). An atmosphere-wave regional coupled model: improving predictions of wave heights and surface winds in the Southern North Sea. *Ocean Sci.*, 13, 289–301. <https://doi.org/10.5194/os-13-289-2017>.

Warner, J.C., Armstrong, B., He, R., J. B. Zambon, J.B. (2010). Development of a Coupled Ocean-Atmosphere-Wave-Sediment Transport (COAWST) ~~modeling~~modelling system. *Ocean Model.*, 35(3), 230-244. <https://doi.org/10.1016/j.ocemod.2010.07.010>.

Warner, J.C., Sherwood, C.R., Signell, R.P., Harris, C., Arango, H.G. (2008). Development of a three-dimensional, regional, coupled wave, current, and sediment-transport model. *Comput. Geosci.* 34(10), 1284-1306. <https://doi.org/10.1016/j.cageo.2008.02.012>.

Westerink, J.J., Luetich, R.A., Blain, C.A., Scheffner, N.W. (1994). ADCIRC: An Advanced Three-Dimensional Circulation Model for Shelves, Coasts and Estuaries. Report 2: Users' Manual for ADCIRC-2DDI, Department of the Army US Army Corps of Engineers, Washington, D.C. Available at <https://apps.dtic.mil/sti/citations/ADA276150> (last access: 25/07/2024).

Wilkin, J. L., Arango, H. G., Haidvogel, D. B., Lichtenwalner, C. S., Glenn, S. M., and Hedström, K. S. (2005). A regional ocean ~~modeling~~modelling system for the Long-term Ecosystem Observatory. *J. of Geophy. Res.: Oceans*, 110(C6), <https://doi.org/10.1029/2003JC002218>.

Zeng, X., Atlas, R., Birk, R. J., Carr, F. H., & others (2020). Title: Use of observing system simulation experiments in the United States. Journal: Bulletin of the American Meteorological Society, 101(8). doi: 10.1175/BAMS-D-19-0155.1

Zhang, Y. and Baptista, A. M. (2008). SELFE: a semi-implicit Eulerian– Lagrangian finite-element model for cross-scale ocean circulation. *Ocean Model.*, 21(3-4), 71-96. <https://doi.org/10.1016/j.ocemod.2007.11.005>.

Zhang, Y. J., Stanev, E., and Grashorn, S. (2016a). Unstructured-grid model for the North Sea and Baltic Sea: validation against observations. *Ocean Model.*, 97, 91-108. <https://doi.org/10.1016/j.ocemod.2015.11.009>.

Zhang, Y. J., Ye, F., Stanev, E. V., and Grashorn, S. (2016b). Seamless cross-scale modelling with SCHISM. *Ocean Model.*, 102, 64-81. <https://doi.org/10.1016/j.ocemod.2016.05.002>.

758 **Competing interests**

759 The contact author has declared that none of the authors has any competing interests.

761 **Author contributions**

762 JS conceptualized the study, analyzed data, and wrote this article. All authors contributed to the writing of the article and
763 quality control.

765 **Competing interests**

766 The contact author has declared that neither of the authors has any competing interests.

767

768 **Disclaimer**

769 Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European
770 Union or the European Health and Digital Executive Agency (HaDEA). Neither the European Union nor HaDEA can be held
771 responsible for them.

772

773 Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published
774 maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes
775 every effort to include appropriate place names, the final responsibility lies with the authors.

776

777 Acknowledgements

778 JS and AM acknowledge —Horizon Europe Project FOCCUS “Forecasting and observing the open-to-coastal ocean for
779 Copernicus users” (Grant Agreement 101133911).

780