The <u>Role-Representation</u> of Rivers in <u>Operational</u> Ocean Forecasting <u>Systems: A Review</u>

Pascal Matte¹, John Wilkin², Joanna Staneva³

¹Meteorological Research Division, Environment and Climate Change Canada, Québec, QC, Canada

⁵ ²Department of Marine and Coastal Sciences, Rutgers, The State University of New Jersey, New Brunswick, NJ, USA ³Institute of Coastal Systems - Analysis and Modeling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

Correspondence to: Pascal Matte (pascal.matte@ec.gc.ca)

10

Abstract. The connection between the ocean and the land is made possible thanks to rivers, which are a vital component of the Earth's system. They govern the hydrological and biogeochemical contributions to the coastal ocean through surface and subsurface water discharge and influence local circulation and the distribution of water masses, modulating processes such as upwelling and mixing. This paper provides an overview of recent advancements in river modellingapproaches to representing

- 15 coastal river discharges and processes in operational ocean forecasting systems (OOFS), with a particular focus on estuaries. The methods discussed range frominclude those currently adopted in coarse-resolution ocean forecasting systems, where mixing processes are primarily parameterized, toas well as more advanced modelling and -coupling approaches tailored to high-resolution that are better suited for-coastal systems. A review of river data availability is also presented, illustrating various sources of freshwater discharge and salinity, from observational data to climatological datasets, and more
- 20 precisealongside river operational river discharge products modelling approaches that are improvingenhance the representation of water discharges in operational systems. New satellite-derived datasets and emerging river modelling techniques are also introduced. FinallyIn addition, a compendium of currentresponses from a survey of existing operational systemsOOFS providers is provided are synthetized, with a focus on how river forcing is treated, from global to coastal scales. Challenges such as data accuracy, standardization, and model coupling are discussed, highlighting the need for improved interfaces
- 25 <u>between monitoring and modelling systems. Finally, some recommendations and ways forward are formulated in relation to</u> <u>identified limitations in current OOFS.</u>

1 Introduction

Rivers provide form the primary link between land and sea, and delivering annually an average of approximately 36,000 thousand km³ of freshwater and over 20 billion tons of solid and dissolved material to the world-global ocean each year

30 (Milliman and Farnsworth, 2011). River discharge into the ocean is a major component of the global hydrological and biogeochemical cycles, which have undergone significant changes under the influence of climate change and human activities (Shi et al., 2019; Yan et al., 2022; Qin et al., 2022; Chandanpurkar et al., 2022). Mediated through estuaries, Estuaries act as

transitional zones where freshwater fluxes influence the ocean circulation and, salinity, and in particular the upper-ocean stratification, which in turn affects the mixed layer depth, ocean currents, and air-sea interaction (Chandanpurkar et al., 2022;

35 Dzwonkowski et al., 2017; Sprintall and Tomczak, 1992; Sun et al., 2017<u>: Pein et al., 2021, 2024</u>). Freshwater inputs to the ocean also modulate coastal upwelling events<u>. Altogether</u>, thus<u>these factors</u> impacting productivity of the coastal marine environment (Sotillo et al., 2021<u>a</u>).

Despite rivers' influence on the coastal and basin-wide circulation and dynamics, in global and regional scale models, effectively accounting for riverine freshwater discharge into the oceans is a challenging problem (Sun et al., 2017; Verri et al.,

- 40 2020). Accurately incorporating river flow into numerical ocean models requires appropriate parameterizations and boundary conditions. The setup of practical open boundary conditions (OBC) is dependent on flow dynamics, model resolution, data availability, and other factors (Blayo and Debreu, 2005). At coarse scales that cannot resolve the estuarine dynamics, but even at finer scales in some cases, river outlets are typically-often represented in a simplistic way, with climatological runoff and zero or constant salinity values, implicitly neglecting estuarine mixing andor exchange as well as seasonal and non-seasonal
- 45 variability (Sun et al., 2017; Verri et al., 2020; Verri et al., 2021; Pein et al., 2021, 2024). ConsequentlyAs a result, important key natural processes are <u>often</u> omitted, and, depending on how river forcing is defined, ocean model <u>results outputs</u> may <u>differ</u> vary significantly from one another,. These discrepancies are most pronounced predominantly in the shelf areas, particularly in (or rRegions of freshwater Freshwater influenceInfluence, (ROFI), but <u>can</u> also propagate to at regional and global scales (Tseng et al., 2016).

50

The next sections present a This paper reviews existing methods and datasets used in Operational Ocean Forecasting Systems (OOFS) to represent river forcing. As the focus is on freshwater discharges, the river supply of nutrients and other materials are neglected in this review but are partly addressed in a separate contribution by Cossarini et al. (2024).

The paper is structured as follows: Section 2 reviews of approaches to treatfor representing river forcing in global, regional,

- 55 and coastal ocean models, including dynamic methods to represent the mixing processes in estuaries throughestuarine mixing parameterizations and coupling techniques. Section 3 A description of describes available data sources from operational centers and data providers as well as emerging techniques for estimating river dischargeused for this purpose is also provided, with eSection 4 presents examples of river forcing methods and data sources taken from implemented in existing OOFS, summarizing. While the first sections review the scientific literature and ancillary information, for example, taken from
- 60 operational centers and data providers websites, the last section summarizes results findings from a survey sent to the conducted within the OceanPredict community. Finally, Section 5 provides a summary and recommendations regarding identified limitations in current OOFS.

2 River forcing in ocean models

2.1 Capturing seasonal and non-seasonal river variability

- 65 Accurate representation of river discharges and associated variables (e.g., salinity, temperature), whether model-derived or observation-based, is crucial for capturing both seasonal and non-seasonal effects in the coastal ocean. The Bay of Bengal is one example where the inclusion of seasonal river discharges and salinity in regional model simulations significantly improves the representation of sea surface temperatures, near-surface salinity, stratification, mixed-layer depth, and barrier-layer thickness, leading to a better simulation of the formation, progression and dispersion of the freshwater plume (Jana et al.,
- 70 <u>2015).</u>

Seasonal variability in river discharge not only impacts coastal salinity and temperature but also contributes to the sea level changes both locally and remotely, mostly via a halosteric sea level contribution. This effect was observed, for example, between the mouth of the Amazon River and the continental shelves of the Gulf of Mexico and Caribbean Sea (Giffard et al., 2019). Similarly, in the U.S. Atlantic and Gulf coasts, river discharge and sea level changes were found to be significantly

75 correlated (Piecuch et al. 2018). Such dynamic sea surface height (SSH) signals driven by river discharge can explain 10-20% of the regional-scale seasonal variance around major rivers, such as the Amazon, Ganges, Brahmaputra, Irrawaddy, Ob, Lena, and Yenisei (Piecuch and Wadehra, 2020).

While the seasonal effects of river discharge on ocean processes have been extensively documented, non-seasonal influences of river runoff on sea level changes remain largely unexplored due to the lack of consolidated discharge databases (Durand et

80 <u>al., 2019</u>). These influences, however, can be significant when considering river runoff jointly with wind-driven transport and heat fluxes, which also play a major role in modulating regional sea level variability (Verri et al., 2018).

2.2 Freshwater input in coarse resolution models: towards a parameterization of estuarine mixing processes

Because many ocean models operate at resolutions too coarse to resolve estuarine processes explicitly, appropriate parameterization of estuarine mixing is required to capture their influence on freshwater transport. In nature, estuaries transport

- and transform water properties along their length, due to tidal mixing, deposition and resuspension, and up- and down-estuary advection. Saltwater intrusion driven by tides and other coastal signals (e.g. storm surges) controls the estuarine water exchange and affects the net estuarine outflow and corresponding salinity values (Sun et al., 2017; Verri et al., 2020). However, although water properties at the head differ from those at the mouth, in models too coarse to resolve the estuaries, river discharge observed far from the river outlet is typically inputted at the coast with zero salinity (Verri et al., 2021; Herzfeld, 2015).
- 90 <u>Alternatively, salinity values can be prescribed based on constant annual or monthly values derived from sensitivity tests</u> and/or in situ campaigns, when available (Verri et al., 2018).

2.1 Freshwater input in coarse resolution models: towards a parameterization of estuarine mixing processes

Herzfeld (2015) describes and assesses the performance of various methods for inputting freshwater into regional ocean

- 95 models. A first approach, referred to as a point source input, adds a term of freshwater flux, entering as surface point sources into one or more layers of the model, to the divergence of flow in via the vertically integrated continuity equation, with no associated velocity profile. It affects the vertical velocity surface boundary condition of the free surface equation, and the surface boundary conditions for the diffusive heat and salt fluxes. A second approach, the flow input, considers the inertia of the river flow and prescribes a velocity profile at the boundary whose vertical integral is equal to the inflow flux. These two
- 100 methods must have a predefined depth at the boundary over which to distribute the volume inflow. A more accurate approach is to add an artificial channel to the coastline to give momentum to the flow and initiate mixing between fresh and salt waters (Lacroix et al., 2004; Sobrinho et al., 2021).

The horizontal distribution of the runoff plays an important role in the regional salinity distribution and in the vertical stratification and mixing (Tseng et al., 2016). Additional subtleties arise for large rivers or deltas, where the coastal source

- 105 points need to be spread laterally to avoid numerical instabilities if inflow values are locally too large (Polton et al. 2023). In global ocean models, however, freshwater inflow is frequently added at the ocean surface, either as an increased precipitation rate over a specified area or by reducing surface salinity (i.e. a virtual salt flux), rather than being introduced as a lateral inflow at the coastal boundary. This freshwater can be distributed vertically over several layers or diffused horizontally using enhanced mixing (Sun et al., 2017; Tseng et al., 2016; Yin et al., 2010).
- 110 Several plume responses may result from the choice of the horizontal and vertical distribution of freshwater input. However, most model applications produce plumes whose types differ from plumes associated with real river discharges (Tseng et al., 2016; Garvine, 2001; Schiller and Kourafalou, 2010). Larger scale offshore stratification is also expected to be impacted by this choice.

MacCready and Geyer (2010) establish the theoretical foundation for estuarine mixing parameterizations, which underpins

- some physics-based methods used to simulate unresolved estuarine processes in regional and global ocean models, such as the estuary box model (EBM); see, for example, Figure 1 (Sun et al., 2017). These models attempt to parameterize mixing processes and to account for baroclinic and barotropic flow, typically using a two-layer formulation; (e.g. Verri et al., (2020;), Verri et al., (2021;), Herzfeld, (2015;), Rice et al., (2008;), Hordoir et al., (2008). From these representations, analytical solutions can be found for the volume fluxes and outflow salinity. Applied globally to the Community Earth System Model
- 120 (CESM), such an approach revealed substantial localized, regional, and long-range effects when compared to cases without parameterization, highlighting once again the strong sensitivity of ocean models to the treatment of rivers (Sun et al., 2017). New hybrid approaches, such as the Hybrid-EBM (Maglietta et al., 2025; Saccotelli et al., 2024), combine physics-based models with machine learning techniques to predict the salt-wedge intrusion length and salinity at river mouths. Hybrid-EBM outperforms the classical EBM and addresses the shortcomings of the dimensional equations in the physics-based EBM, which





Figure 1: Schematic diagram of the estuary box model (EBM) implemented in the Community Earth System Model (CESM) (Sun et al., 2017). The EBM is, depicted as a two-layer rectangular box with constant width, uniform local depth (H), and a time-varying
length (L). Each layer has a fixed thickness (h for the lower layer and H-h for the upper layer), with vertically uniform but horizontally variable salinity and density. Thick solid lines represent closed boundaries, dotted lines mark open boundaries, and the dashed line shows the interface between layers. Volume fluxes (Q) and salinities (S) are indicated by arrows at open boundaries: riverine freshwater discharge (QR) enters at the estuary head, oceanic saltwater flows into the lower layer at the mouth (QLM), and QUt represents the average tidal volume flux during half a tidal cycle, driving net horizontal salt flux into the upper layer at the mouth. Shear-induced turbulent mixing (shown by paired upward and downward open arrows) and upward advection from exchange flow (solid upward arrows) link the upper and lower layers. The color gradient illustrates salinity variation, from fresher

(lighter shades) to saltier (darker shades) waters.¹

2.32 Freshwater input in high resolution models: unstructured modelling of the river-sea continuum

In contrast, when the model resolution is higher than the estuary width, the latter can be resolved explicitly by extending the

140 grid for some distance inland using either real bathymetry or a straight channel approximation. When extending it beyond the salinity intrusion limit and/or the head of tides, a freshwater flux can be directly specified at the upstream boundary. This is the preferred option in many east coast US studies (Herzfeld, 2015) (e.g. RISE - Liu et al., 2004; LATTE - Choi and Wilkin, 2007; MerMADE - Hetland and MacDonald, 2008).

The use of unstructured grids offers various advantages, including a more accurate treatment of the freshwater inputs from

145 rivers, a realistic representation of river-sea interactions and estuarine processes at spatial and temporal scales usually not resolved in the ocean, and an improved interface between estuaries and the open ocean, sometimes with higher-order spatial discretizations (Staneva et al., 2024). In addition, the unstructured grid modelling combined with an efficient vertical coordinate system can better resolve the coastal sea dynamics (Verri et al., 2023).

With seamless grid transitions between models or domains, flexibility and cross-scale capabilities are augmented (Zhang et

150 al., 2016). As examples, a river-coastal-ocean continuum model has been developed for the Tiber River delta, reproducing the

¹ Reprinted from Ocean Modelling, Vol 112, Sun, Q., Whitney, M. M., Bryan, F. O., and Tseng, Y., A box model for representing estuarine physical processes in Earth system models, Page 140, Copyright Elsevier Ltd. (2017), with permission from Elsevier.

coastal dynamic processes better than the classic coastal-ocean representation, including the salt wedge intrusion, and revealing new features near the river mouth induced by river discharge and coastal morphology (Bonamano et al., 2024). In the Columbia River estuary, where both shelf and estuarine circulations are coupled, a multi-scale model has proved to reproduce key processes driving the river plume dynamics in a region characterized by complex bathymetry and marked

- 155 gradients in density and velocity (Vallaeys et al., 2018). Likewise, Vallaeys et al. (2021) used a similar model in a topographically challenging area of the Congo River estuary, characterized by high river discharge, strong stratification and large depth. Similarly, Maicu et al. (2021) simulated the circulation in the Goro Lagoon and Po River Delta branches using downscaling and a seamless chain of models integrating local forcings and dynamics into a coarser OOFS based on a cascading approach.
- 160 While these examples were successful in representing dynamical processes across temporal and spatial scales, in some contexts, the large inward tidal extent and/or complex bathymetries and coastlines, often featuring coastal infrastructures, pose significant challenges for explicitly resolving estuaries, making it impractical in many coastal models. As a result, this approach has yet to become standard practice in OOFS.

165 2.43 One-way and two-way coupling

Coupling techniques can be used to link two or more models to allow one-way data exchange, for example, between a hydrological model and an ocean model. That wayIn this approach, external forcing is reduced to fewer-a limited set of variables, simplifying computational requirements but potentially overlooking key-Limitations of this approach include processes that cannot be resolved at the land-sea interface. Additionally, it requires and the need to extending the ocean domain 170 limits boundaries far landwardinland, beyond the limit of tide and storm-surge propagation. While some parameterizations (cf. Section 2.2) or use of unstructured grids (cf. Section 2.3) can partly alleviate these shortcomings, Iin a compound flooding context, two-way coupled models are preferred because both land and ocean processes can be represented along with their interactions (Bao et al., 2022; Cheng et al., 2010). When adding The inclusion of momentum flux exchanges between land and ocean processes, the ability to reproduce improves the simulation of estuarine water levels in the estuary is enhanced by 175 capturing nonlinear feedbacks between runoff and residual ocean water levels. In a case study of Hurricane Florence, (Bao et al., (2022) achieved significant improvement in simulated water levels (20%-40% at the head of Cape Fear River Estuary) during the post-hurricane period by using a two-way coupled model, compared to a stand-alone and linked (one-way coupled) approach. - Moreover, with seamless grid transitions between the different models, flexibility and cross scale capabilities are augmented (Zhang et al., 2016).

180 Alternative approaches for assessing the risk of compound flooding have been proposed, including integrated hydrodynamic and machine learning methods to predict water level dynamics (Sampurno et al., 2022). Such approaches are particularly valuable in data-scarce regions, where developing fully calibrated, computationally intensive models can be impractical or infeasible.

185 **3 Data sources**

190

3.1 Freshwater discharge

An unsolved problem and one of the classical limitations of <u>A persistent challenge in</u> OOFS with respect to river forcing is the absence-lack of a global networks of for observed river flows to the oceans. While advances are being made in creating such a network, several challenges remain pertaining to data quality, accessibility, and timeliness, at the required spatial and temporal scales.

In situ river discharge observations are necessary to build climatologies<u>and t</u>hey represent a key component of the calibration of hydrological models, and thereby of any reanalysis, near-real-time (NRT) analysis and forecast products. The various types of discharge products used in OOFS are described in the following.

3.1.1 Climatologies

- 195 Most ocean models use climatologies to introduce river forcing based on multi-decadal averages of observed and/or modeled freshwater discharges, along with zero or constant salinity values. Although <u>use of</u>-climatological data is <u>still</u>-commonly acceptedused, even <u>when in cases wherethe</u> estuarine dynamics <u>is are not explicitly</u> resolved, more realistic <u>and less subjective</u> estimates of freshwater volume flux and salinity <u>inputs estimates</u> would <u>produce a more accurate representationimprove the</u> modelling of <u>coastal (e.g. river plumes) to basin-wide circulation and dynamics (e.g. dense water formation, overturning</u>
- <u>circulation cells, water exchange at straits</u>) in ocean models (Verri et al., 20<u>1821</u>), especially during non_seasonal (e.g. storm induced) events (Chandanpurkar et al., 2022). Moreover, given the global decline of the hydrometric networks, building climatologies is not always possible, especially for small or less_-studied rivers, and even for large rivers in regions where routine monitoring is absent (Campuzano et al., 2016; Mishra and Coulibaly, 2009). Furthermore, monthly climatological products are not adequate for high resolution coastal models where temporal variability at daily or even higher frequency is needed (Sotillo et al., 2021a).

3.1.2 River discharge databases

In contrast, Rriver databases and services are progressively becoming available and provide better estimates of coastal runoff and river discharges at the global scale (Sotillo et al., 2021<u>a</u>). These databases typically assemble information from multiple data providers into coherent, gap-free and quality-controlled datasets. A few eExamples below are given herecategorized by

210 data source:

In situ databases:

- The Global Runoff Data Center² (GRDCGRDC), under the auspices of the WMO, is an international archives of quality_-controlled historical mean daily and monthly discharge data, from over 10,000 stations distributed inacross 159 countries, facilitating exchanges between data providers and data users. The most recent published version of the Exchanges into the Warld's Oceans³ detect head on the meter
- 215 Freshwater Fluxes into the World's OceansFreshwater Fluxes into the World's Oceans³ dataset, based on the water balance model WaterGAP, contains-provides annual runoff values estimates covering the from 1901-2016 period.
 - The Global Streamflow Indices and Metadata archive (GSIM), a collection of metadata and indices derived from more than 35 000 daily streamflow time series worldwide, gathered from 12 open databases (7 national and 5 international collections) (Do et al., 2018; Gudmundsson et al., 2018).
- A global dataset of monthly streamflow for 925 of the world's largest rivers connecting to the ocean was built by Dai et al. (2009), updated from Dai and Trenberth (2002).
 - A global database of monthly mean runoff for 986 rivers was incorporated in the NCOM, now HYCOM, U.S. model (Barron and Smedstad, 2002). It expands on the work of Perry et al. (1996) with corrections and additions derived from monthly mean streamflow from the U.S. Geological Survey (USGS) (Wahl et al., 1995), and extends the basic
- 225 <u>RivDIS database (Vörösmarty et al., 1998) to adjust for missing discharge attributed to small (ungauged) rivers.</u> *Model-derived databases:*
 - A 35-year daily and monthly global reconstruction of river flows (GRADES) at 2.94 million river reaches, with bias correction from machine-learning derived global runoff characteristics maps, was developed in support of the Surface Water and Ocean Topography (SWOT) satellite mission (Lin et al., 2019).
- A dataset of historical river discharge from 1958 to 2016 was created using the CaMa-Flood global river routing model and adjusted runoff from the land component of JRA-55 (Suzuki et al., 2018; Tsujino et al., 2018).
 - A global freshwater budget is included in the CORE.v2 datasets that have an accompanying database for continental runoff from rivers, groundwater and icebergs. These are estimated from continental imbalances between precipitation, evaporation and storage, and then distributed between bordering ocean basins based on river routing schemes and flow estimates (Large and Yeager, 2009).

Hybrid database:

• <u>EMODnet Physics⁴ provides ocean physics data and data products built with common standards, consisting of collections of in situ data, reanalysis, and aggregated in situ data and model outputs. As part of the available parameters, the operational river runoff data includes near-real time data from European river stations and a subset of</u>

the GRDC focusing on coastal areas and stations located near river mouths, which extend beyond European borders.

240

About 1,200 rivers worldwide are connected and operationally available.

² https://grdc.bafg.de/

³ https://fwf.bafg.de/

⁴ https://emodnet.ec.europa.eu/geoviewer/

- A 35 year daily and monthly global reconstruction of river flows (GRADES), with bias correction from machinelearning derived global runoff characteristics maps, was developed in support of the Surface Water and Ocean Topography (SWOT) satellite mission (Lin et al., 2019).
- A global dataset of monthly streamflow for 925 of the world's largest rivers connecting to the ocean was built by Dai et al. (2009), updated from Dai and Trenberth (2002).
 - A dataset of historical river discharge from 1958 to 2016 was created using the CaMa Flood global river routing model and adjusted runoff from the land component of JRA-55 (Suzuki et al., 2018; Tsujino et al., 2018).
- A global database of monthly mean runoff for 986 rivers was incorporated in the NCOM, now HYCOM, U.S. model
 (Barron and Smedstad, 2022), that expands on the work of Perry et al. (1996) with corrections and additions derived from monthly mean streamflow from the USGS (Wahl et al., 1995), and extends the basic RivDIS database (Vörösmarty et al., 1998) to make adjustments for missing discharge attributed to small (ungauged) rivers.
 - A database of pan Arctic river discharge (R Arcticnet).
 - Satellite-derived database:

- A database for Greenland liquid water discharge from 1958 through 2019 (Mankoff et al., 2020).
 - The largest known dataset compiles publicly available river gauge data, with satellite-based rating curves used to fill in the temporal gaps (Riggs et al., 2023).

Regional databases also exist, such as:

- Long-term (1993-2011) satellite-derived estimates of continental freshwater discharge into the Bay of Bengal (Papa et al., 2012).
 - A database of pan-Arctic river discharge (R-Arcticnet⁵).
 - A database for Greenland liquid water discharge from 1958 through 2019 (Mankoff et al., 2020).
 - A river discharge climatology and corresponding historical time series for all rivers flowing into the Adriatic Sea with an average climatological daily discharge exceeding 1 m³s⁻¹ (Aragão et al., 2024).
- Of particular importance is the fact that some of these databases use model-simulated runoff ratios (e.g. from Community Land Model (CLM) or river routing model) over gauged and ungauged drainage areas to estimate the contribution from the areas not monitored by the hydrometric network and adjust the station flow to represent river mouth outflow (-e.g. Dai et al. (2009). This allows more precise derivation of the total discharge into the global oceans, through the sum of both gauged and ungauged discharges.
- 270 <u>Unless explicitly stated (e.g. for EMODnet Physics)</u>, <u>It is not evident that anymost</u> of these databases are updated on a regularlack clearly stated update schedules; some remain static, <u>while</u> others are updated at irregularly <u>intervals</u>. Such databases are useful in the context of a reanalysis, but less so in an operational context where near-real-time data feeds are required. Furthermore, a detailed comparative assessment of these various data sources is still lacking.

⁵ https://www.r-arcticnet.sr.unh.edu/v4.0/index.html

Alternatively, indirect approaches using tidal statistics at the estuarine entrance from tidal stations rather than direct flow

275 measurements have been developed to estimate the net freshwater discharge at the mouth of an estuary, with the advantage of integrating processes at the basin scale, downstream of the last hydrometric station (Moftakhari et al., 2013; Moftakhari et al., 2016). Because tide gauge records at the coasts were often installed well before the onset of systematic river gauging (Talke and Jay, 2013), such inverse techniques make it possible to extend flow records back in time.

3.1.3 Operational river discharge products

- 280 While most river discharge databases are static, operational products have been developed for near-real-time applications. For example, <u>Ft</u>he Global Flood Awareness System, GloFAS-ERA5, is an operational global river discharge reanalysis produced consistently with the ECMWF ERA5 atmospheric reanalysis and providing global gridded data products from 1979 to near-real-time (within a 7-day delay) (Harrigan et al., 2020). Figure 2 illustrates the resolution of the river network that emerges in the GloFAS gridded data, and the association of discharge at the coast to point sources in a regional model of the northwest
 285 Atlantic Opera that is in development for future operations.
- 285 Atlantic Ocean that is in development for future operations.



Figure 2: Annual mean surface water discharge (<u>m³s⁻¹m³s-1</u>) in 0.1° x 0.1° cells of the GloFAS analysis from Harrigan et al. (2020) for the year 2023. Filled circles show the locations of 93 point sources in the prototype East Coast Community Ocean Forecast System (ECCOFS) ROMS model (domain denoted by the gray perimeter box) associated to GloFAS points near the coast that have long-term mean (2009-2019) discharge exceeding 50 <u>m³s⁻¹m³s-1</u>. <u>River networks come from GloFAS</u>.

Several centers are also producing continental- and global-scale hydrological (ensemble) forecasts operationally: the European Flood Awareness System (EFAS) (Thielen et al., 2009), the European Hydrological Predictions for the Environment (E-HYPE) (Donnelly et al., 2015), the Hydrologic Ensemble Forecast Service (HEPS) in the U.S. (Demargne et al., 2014), the Flood Forecasting and Warning Service (FWWS) in Australia, the National Surface and River Prediction System (NSRPS) in Canada

(Fortin et al., 2023); and globally, the World-Wide HYPE (WWH) (Arheimer et al., 2020) and GloFAS (Harrigan et al., 2023).
 Notably, as part of the GloFAS service evolution, global daily ensemble river discharge reforecasts (20-year) and real-time forecast (2020-present) datasets are made freely and openly available through the Copernicus Climate Change Service (C3S) Climate Data Store (CDS) (Harrigan et al., 2023).

Other projects have been supported by the Copernicus Marine Environment Monitoring Service (CMEMS), for example, the LAMBDA project regionally focused on the European Atlantic Façade and the North Sea. The resulting freshwater model estimates and in-situ observations are operationally updated and made available via the project viewer web interfaceviewer web interface⁶ (Sotillo et al., 2021a).

The FOCCUS project (Forecasting and Observing the Open-to-Coastal Ocean for Copernicus Users⁷) further enhances operational hydrological models by addressing the land-ocean continuum through improved river runoff estimations and the

- 305 development of advanced coupling between hydrological and coastal ocean models. FOCCUS builds on existing pan-European hydrological frameworks, such as E-HYPE and LISFLOOD, to provide dynamic freshwater inputs, including nutrient and inorganic matter transport. Additionally, the project integrates novel AI techniques to optimize estuarine modelling and freshwater forcing for coastal systems. These innovations directly contribute to refining CMEMS and supporting all European coastal services with more accurate and seamless coastal monitoring and forecasting capabilities.
- 310 In some instances, the regional products may appear to be the preferred option for some regional or local studies, as they were designed to specifically represent the hydrological characteristics of a given region, sometimes with higher resolution and accuracy. However, a global solution is attractive in data scarce areas and where consistency between discharge products and across all forcing variables is required over large domains (Polton et al., 2023).

3.1.4 Remotely-sensed discharges

315 <u>Remote sensing of river discharge is a rapidly advancing research field (see Gleason and Durand, 2020, and references therein).</u> With the <u>successful launch of the SWOT satellite launched</u> in December 2022, <u>new global</u> discharge products will <u>soon</u> become available <u>globally</u> at a nominal resolution of 10 km <u>over for</u> river reaches wider than 100 m, thus vastly expanding measurements of global rivers in both gauged and ungauged basins (Durand et al., 2023). Significant improvements on global uncalibrated models are expected (Emery et al., 2018). <u>Furthermore, the temporal variations in</u> SWOT<u>-derived</u> discharge <u>data</u>

⁶ http://www.cmems-lambda.eu/home.html

⁷ https://foccus-project.eu/

320 <u>is expected to improve-could lead to a better representation of the</u> global hydrological cycle <u>representation</u> and to enhanced ocean model solutions near the coast-when forced by SWOT discharges.

3.1.5 Machine learning-derived discharge estimates

Machine learning is increasingly used in hydrology for rainfall-runoff modelling, with Long Short-Term Memory (LSTM) networks (Greff et al., 2016; Hochreiter and Schmidhuber, 1997) proving particularly effective in capturing both periodic and

- 325 <u>chaotic patterns in time-series data while accurately learning long-term dependencies (Fang et al., 2017; Hu et al., 2019;</u> Mouatadid et al., 2019). In numerous hydrological studies, LSTM has demonstrated superior performance over traditional process-based models in simulating runoff, primarily in data-rich regions (Feng et al., 2020, 2021; Frame et al., 2022; Gauch et al., 2021; Hunt et al., 2022; Konapala et al., 2020; Kratzert et al., 2019; Lees et al., 2021; Li et al., 2023; Luppichini et al., 2024; Nearing et al., 2021; Reichstein et al., 2019). However, limited efforts have explored the transferability of LSTM models
- 330 to data-scarce regions (e.g. Akpoti et al., 2024), with Ma et al., (2021) and Muhebwa et al. (2024) (and references therein) being a few such exceptions. Recently, researchers have explored the potential of LSTM models for global river discharge estimations (Rasiva Koya and Roy, 2024; Tang et al., 2023; Yang et al. 2023; Zhao et al. 2021). However, extensive validation beyond the training basins is required to fully evaluate their suitability for global-scale implementations.

335 3.2 Salinity and temperature

Estuarine mixing processes affect the<u>influences</u> salinity distribution and its seasonal <u>variations-variability</u> near river mouths (Sun et al., 2019). Models are particularly sensitive to salinity in shelf areas and ROFI zones, most often due to the diverse treatment of OOFS given to coastal and river freshwater forcing (Sotillo et al., 2021<u>a</u>). Therefore, to assess the impact of a chosen formulation and evaluate model performances, sea surface salinity (SSS) and temperature (SST) are typically used.
The <u>climatology of the</u>-World Ocean Atlas <u>climatology</u> (Locarnini et al., 2013; Zweng et al., 2023) <u>can have a high positive often overestimates nearshore</u> salinity <u>bias nearshore and is not adequate, making it unsuitable</u> for model evaluation in these areascoastal regions. As an alternative, Sun et al. (2019) built on the original World Ocean Database and developed an improved salinity and temperature climatology with an enhanced representation of the coastal ocean. In-situ data and satellite

observations from SMOS, Aquarius and SMAP (Bao et al., 2019) can also be used to assess the impact of river forcing on sea

- 345 surface salinity (Feng et al., 2021). However, <u>seasonal variability in the skill of SSS retrievals can be associated with SST-dependent bias and strong land-sea differences in microwave emissivity, makinge satellite observations unreliable within some 70 km of the coast (<u>Grodsky et al., 2018; Menezes, 2020;</u> Vazquez-Cuervo et al., 2018). Higher resolution coastal satellite products have been <u>demonstrated_developed</u> based on empirical relationships between local salinity and ocean color observations (<u>e.g.</u>-Geiger et al, 2011; <u>Chen et al., 2017</u>), using deep neural networks trained on Sentinel-2 Level 1-C Top of</u>
- 350 Atmosphere (TOA) reflectance data (Medina-Lopez and Ureña-Fuentes, 2019; Medina-Lopez, 2020), or by relating the

reflectance of the visible bands from Sentinel-2 imagery with electrical conductivity, influenced by the concentration and composition of dissolved salts (Sakai et al., 2021), but although these are not applied globally applicable.

A recent study in the German Bight (Thao et al., 2024) demonstrated the critical role of high-resolution salinity inputs at estuarine mouths in improving the predictive capabilities of coupled wave-ocean models. Using the GCOAST model system,

355 which seamlessly integrates estuarine and coastal dynamics with regional ocean models, researchers validated salinity and temperature fields against in-situ observations. The results highlighted that estuarine inflows significantly enhance the accuracy of coastal ocean models.

Alternatively, salinity predictions in estuaries and at river mouths have been successfully estimated using machine learning approaches. A few examples can be found in the recent literature: Qiu and Wan (2013) developed an autoregressive model

- 360 relating salinity at a given time to past observations of salinity and physical drivers (freshwater inflow, rainfall, tidal elevation) in the Caloosahatchee River Estuary; Fang et al. (2017) used a genetic algorithm coupled with support vector machine to predict salinity in the Min River Estuary; Qi et al. (2022) applied four neural network models to emulate salinity simulations in the Sacramento-San Joaquin Delta from a process-based river, estuary and land modelling system; Guillou et al. (2023) were able to reproduce the seasonal and semi-diurnal variations of sea surface salinity at the mouth of the Elorn estuary (bay
- 365 of Brest), with support vector regression performing best among all tested algorithms. Despite these advancements, Improved and sustained operational high-resolution salinity monitoring for salinity observations is still-needed with enhanced spatial coverage and temporal repetition in order to build confidence in numerical solutions near the coast. Integrating salinity, temperature, and additional parameters such as nutrients and sediments directly into river outflows could further improve model accuracy (Verri et al., 2018; Thao et al., 2024). While these factors play a secondary
- 370 role in influencing oceanographic processes, their inclusion could advance research on coastal hypoxia, carbon cycling, and regional weather and climate, ultimately supporting seamless predictions of land-ocean-atmosphere feedbacks in nextgeneration Earth system models (Feng et al. 2021).

<u>43.3</u> Examples of current OOFS

This section describes howe status of implementation of river forcing is implemented in current OOFS is described in this section. The objective is to get a picture of the current landscape of approaches and data sources. While Cirano et al. (2024) provide a comprehensive overview of existing OOFS worldwide, the representation of rivers in these systems remains poorly documented and often buried in model configuration files. The list of systems presented below in Appendix A is therefore not exhaustive and is limited to a compilation of comments received as part of a survey conducted among members of the OceanPredict community in May 2023. It is meant to illustrate the diversity of methods employed for treating freshwater fluxes

380 in OOFS and associated input data sources, in <u>4 global</u>, <u>12 regional</u>, <u>4 coastal and <u>1 inland systems</u>. <u>Although the survey covers</u> <u>a limited number of systems</u>, <u>-t</u>The literature review <u>presented</u> in <u>the previous sections provides offers other additional</u> examples to complete the picture.</u> Figure 3 provides a graphical summary of the 6 river forcing methods and 4 data sources used in the OOFS listed in Appendix A. In terms of river forcing methods, most systems specify vertical or lateral freshwater fluxes to account for riverine inputs.

385 Only a few of them rely on more sophisticated approaches that use channel extensions within the ocean model or routing schemes from hydrological models to transport the water from the watershed to the coast. Furthermore, none of the global systems surveyed use lateral boundary conditions, likely due to insufficient spatial resolution near river mouths.

In terms of the data sources used in OOFS, what stands out from the survey is the use of in situ data as a primary source in most systems, and climatology either as a primary or fallback source of freshwater discharge. Global systems tend to opt for climatologies in comparison with regional or coastal systems that favour observed data when available, which allows to capture

both seasonal and non-seasonal events and their potential local or regional impacts. Only a few regional and inland systems use hydrological models or reanalyses as primary data sources.



Figure 3: Graphical summary from a survey on river forcing methods (left panel) and data sources (right panel) used in global,
 regional, coastal and inland OOFS listed in Appendix A. Coloured bars indicate the primary data sources or methods, whereas dashed bars represent secondary data sources used as a fallback when primary sources are unavailable.

Additional considerations were also highlighted by the respondents, essential for appropriately representing river inflow in ocean models and addressing challenges such as numerical instabilities and data limitations. For example, spatial smoothing around the river source, or equivalently, optimizing the integration distance for equivalent coastal precipitation may be required to prevent numerical instabilities. Similarly, an increased diffusivity within the surface mixing layer can be implemented to simulate the effects of river inflow. Salinity and temperature of the input freshwater can either be set to zero and to the local SST, respectively, or derived from a combination of real-time gauge data and monthly averages when available. For ungauged areas, river gauge data can be scaled, or additional coastal runoff can be incorporated. In contrast, some systems directly convert precipitation data into river discharges, disregarding hydrological processes and assuming an instantaneous response.

405 In sum, the representation of rivers in OOFS requires careful consideration of various numerical methods, data sources, and modelling approaches. However, some simplifications may limit accuracy in applications requiring high regional precision.

5 Summary and recommendations

The assessment of river forcing implementation in OOFS highlights the complexity and challenges of accurately integrating riverine freshwater discharges into ocean models. Despite the growing demand for operational oceanographic products,

- 410 especially in coastal areas (Ciliberti et al., 2023), OOFS river forcing still faces shortcomings related to the representation of physical processes, data availability, and data quality. The parameterization of river inputs and the interaction between model components, often nonlinear, remain unresolved issues, underscoring the absence of standardized practices for river forcing. Addressing these gaps requires advancements in model physics, improved spatial and temporal resolution, and enhanced coupling between land, ocean and atmosphere. Furthermore, the incorporation of river flow varies regionally, largely due to
- 415 differences in the availability and quality of river discharge, salinity and bathymetric datasets, and is further influenced by model scale and resolution. As the demand for reliable coastal forecasts grows, real-time, high-quality river discharge data becomes increasingly pressing. Standardized methodologies and improved integration of riverine parameters—including salinity, temperature, and biogeochemical components—will facilitate seamless watershed-ocean coupling and improve predictions of coastal dynamics, particularly under extreme conditions.
- 420 Service evolution roadmaps, such as those outlined by CMEMS, emphasize the need for a better characterization of coastal freshwater exchanges to improve forecasts, especially during severe weather events (Sotillo et al., 2021b). A key step forward involves the progressive replacement of static climatologies with real-time, updated time series (past, present, and forecasts) of river inputs, covering both major and minor or ephemeral streams. Recommendations have been made towards standardized freshwater inputs (and associated river inputs of nutrients and sediment loading), harmonized river forcing approaches, and a
- 425 more integrated watershed-ocean strategy (Campuzano et al., 2016; Capet et al., 2020; Sobrinho et al., 2021). Additionally, ensuring validated observational error estimates for estuary-mouth forcing, including river discharge and auxiliary variables such as coastal salinity, is crucial for model accuracy (De Mey-Frémaux et al., 2019; Polton et al., 2023). Improved interfaces between coastal monitoring and modelling systems are therefore essential. The FOCCUS project exemplifies progress in addressing these challenges through advancements in hydrological and estuarine modelling, dynamic freshwater inputs, and
- the integration of AI-driven tools to refine river discharge estimations and coastal system forecasts.
 Future efforts must focus on refining model physics, resolution, and coupling strategies to better integrate the land-ocean continuum. Standardized methodologies and integrated high-quality data sources, together with continued interdisciplinary collaboration and technological advancements, will be key to overcoming existing limitations and ensuring more accurate and reliable ocean predictions. Such efforts are critical for improving predictions of coastal dynamics and for fostering a deeper understanding of their implications on global climate and ecosystem functioning.
 - 15

Appendix A: Survey on river forcing methods and data sources in current OOFS

This Appendix presents results of a survey conducted among members of the OceanPredict community in May 2023. The
 responses are reported in the following tables as given by the participants; nearly no changes were made to each contributed entry, except for a few added references and acronym definitions.

3.3

A.1 Global systems

System	<u>Institution</u>	Domain(s)	Resolution	Circulation Model	Method for river forcing	Data sources
MOVE/	<u>Japan</u>	Global	1/4°	MRI.COM Ver.	River discharge is expressed as a	Climatology of JRA-55do river
MRI.COM-	Meteorological			4	part of the surface freshwater	runoff data
<u>G3 8</u>	Agency (JMA)'s					
(Multivariate	Meteorological					
Ocean	Research					
<u>Variational</u>	Institute					
Estimation/						
Meteorological						
Research						
Institute						
Community						
Ocean Model -						
Global version						
<u>3)</u>						
GEOS 9	NASA's Global	Global	25 km –	MOM6	GEOS-land component run off,	In situ data, land/catchment
(NASA	Modeling and		4 km		routed to catchments	model
Goddard Earth	Assimilation					
Observing	<u>Office</u>					
<u>System)</u>						

<u>⁸ https://ds.data.jma.go.jp/tcc/tcc/products/elnino/move_mricom-g3_doc.html</u> <u>⁹ https://gmao.gsfc.nasa.gov/GEOS_systems/</u>

RTOFSv2 ¹⁰	NOAA's	Global	0.08°	HYCOMv2.2	Rivers are implemented as	RTOFS uses global climatology
(Real-Time	National Centers				virtual salt flux at the ocean	of monthly mean river discharge
<u>Ocean</u>	<u>for</u>				surface. River runoff is	created at the Naval Research
Forecast	Environmental				distributed over several ocean	Laboratory (NRL) (Barron and
<u>System)</u>	Prediction				grid points around the river	Smedstad, <u>20222002</u>). It
					source by applying spatial	provides monthly runoff for 986
					smoothing to spread out the	rivers. The dataset is based on
					effect of the river and prevent	the Perry (1996) data with
					negative salinities due to	corrections and additions
					numerical overshooting. To	derived from: (1) monthly mean
					mimic the river inflow, river	streamflow over all years,
					freshwater is mixed from the	accessible from the USGS
					surface down to a depth specified	(Wahl et al., 1995); (2)
					by the user (set to 6 meters in	the Global River Discharge
					RTOFS). In the grid cells with	(RivDIS) database (Vörösmarty
					not-zero river runoff and in the	et al., 1998); (3) the Regional,
					upper layers, river freshwater is	Hydrometeorological Data
					mixed within increased vertical	Network (R-Arcticnet <u>R-</u>
					diffusivity. Alternatively, rivers	<u>Arcticnet¹¹</u>) database provides
					can be added directly to the input	most of the information
					precipitation fields, which is a	ultimately used on rivers
					better option for a higher (than	flowing into the Arctic,
					monthly) frequency river flow	primarily rivers in Russia and
					data. It is possible to treat rivers	Canada.
					(as well as evaporation minus	
					precipitation, E-P) as a mass	
					exchange (not activated in	
					RTOFS).	

¹⁰ https://polar.ncep.noaa.gov/global/about/ ¹¹ http://www.r-arcticnet.sr.unh.edu/

FOAM-CPL-	UK Met Office	Global	1/4°	NEMO v3.6	Fresh water runoff from land is	Climatological river runoff
<u>NWP ¹²</u>					input in the surface layer of the	fields were derived by
(Forecast					ocean with the assumption that	Bourdalle-Badie and Treguier
<u>Ocean</u>					the runoff is fresh and at the	(2006) based on estimates given
Assimilation					same temperature as the local sea	in Dai and Trenbert (2002)
Model,					surface temperature. An	(Blockley et al., 2014)
<u>Coupled</u>					enhanced vertical mixing of	
Numerical					$2x10^{-3}$ m ² s ⁻¹ is added over the top	
Weather					10 m of the water column at	
Prediction					runoff points to mix the runoff	
					vertically and avoid instabilities	
					associated with very shallow	
					fresh layers at the surface	
					(Storkey et al., 2018).	

3.3A.2 Regional systems

 A.2: Examples of river forcing methods and data sources in regional OOFS.

System	Institution	Domain(s)	Resolution	Circulation	Method for river forcing	Data sources
				Model		
MOVE/MRI.C	<u>Japan</u>	North	2 km - 10 km	MRI.COM	River discharge is expressed as	Climatology of JRA-55do river
OM-NP/JPN ¹³	Meteorological	Pacific		Ver. 5	a part of the surface freshwater	runoff data
(Multivariate	Agency					
<u>Ocean</u>	(JMA)'s					
Variational	Meteorological					
Estimation/						

¹² https://www.metoffice.gov.uk/services/data/met-office-data-for-reuse/model ¹³ https://www.data.jma.go.jp/kaiyou/data/db/kaikyo/knowledge/move_jpn/system.html

	-			-		
Meteorological	Research					
Research	Institute					
Institute						
Community						
<u>Ocean Model –</u>						
North Pacific/						
Japan)						
			 !			
$\underline{\mathrm{TOPAZ}^{14}}$	Norwegian's	Arctic and	12 km	НҮСОМ	Removal of salt from the surface	Swedish Meteorological and
	<u>Nansen</u>	Nordic Seas			(an ellipse around the river	Hydrological Institute (SMHI)
	Environmental				mouth) and barotropic water	(Arctic-HYPE and E-HYPE),
	and Remote				flux. We use nutrients (N, P and	GRACE satellite for Greenland
	Sensing Center				Si) from the globalNEWS	mass loss and a home-made
	(NERSC)				model and scale them by river	climatology for Greenland
					discharge.	surface mass balance.
eSA-Marine ¹⁵	South	South	2.5 km and 0.5	ROMS	None, intermittent river input is	None
	Australian	Australian	km		usually weak to non-existent.	
	Research and	Gulfs and			-	
	Development	Shelf				
	Institute					
<u>DMI</u> _	<u>Danish</u>	Arctic and	4-10 km: ~5	HYCOM +	River forcing is converted to	River forcing is taken from
HYCOM	Meteorological	Atlantic	km	CICE fully	monthly means precipitation	various sources using a dataset
$\underline{\text{CICE}^{16}}$	Institute (DMI)	Oceans	throughout	coupled using	equivalents [m/s] for ~50,000	from the Geological Survey of
	-			Eastly Countries	minum aff and and	
			Arctic and	Earth System	river-runoii outiets and	Denmark and Greenland
			Arctic and northern	<u>Modeling</u>	distributed to the nearest coastal	(Mankoff et al., 2020),
			Arctic and northern Atlantic	<u>Modeling</u> <u>Framework</u>	distributed to the nearest coastal	(Mankoff et al., 2020),

 ¹⁴ https://nersc.no/en/products-and-services/analysis-tools-and-models/ocean-models/
 ¹⁵ https://pir.sa.gov.au/research/services/esa_marine/about_esa-marine
 ¹⁶ https://ocean.dmi.dk/models/hycom.uk.php

					(ESMF)	model grid point(s) (Ponsoni et	converted to monthly means
					coupler. CICE	al., 2023).	precipitation equivalents [m/s]
					runs on a		
					subset of the		
					full HYCOM		
					domain		
	DKSS ¹⁷ (Danish	<u>Danish</u>	North Sea -	3 nautical	HBM ,	River forcing is treated as a	European hydrological model
	Storm Surge	Meteorological	Baltic Sea,	miles	(Hiromb-	freshwater flux into coastal grid	E-HYPE3, from which an
	<u>System)</u>	<u>Institute</u>	with	(coarsest) to	Baltic Model)	cells. Water temperature equal	annual plus a calendar day ~30y
			multiple	0.1 nautical		to receiving cell (river	climatology has been derived
			nested	mile (finest)		temperature data not used) with	and used as a back-up for a
			subdomains			0°C as lower limit to avoid	daily forecast. The forecast
						instantaneous freezing.	model is run by the Swedish
							Hydrological and
							Meteorological Institute, and
							the day-to-day service comes
							with an annual fee.
-							
	IBI Near-Real-	Iberia Biscay	European	1/36°, Surface	NEMO v3.6	Freshwater river discharge	Data come from different
	Time ¹⁸	Irish (IBI) Sea	Atlantic	and 3D fields		inputs are implemented as	sources, depending on their
		 Monitoring 	façade (the	(50 vertical		lateral open boundary	availability, in the following
		Forecasting	Iberia-	levels)		conditions for the main 33 rivers	order: (1) Model data: SMHI
		<u>Center</u>	Biscay-			of the IBI area. The system also	hydrologic model; (2) Monthly
			Ireland			incorporates an extra coastal	climatological data taken from
			zone): Lat:			runoff rate (derived from the Dai	GRDCGRDC, French "Banque
			from 26N to			and Trenberth (2002)	HydroBanque Hydro" ¹⁹
- 1				1	1		

 $[\]frac{17}{\text{https://opendatadocs.dmi.govcloud.dk/Data/Forecast}} Data Storm Surge Model DKSS#:~:text=DKSS\%20is\%20DMI\%27s\%20forecast\%20m}{DKSS} DKSS\%20is\%20DMI\%27s\%20forecast\%20m}{DKSS} DKSS\%20DMI\%20s$ {DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS\%20DMI\%20s{DKSS} DKSS odel,ice%20thickness%20and%20ice%20concentration

¹⁸ https://marine.copernicus.eu/about/producers/ibi-mfc ¹⁹ http://www.hydro.eaufrance.fr/

	1					
		56N, Lon: from 19W to 5E			climatology, on a monthly basis), which makes the IBI forcing consistent with the ones imposed in the parent Copernicus Marine GLOBAL system.	dataset, Copernicus Marine Service and EmodnetEMODnet.
IBI Multi-Year ²⁰	Iberia Biscay Irish (IBI) Sea – Monitoring Forecasting Center	European Atlantic facade (the Iberia- Biscay- Ireland zone): Lat: from 26N to 56N, Lon: from 19W to 5E	1/12°, Surface and 3D fields (50 vertical levels)	NEMO v3.6	Same as IBI-NRT, but with an additional river (LAGAN)	Data come from different sources, depending on their availability, in the following order: (1) In-situ data: daily measurements from Copernicus Marine Service, Emodnet EMODnet_or national web sites; (2) Model data: SMHI hydrologic model.
CBEFS21(ChesapeakeBayEnvironmentalForecastSystem)	<u>Virginia</u> <u>Institute of</u> <u>Marine</u> <u>Science</u>	Chesapeake Bay	600 m x 600 m	ROMS	Freshwater - Real time USGS river gauge data is scaled to better represent total freshwater inflows over a larger area based on a watershed model. The scaled discharge is then disaggregated into the main river inflow and smaller streams based on proportions developed from the watershed model. The	In situ gauge data. Hindcast watershed model information. Artificial Neural Networks.

²⁰ https://marine.copernicus.eu/about/producers/ibi-mfc ²¹ https://www.vims.edu/research/products/cbefs/

	1		r			
					forecast is a simple autoregressive model based on the past few days. Riverine Biogeochemistry - Inputs are specified using Artificial Neural Network AI models based on the discharge and date, which recreate what the watershed model would have predicted had the current and forecast conditions been simulated by the watershed model. Temperature - Water	
					data and monthly averages depending on what is available.	
DREAMS22(RIAMReal-TimeOceanForecasting)	Kyushu University's Research Institute for Applied Mechanics (RIAM)	East Asian marginal seas	0.3 – 22 km	RIAM Ocean Model	Coastal precipitation is directly converted into the amount of river discharges. The integration distance was optimized by using model Green's functions (Hirose, 2011).	Grid point value (GPV) precipitation data of Japan Meteorological Agency (JMA)

²² https://dreams-c1.riam.kyushu-u.ac.jp/vwp/

FOAM-	UK Met Office	Northwest	1.5 km	NEMO v3.6	For each river input location, a	River runoff is primarily
<u>AMM15</u> 23		European			daily freshwater flux is	derived from a daily
(Forecast Ocean		Shelf Seas			assigned, with depth determined	climatology of gauge
Assimilation					by the average ratio of runoff to	measurements averaged for
Model- Atlantic					tidal range (as per the estuary	1980–2014. UK data were
Margin model					classifications of Cameron and	processed from raw data
<u>1.5km)</u>					Pritchard, 1963). The runoff	provided by the Environment
					temperature is assumed to align	Agency, the Scottish
					with the local sea surface	Environment Protection
					temperature (SST), as the	Agency, the Rivers Agency
					climatology does not include	(Northern Ireland), and the
					temperature data (Graham et al.,	National River Flow Archive
					2018).	(gauge data were provided by
						Sonja M. van Leeuwen,
						CEFAS, Lowestoft, UK,
						personal communication,
						2016). For major rivers that
						were missing from this data set
						(e.g. along the French and
						Norwegian coasts), data have
						been provided from an earlier
						climatology (Vörösmarty et al.,
						2020; Young and Holt, 2007),
						based on a daily climatology of
						gauge data averaged for the
						period 1950-2005 (Tonani et
						al., 2019).
	1					1

 ²³ https://www.metoffice.gov.uk/services/data/met-office-data-for-reuse/model

 24

FOAM- AMM7 ²⁴ (Forecast Ocean Assimilation Model- Atlantic Margin model 7km)	<u>UK Met Office</u>	Northwest European Shelf Seas	7 km	NEMO v3.6 (coupled to ERSEM 20.10 for biogeochemis try)	For each river input location, a daily freshwater flux is assigned, with depth determined by the average ratio of runoff to tidal range (as per the estuary classifications of Cameron and Pritchard, 1963). The runoff temperature is assumed to align with the local sea surface temperature (SST), as the climatology does not include temperature data (Graham et al., 2018).	Daily timeseries of river discharge, nutrient loads (nitrate, phosphate, silicate, ammonia), alkalinity (bioalkalinity, dissolved organic carbon) and oxygen were produced from an updated version of the river dataset used in Lenhart et al. (2010), combined with climatology of daily discharge data from the Global River Discharge Database (Vörösmarty et al., 2020) and from data prepared by the Centre for Ecology and Hydrology as used by Young and Holt, 2007. The climatology has an annually- varying component until 2018 to account for historic changes in nutrient loads, values for 2018 are used as a climatology in the operational system (Kay et al., 2020).
DOPPIO ²⁵ Rutgers University-and	<u>Rutgers</u> <u>University</u>	Northeast USA and	7 km	ROMS	Discharge is introduced as volume flux divergence (method LwSrc in ROMS) at	et al., 2020). Daily USGS discharge data are scaled for ungauged portions of the watershed based on the

²⁴ https://www.metoffice.gov.uk/services/data/met-office-data-for-reuse/model ²⁵ https://gmd.copernicus.org/articles/13/3709/2020/

MARACOOS ²⁶	Nova Scotia,		27 point sources in model cells	statistics of a 10-year
(Mid-Atlantic	Canada		adjacent to the coast.	hydrological model analysis.
Regional				
Association				
Coastal Ocean				
Observing				
System)				

3.3A.3 Coastal systems

 A.3: Examples of river forcing methods and data sources in coastal OOFS.

System	<u>Institution</u>	Domain(s)	Resolution	Circulation Model	Method for river forcing	Data sources
DFO's Port	Government of	Kitimat Fjord,	20 - 200 m	NEMO 3.6	NEMO's runoff feature for	Gauge data (from
<u>Ocean</u>	<u>Canada's</u>	Vancouver			some rivers, and a SSH open	Environment and Climate
Prediction	Department of	Harbour, Lower			boundary condition for	<u>Change Canada,</u> ECCC)
Systems ²⁷	Fisheries and	Fraser River, St			others	where available,
	Oceans (DFO)	Lawrence				climatology elsewhere
		Estuary, Port of				
		Canso, Saint				
		John harbour				
CIOPS 28	Environment and	East/West +	1/36° +	NEMO 3.6	Same as DFO port models	Gauge data for Fraser
(Coastal Ice-	Climate Change	SalishSea500	500m for			River, climatology
Ocean	Canada (ECCC)		SS500			elsewhere
Prediction						
System)						

²⁶ https://maracoos.org/ ²⁷ https://publications.gc.ca/site/eng/9.905464/publication.html ²⁸ https://eccc-msc.github.io/open-data/msc-data/nwp_ciops/readme_ciops_en/ 26

<u>FANGAR _</u> <u>BAY²⁹</u>	<u>Universitat</u> Politècnica de <u>Catalunya</u>	Ebro Delta	350m / 70m	COAWST (ROMS/ SWAN)	Climatological freshwater from Ebro River	In situ data
NARF 30 (Northern Adriatic Reanalysis and Forecasting system)	Istituto Nazionale di Oceanografia e di Geofisica Sperimentale	Northern Adriatic Sea (Mediterranean Sea)	1/128° (~750 m)	MITgcm-BFM (coupled hydrodynamic- biogeochemical)	The downstream end of the rivers flowing into the basin is simulated as a narrow channel: one or two cells in the horizontal direction and a few vertical levels. Freshwater discharge rates from NRT data or climatologies are converted into horizontal velocities (the section of the riverbed is known) and applied as lateral open boundary conditions. Salinity is constant (5 PSU), temperature has a yearly sinusoidal cycle (maxima and minima in summer and winter, respectively) and biogeochemical concentrations are derived from literature/climatologies.	In-situ NRT discharge data for the Po River (main contributor), climatologies for the others (with sinusoidal modulation: maxima in spring/fall, minima in summer/winter). Daily frequency.

²⁹ https://doi.org/10.5194/egusphere-egu24-11220
³⁰ https://medeaf.ogs.it/got

450 3.3A.4 Inland systems

Table A.4: Example of river forcing methods and data sources in inland OOFS.

System	<u>Institution</u>	Domain(s)	Resolution	Circulation Model	Method for river forcing	Data sources
WCPS 31 (Water Cycle	Environment and Climate Change Canada (ECCC)	Great-Lakes+ <u>Northwest</u> <u>Atlantic (</u> NWA <u>)</u>	1/36° + 1km	NEMO 3.6	Fully coupled hydrologic model for GL, climatology for NWA	Hydrological model uses gauge data
Prediction System)						

³¹ https://eccc-msc.github.io/open-data/msc-data/nwp_wcps/readme_wcps_en/

References

460

470

- Akpoti, K., Mekonnen, K., Leh, M., Owusu, A., Dembélé, M., Tinonetsana, P., Seid, A., & Velpuri, N. M. State of continental
- <u>discharge estimation and modelling: challenges and opportunities for Africa. *Hydrological Sciences Journal*, 69(15),
 2124–2152. https://doi.org/10.1080/02626667.2024.2402938, 2024.
 </u>
 - Alvarez Fanjul, E., Ciliberti, S., Bahurel, P.: Implementing Operational Ocean Monitoring and Forecasting Systems. IOC-UNESCO, GOOS-275. <u>https://doi.org/10.48670/ETOOFS</u>, 2022.

Aragão, L., Mentaschi, L., Pinardi, N., Verri, G., Senatore, A., & di Sabatino, S. The freshwater discharge into the Adriatic Sea revisited. *Frontiers in Climate*, 6. https://doi.org/10.3389/fclim.2024.1368456, 2024.

- Arheimer, B., Pimentel, R., Isberg, K., Crochemore, L., Andersson, J. C. M., Hasan, A., and Pineda, L.: Global catchment modelling using World-Wide HYPE (WWH), open data, and stepwise parameter estimation, Hydrol. Earth Syst. Sci., 24, 535-559. <u>https://doi.org/10.5194/hess-24-535-2020</u>, 2020.
- Bao, D., Xue, Z.G., Warner, J.C., Moulton, M., Yin, D., Hegermiller, C.A., Zambon, B., He, R.: A Numerical Investigation of
- Hurricane Florence-Induced Compound Flooding in the Cape Fear Estuary Using a Dynamically Coupled Hydrological Ocean Model. Journal of Advances in Modeling Earth Systems, 14(11), e2022MS003131.
 https://doi.org/10.1029/2022MS003131, 20220.
 - Bao, S., Wang, H., Zhang, R., Yan, H., and Chen, J.: Comparison of Satellite-Derived Sea Surface Salinity Products from SMOS, Aquarius, and SMAP. Journal of Geophysical Research: Oceans, 124(3), 1932-1944. https://doi.org/10.1029/2019JC014937, 2019.
 - Barron, C. N., and Smedstad, L. F.: Global river inflow with<u>in the</u> Navy Coastal Ocean Model. OCEANS '02 MTS/IEEE, Biloxi, MI, USA, 2002, pp. 1472-1479 vol.3. DOI: 10.1109/OCEANS.2002.1191855, 20222002.
 - Blayo, E., and Debreu, L.: Revisiting open boundary conditions from the point of view of characteristic variables. Ocean Modelling, 9(3), 231-252. <u>https://doi.org/10.1016/j.ocemod.2004.07.001</u>, 2005.
- 475 Blockley, E. W., Martin, M. J., McLaren, A. J., Ryan, A. G., Waters, J., Lea, D. J., Mirouze, I., Peterson, K. A., Sellar, A., and Storkey, D.: Recent development of the Met Office operational ocean forecasting system: an overview and assessment of the new Global FOAM forecasts, Geosci. Model Dev., 7, 2613-2638. <u>https://doi.org/10.5194/gmd-7-2613-2014</u>, 2014.

Bonamano, S., Federico, I., Causio, S., Piermattei, V., Piazzolla, D., Scanu, S., Madonia, A., Madonia, N., de Cillis, G., Jansen, E., Fersini, G., Coppini, G., & Marcelli, M. River–coastal–ocean continuum modeling along the Lazio coast (Tyrrhenian

- 480 <u>Sea, Italy</u>): Assessment of near river dynamics in the Tiber delta. *Estuarine, Coastal and Shelf Science*, 297, 108618.
 <u>https://doi.org/https://doi.org/10.1016/j.ecss.2024.108618, 2024.</u>
 - Cameron, W. M., and Pritchard, D. W.: Estuaries. in The Sea, vol. 2, M. N. Hill, Ed. New York: John Wiley & Sons, 1963, 306-324, 1963.

Campuzano, F., Brito, D., Juliano, M. et al.: Coupling watersheds, estuaries and regional ocean through numerical modelling

- 485 for Western Iberia: a novel methodology. Ocean Dynamics 66, 1745-1756. <u>https://doi.org/10.1007/s10236-016-1005-4</u>, 2016.
 - <u>Capet, A., Fernández, V., She, J., Dabrowski, T., Umgiesser, G., Staneva, J., Mészáros, L., Campuzano, F., Ursella, L., Nolan, G., and El Serafy, G.: Operational Modeling Capacity in European Seas—An EuroGOOS Perspective and Recommendations for Improvement, Front. Mar. Sci., 7, 1–19, https://doi.org/10.3389/fmars.2020.00129, 2020.</u>
- 490 Chandanpurkar, H. A, Lee, T., Wang, X., Zhang, H., Fournier, S., Fenty, I., Fukimori, I., Memnemnlis, D., Piecuc, C.G., Reager, J.T., Wang, O. and Worden, J.: Influence of Nonseasonal River Discharge on Sea Surface Salinity and Height. Journal of Advances in Modeling Earth Systems, 14(2), e2021MS002715. <u>https://doi.org/10.1029/2021MS002715</u>, 2022.
 - Chen, S., & Hu, C. Estimating sea surface salinity in the northern Gulf of Mexico from satellite ocean color measurements. *Remote Sensing of Environment*, 201, 115–132. https://doi.org/https://doi.org/10.1016/j.rse.2017.09.004, 2017.
- 495 Cheng, H., Cheng, J. C., Hunter, R. M., and Lin, H.: Demonstration of a Coupled Watershed-Nearshore Model. Available at https://apps.dtic.mil/sti/citations/ADA518953, 2010 (last access: 28/07/2024).
 - Choi, B.-J., and Wilkin, J. L.: The Effect of Wind on the Dispersal of the Hudson River Plume. Journal of Physical Oceanography, 37(7), 1878–1897. <u>https://doi.org/10.1175/JPO3081.1</u>, 2007.
 - Ciliberti, S., Alvarez Fanjul, E., Pearlman, J., Wilmer-Becker, K., Bahurel, P., Ardhuin, F., Arnaud, A., Bell, M., Berthou, S.,
- Bertino, L., Capet, A., Chassignet, E., Ciavatta, S., Cirano, M., Clementi, E., Cossarini, G., Coro, G., Corney, S., Davidson, F., Drevillon, M., Drillet, Y., Dussurget, R., El Serafy, G., Fennel, K., Garcia Sotillo, M., Heimbach, P., Hernandez, F., Hogan, P., Hoteit, I., Joseph, S., Josey, S., Le Traon, P.-Y., Liying, W., Libralato, S., Mancini, M., Matte, P., Melet, A., Miyazawa, Y., Moore, A., Novellino, A., Porter, A., Regan, H., Romero, L., Schiller, A., Siddorn, J., Staneva, J., Thomas-Courcoux, C., Tonani, M., Veitch, J., von Schuckmann, K., Wilkin, J., and Zufic, R.: Evaluation of
- 505operational ocean forecasting systems from the perspective of the users and the experts, State Planet, Copernicus Mar.
Serv. Ocean State Rep. 7, 1-osr7, https://doi.org/10.5194/sp-1-osr7-2-2023, 2023.
- <u>Cirano, M., Alvarez-Fanjul, E., Capet, A., Ciliberti, S., Clementi, E., Dewitte, B., Dinápoli, M., el Serafy, G., Hogan, P., Joseph, S., Miyazawa, Y., Montes, I., Narvaez, D., Regan, H., Simionato, C. G., Tanajura, C. A. S., Thupaki, P., Urbano-Latorre, C., & Veitch, J. A description of existing Operational Ocean Forecasting Services around the Globe. *State Planet Discuss.*, 2024, 1–44, https://doi.org/10.5194/sp-2024-26, 2024.
 </u>
 - Cossarini, G., Moore, A., Ciavatta, S., & Fennel, K. Numerical Models for Monitoring and Forecasting Ocean Biogeochemistry: a short description of present status. *State Planet Discuss.*, 2024, 1–13. https://doi.org/10.5194/sp-2024-8, 2024.
 - Dai, A., and Trenberth, K. E.: Estimates of Freshwater Discharge from Continents: Latitudinal and Seasonal Variations.
- 515 Journal of Hydrometeorology, 3(6), 660-687. <u>https://doi.org/10.1175/1525-7541(2002)003<0660:EOFDFC>2.0.CO;2</u>, 2002.

- Dai, A., Qian, T., Trenberth, K.E., and Milliman, J.D.: Changes in Continental Freshwater Discharge from 1948 to 2004. Journal of Climate, 22(10), 2773-2792. <u>https://doi.org/10.1175/2008JCL12592.1</u>, 2009.
- De Mey-Frémaux, P., Ayoub, N., Barth, A., Brewin, R., Charria, G., Campuzano, F., and Al., E.: Model-observations synergy
 in the coastal ocean, Front. Mar. Sci., 6, 436, https://doi.org/10.3389/fmars.2019.00436, 2019.
 - Demargne, J., Wu, L., Regonda, S.K., Brown, J.S., Lee, H., He, M., Seo, D.K., et al.: The Science of NOAA's Operational Hydrologic Ensemble Forecast Service. Bull. Am. Meteorol. Soc., 95(1),- 79-98. <u>https://doi.org/10.1175/BAMS-D-12-00081.1</u>, 2014.
- Do, H. X., Gudmundsson, L., Leonard, M., & Westra, S. The Global Streamflow Indices and Metadata Archive (GSIM) Part
 1: The production of a daily streamflow archive and metadata. *Earth Syst. Sci. Data*, 10(2), 765–785. https://doi.org/10.5194/essd-10-765-2018, 2018.
 - Donnelly, C., Andersson, J. C. M., and Arheimer, B.: Using flow signatures and catchment similarities to evaluate the E-HYPE multi-basin model across Europe. Hydrological Sciences Journal, 61(2), 255-273. https://doi.org/10.1080/02626667.2015.1027710, 2015.
- 530 Durand, F., Piecuch, C. G., Becker, M., Papa, F., Raju, S. v, Khan, J. U., & Ponte, R. M. Impact of Continental Freshwater Runoff on Coastal Sea Level. Surveys in Geophysics, 40(6), 1437–1466. https://doi.org/10.1007/s10712-019-09536-w, 2019.

550

- Durand, M., Gleason, C.J., PAvelsky, T.M., de Moraes Frasson, R.P., Turnmon, M., et al.: A Framework for Estimating Global River Discharge From the Surface Water and Ocean Topography Satellite Mission. Water Resources Research, 59(4), e2021WR031614. https://doi.org/10.1029/2021WR031614, 2023.
- Dzwonkowski, B., Greer, A.T., Briseño-Avena, C. Krause, J.W., Soto, I.M., Hernandez, F.J., Deary, A.L., Wiggert, J.D., Joung, D., Fitzpatrick, P.J., O'Brien, S.J., Dykstra, S.L., Lau, Y., Cambazoglu, M.K., Lockridge, G., Howden, S.D., Shiller, A.M., Graham, W.M.: Estuarine influence on biogeochemical properties of the Alabama shelf during the fall season. Continental Shelf Research, 140, 96-109. <u>https://doi.org/10.1016/j.csr.2017.05.001</u>, 2017.
- 540 Emery, C. M., Paris, A., Biancamaria, S., Boone, A., Calmant, S., Garambois, P.-A., and Santos da Silva, J.: Large-scale hydrological model river storage and discharge correction using a satellite altimetry-based discharge product, Hydrol. Earth Syst. Sci., 22, 2135-2162. <u>https://doi.org/10.5194/hess-22-2135-2018</u>, 2018.
 - Fang, Y., Chen, X., & Cheng, N.-S. Estuary salinity prediction using a coupled GA-SVM model: a case study of the Min River Estuary, China. Water Supply, 17(1), 52–60. https://doi.org/10.2166/ws.2016.097, 2016.
- 545 Fang, K., Shen, C., Kifer, D., & Yang, X. Prolongation of SMAP to Spatiotemporally Seamless Coverage of Continental U.S. Using a Deep Learning Neural Network. *Geophysical Research Letters*, 44(21), 11, 11–30, 39. <u>https://doi.org/https://doi.org/10.1002/2017GL075619, 2017.</u>

Feng, D., Fang, K., & Shen, C. Enhancing streamflow forecast and extracting insights using long-short term memory networks
 with data integration at continental scales. *Water Resources Research*, 56(9), e2019WR026793.
 https://doi.org/10.1029/2019WR026793, 2020.

- Feng, D., Lawson, K., & Shen, C. Mitigating Prediction Error of Deep Learning Streamflow Models in Large Data-Sparse Regions With Ensemble Modeling and Soft Data. *Geophysical Research Letters*, 48(14), e2021GL092999. https://doi.org/10.1029/2021GL092999, 2021.
- Feng, Y., Menemenlis, D., Xue, H., Zhang, H., Carroll, D., Du, Y., and Wu, H.: Improved representation of river runoff in
 Estimating the Circulation and Climate of the Ocean Version 4 (ECCOv4) simulations: implementation, evaluation, and
 impacts to coastal plume regions, Geosci. Model Dev., 14, 1801-1819. https://doi.org/10.5194/gmd-14-1801-2021, 2021.
 - Fortin, V., Innocenti, S., Gaborit, É., Durnford, D., Keita, S., Bruxer, J., Boucher, M.-A., Harrigan, S., Zsoter, E., Dimitrijevic,
 M., Sévigny, C., O'Brien, N., and Gervasi, N.: Evaluation of continental-scale ensemble hydrological forecasts from
 Environment and Climate Change Canada: a comparison with forecasts from the Global Flood Awareness System
- 560 (GloFAS), EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-2950, https://doi.org/10.5194/egusphere-egu23-2950, 2023.
 - Frame, J. M., Kratzert, F., Klotz, D., Gauch, M., Shalev, G., Gilon, O., Qualls, L. M., Gupta, H. v, & Nearing, G. S. Deep learning rainfall-runoff predictions of extreme events. *Hydrol. Earth Syst. Sci.*, 26(13), 3377–3392. https://doi.org/10.5194/hess-26-3377-2022, 2022.
- 565 Garvine, R. W.: The impact of model configuration in studies of buoyant coastal discharge. Journal of Marine Research, 59, 193-225. <u>https://elischolar.library.yale.edu/journal of marine research/2389</u>, 2001.
 - Gauch, M., Mai, J., & Lin, J. (2021). The proper care and feeding of CAMELS: How limited training data affects streamflow prediction. *Environmental Modelling & Software*, *135*, 104926.
- Gleason, J. C., & Durand, T. M. Remote Sensing of River Discharge: A Review and a Framing for the Discipline. In *Remote* 570 Sensing (Vol. 12, Issue 7). https://doi.org/10.3390/rs12071107, 2020.
 - Graham, J. A., O'Dea, E., Holt, J., Polton, J., Hewitt, H. T., Furner, R., Guihou, K., Brereton, A., Arnold, A., Wakelin, S., Castillo Sanchez, J. M., and Mayorga Adame, C. G.: AMM15: a new high-resolution NEMO configuration for operational simulation of the European north-west shelf, Geosci. Model Dev., 11, 681-696. <u>https://doi.org/10.5194/gmd-11-681-2018</u>, 2018.
- 575 Greff, K., Srivastava, R. K., Koutník, J., Steunebrink, B. R., & Schmidhuber, J. LSTM: A Search Space Odyssey. IEEE <u>Transactions on Neural Networks and Learning Systems</u>, 28(10), 2222–2232. https://doi.org/10.1109/TNNLS.2016.2582924, 2017.

Grodsky, S. A., Vandemark, D., & Feng, H. Assessing coastal SMAP surface salinity accuracy and its application to monitoring Gulf of Maine circulation dynamics. Remote Sensing, 10(8). https://doi.org/10.3390/rs10081232, 2018.

- 580 <u>Gudmundsson, L., Do, H. X., Leonard, M., & Westra, S. The Global Streamflow Indices and Metadata Archive (GSIM) Part</u> <u>2: Quality control, time-series indices and homogeneity assessment. *Earth Syst. Sci. Data, 10*(2), 787–804. <u>https://doi.org/10.5194/essd-10-787-2018, 2018.</u></u>
 - Guillou, N., Chapalain, G., & Petton, S. Predicting sea surface salinity in a tidal estuary with machine learning. Oceanologia, 65(2), 318–332. https://doi.org/https://doi.org/10.1016/j.oceano.2022.07.007, 2023.

- 585 Harrigan, S., Zsoter, E., Alfieri, L., Prudhomme, C., Salamon, P., Wetterhall, F., Barnard, C., Cloke, H., and Pappenberger, F.: GloFAS-ERA5 operational global river discharge reanalysis 1979–present, Earth Syst. Sci. Data, 12, 2043-2060. https://doi.org/10.5194/essd-12-2043-2020, 2020.
 - Harrigan, S., Zsoter, E., Cloke, H., Salamon, P., and Prudhomme, C.: Daily ensemble river discharge reforecasts and real-time forecasts from the operational Global Flood Awareness System, Hydrol. Earth Syst. Sci., 27, 1-19. https://doi.org/10.5194/hess-27-1-2023, 2023.

615

Herzfeld, M.: Methods for freshwater riverine input into regional ocean models. Ocean Modelling, 90, 1-15. https://doi.org/10.1016/j.ocemod.2015.04.001, 2015.

Hetland R. D., and MacDonald, D. G.: Spreading in the near-field Merrimack River plume. Ocean Modelling, 21(1), 12-21. https://doi.org/10.1016/j.ocemod.2007.11.001, 2008.

595 Hirose, N.: Inverse estimation of empirical parameters used in a regional ocean circulation model, J. Oceanogr., 67, 323–336, https://doi.org/10.1007/s10872-011-0041-4, 2011.

Hochreiter, S., & Schmidhuber, J. Long Short-Term Memory. Neural Computation, 9(8), 1735–1780. https://doi.org/10.1162/neco.1997.9.8.1735, 1997.

Hordoir, R., Polcher, J., Brun-Cottan, JC. et al.: Towards a parametrization of river discharges into ocean general circulation

- 600 models: a closure through energy conservation. Clim Dyn 31, 891-908. <u>https://doi.org/10.1007/s00382-008-0416-4</u>, 2008.
 - Hu, R., Fang, F., Pain, C. C., & Navon, I. M. Rapid spatio-temporal flood prediction and uncertainty quantification using a deep learning method. *Journal of Hydrology*, *575*, 911–920. https://doi.org/10.1016/j.jhydrol.2019.05.087, 2019.
- Hunt, K. M. R., Matthews, G. R., Pappenberger, F., & Prudhomme, C. Using a long short-term memory (LSTM) neural
 network to boost river streamflow forecasts over the western United States. *Hydrol. Earth Syst. Sci.*, 26(21), 5449–5472.
 https://doi.org/10.5194/hess-26-5449-2022, 2022.
 - Jana, S., Gangopadhyay, A., and Chakraborty, A.: Impact of seasonal river input on the Bay of Bengal simulation, Cont. Shelf Res., 104, 45–62, https://doi.org/https://doi.org/10.1016/j.csr.2015.05.001, 2015.
- Kay, S., McEwan, R., and Ford, D.: North West European Shelf Production Centre NWSHELF MULTIYEAR BIO 004 011. 610 Available at https://catalogue.marine.copernicus.eu/documents/OUID/CMEMS-NWS-OUID-004-011.pdf. 2020 (last access: 28/07/2024).
 - Konapala, G., Kao, S.-C., Painter, S. L., & Lu, D. Machine learning assisted hybrid models can improve streamflow simulation in diverse catchments across the conterminous US. *Environmental Research Letters*, 15(10), 104022. https://doi.org/10.1088/1748-9326/aba927, 2020.
 - Kratzert, F., Klotz, D., Shalev, G., Klambauer, G., Hochreiter, S., & Nearing, G. S. Towards learning universal, regional, and local hydrological behaviors via machine learning applied to large-sample datasets. *Hydrology and Earth System Sciences*, 23(12), 5089–5110. https://doi.org/10.5194/hess-23-5089-2019, 2019.

Lacroix, G., Ruddick, K., Ozer, J., and Lancelot, C.: Modelling the impact of the Scheldt and Rhine/Meuse plumes on the

- 620 salinity distribution in Belgian waters (southern North Sea). Journal of Sea Research, 52(3), 149-163. https://doi.org/10.1016/j.seares.2004.01.003, 2004.
 - Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air-sea flux data set, Clim. Dyn., 33, 341– 364, https://doi.org/10.1007/s00382-008-0441-3, 2009.
- Lees, T., Buechel, M., Anderson, B., Slater, L., Reece, S., Coxon, G., & Dadson, S. J. Benchmarking data-driven rainfall–
 runoff models in Great Britain: A comparison of long short-term memory (LSTM)-based models with four lumped conceptual models. *Hydrology and Earth System Sciences*, 25(10), 5517–5534. https://doi.org/10.5194/hess-25-5517-2021, 2021.
 - Lenhart, H.-J., et al.: Predicting the consequences of nutrient reduction on the eutrophication status of the North Sea. Journal of Marine Systems, 81(1), 148-170. <u>https://doi.org/10.1016/j.jmarsys.2009.12.014</u>, 2010.
- 630 Li, J., Yuan, X., & Ji, P. Long-lead daily streamflow forecasting using Long Short-Term Memory model with different predictors. *Journal of Hydrology: Regional Studies*, 48, 101471. https://doi.org/10.1016/j.ejrh.2023.101471, 2023.
 - Lin, P., Ming, P., Beck, H.E., Yang, Y., Yamazaki, D., Frasson, R., David, C.H., Durand, M., Pevelsky, T.M., Allen, G.H., Gleason, C.J., Wood, E.F.: Global Reconstruction of Naturalized River Flows at 2.94 Million Reaches. Water Resources Research, 55(8), 6499-6516. <u>https://doi.org/10.1029/2019WR025287</u>, 2019.
- 635 Liu, Y., MacCready, P., Hickey, B.M., Dever, E.P., Kosro, P.M., and Banas, N.S.: Evaluation of a coastal ocean circulation model for the Columbia River plume in summer 2004. Journal of Geophysical Research: Oceans, 114, C00B04. <u>https://doi.org/10.1029/2008JC004929</u>, 2004.

Locarnini, R. A., et al.: World Ocean Atlas 2013, Volume 1: Temperature. NOAA Atlas NESDIS 73, 2013.

- Luppichini, M., Vailati, G., Fontana, L., & Bini, M. Machine learning models for river flow forecasting in small catchments.
 Scientific Reports, 14(1), 26740. https://doi.org/10.1038/s41598-024-78012-2, 2024.
 - Ma, K., Feng, D., Lawson, K., Tsai, W.-P., Liang, C., Huang, X., Sharma, A., & Shen, C. Transferring Hydrologic Data Across Continents – Leveraging Data-Rich Regions to Improve Hydrologic Prediction in Data-Sparse Regions. *Water Resources Research*, 57(5), e2020WR028600. https://doi.org/https://doi.org/10.1029/2020WR028600, 2021.
- MacCready, P., and Geyer, W. R.: Advances in Estuarine Physics. Annual Review of Marine Science, 2(1), 35-58. https://doi.org/10.1146/annurey-marine-120308-081015, 2010.
 - Maglietta, R., Verri, G., Saccotelli, L., de Lorenzis, A., Cherubini, C., Caccioppoli, R., Dimauro, G., & Coppini, G. Advancing estuarine box modeling: A novel hybrid machine learning and physics-based approach. *Environmental Modelling & Software*, 183, 106223. https://doi.org/https://doi.org/10.1016/j.envsoft.2024.106223, 2025.
- Maicu, F., Alessandri, J., Pinardi, N., Verri, G., Umgiesser, G., Lovo, S., Turolla, S., Paccagnella, T., & Valentini, A.
 Downscaling With an Unstructured Coastal-Ocean Model to the Goro Lagoon and the Po River Delta Branches. *Frontiers* in Marine Science, 8. https://www.frontiersin.org/journals/marine-science/articles/10.3389/fmars.2021.647781, 2021.

- Mankoff, K. D., Noël, B., Fettweis, X., Ahlstrøm, A. P., Colgan, W., Kondo, K., Langley, K., Sugiyama, S., van As, D., and Fausto, R. S.: Greenland liquid water discharge from 1958 through 2019, Earth Syst. Sci. Data, 12, 2811-2841. <u>https://doi.org/10.5194/essd-12-2811-2020</u>, 2020.
- 655 <u>Medina-Lopez, E. Machine learning and the end of atmospheric corrections: A comparison between high-resolution sea surface</u> salinity in coastal areas from top and bottom of atmosphere Sentinel-2 imagery. *Remote Sensing*, 12(18). https://doi.org/10.3390/RS12182924, 2020.
 - Medina-Lopez, E., & Ureña-Fuentes, L. High-resolution sea surface temperature and salinity in coastal areas worldwide from raw satellite data. *Remote Sensing*, *11*(19). https://doi.org/10.3390/rs11192191, 2019.
- 660 <u>Menezes, V. v. Statistical assessment of sea-surface salinity from SMAP: Arabian sea, bay of Bengal and a promising red sea</u> application. Remote Sensing, 12(3). https://doi.org/10.3390/rs12030447, 2020.
 - Milliman J. D., and Farnsworth, K. L.: River Discharge to the Coastal Ocean: A Global Synthesis. Cambridge: Cambridge University Press, 2011. <u>https://doi.org/10.1017/CBO9780511781247</u>, 2011.
 - Mishra, A. K., and Coulibaly, P.: Developments in hydrometric network design: A review. Review of Geophysics, 47(2).
- 665 <u>https://doi.org/10.1029/2007RG000243</u>, 2009.

- Moftakhari, H. R., Jay, D. A., and Talke, S. A.: Estimating river discharge using multiple-tide gauges distributed along a channel. Journal of Geophysical Research: Oceans, 121(4), 2078-2097. <u>https://doi.org/10.1002/2015JC010983</u>, 2016
- Moftakhari, H. R., Jay, D. A., Talke, S. A., Kukulka, T., and Bromirski, P. D.: A novel approach to flow estimation in tidal rivers. Water Resources Research, 49(8), 4817-4832. <u>https://doi.org/10.1002/wrcr.20363</u>, 2013.
- 670 Mouatadid, S., Adamowski, J. F., Tiwari, M. K., & Quilty, J. M. Coupling the maximum overlap discrete wavelet transform and long short-term memory networks for irrigation flow forecasting. *Agricultural Water Management*, 219, 72–85. https://doi.org/https://doi.org/10.1016/j.agwat.2019.03.045, 2019.
 - Muhebwa, A., Gleason, C. J., Feng, D., & Taneja, J. Improving Discharge Predictions in Ungauged Basins: Harnessing the Power of Disaggregated Data Modeling and Machine Learning. *Water Resources Research*, 60(9), e2024WR037122. https://doi.org/https://doi.org/10.1029/2024WR037122, 2024.
 - Nearing, G. S., Kratzert, F., Sampson, A. K., Pelissier, C. S., Klotz, D., Frame, J. M., et al. What Role Does Hydrological Science Play in the Age of Machine Learning? Water Resources Research, 57(3). https://doi.org/10.1029/2020WR028091, 2021.
 - Papa, F., Bala, S. K., Pandey, R. K., Durand, F., Gopalakrishna, V. v, Rahman, A., & Rossow, W. B. Ganga-Brahmaputra
- 680 river discharge from Jason-2 radar altimetry: An update to the long-term satellite-derived estimates of continental freshwater forcing flux into the Bay of Bengal. *Journal of Geophysical Research: Oceans*, 117(C11). https://doi.org/https://doi.org/10.1029/2012JC008158, 2012.
 - Pein, J., & Staneva, J. (2024): Eutrophication hotspots, nitrogen fluxes and climate impacts in estuarine ecosystems: A model study of the Odra estuary system. Ocean Dynamics 74, 335–354 (2024), doi:10.1007/s10236-024-01607-w

685 Pein, J., Eisele, A., Sanders, T., Daewel, U., Stanev, E.V., van Beusekom, J.E.E., Staneva, J., & Schrum, C. (2021): Seasonal Stratification and Biogeochemical Turnover in the Freshwater Reach of a Partially Mixed Dredged Estuary. Front. Mar. Sci. 8:623714, doi:10.3389/fmars.2021.623714

690

- Perry, G. D., Duffy, P. B., and Miller, N. L.: An extended data set of river discharges for validation of general circulation models. Journal of Geophysical Research: Atmospheres., 10(D16), 21339-21349. <u>https://doi.org/10.1029/96JD00932</u>, 1996.
- Polton, J., Harle, J., Holt, J., Katavouta, A., Partridge, D., Jardine, J., Wakelin, S., Rulent, J., Wise, A., Hutchinson, K., Byrne, D., Bruciaferri, D., O'Dea, E., De Dominicis, M., Mathiot, P., Coward, A., Yool, A., Palmiéri, J., Lessin, G., Mayorga-Adame, C. G., Le Guennec, V., Arnold, A., and Rousset, C.: Reproducible and relocatable regional ocean modelling: fundamentals and practices, Geosci. Model Dev., 16, 1481–1510, https://doi.org/10.5194/gmd-16-1481-2023, 2023.
- 695 Ponsoni, L., et al.: Greenlandic sea ice products with a focus on an updated operational forecast system. Frontiers in Marine Science, 10. <u>https://doi.org/10.3389/fmars.2023.979782</u>, 2023.
 - Qi, S., He, M., Bai, Z., Ding, Z., Sandhu, P., Zhou, Y., Namadi, P., Tom, B., Hoang, R., & Anderson, J. Multi-Location Emulation of a Process-Based Salinity Model Using Machine Learning. Water, 14(13). https://doi.org/10.3390/w14132030, 2022.
- 700 Qin, T., Fan, J., Zhang, X., et al.: Global Freshwater Discharge into the World's Oceans Reached a Record Low Over Past Nearly 70 Years. Available at Research Square <u>https://doi.org/10.21203/rs.3.rs-1402652/v1</u>, 2022.
 - Qiu, C., & Wan, Y. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. Water Resources Research, 49(9), 5804–5816. https://doi.org/10.1002/wrcr.20415, 2013.

Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., & Prabhat. Deep learning and process

- 705 <u>understanding for data-driven Earth system science</u>. *Nature*, *566*(7743), 195–204. https://doi.org/10.1038/s41586-019-0912-1, 2019.
 - Rasiya Koya, S., & Roy, T. Temporal Fusion Transformers for streamflow Prediction: Value of combining attention with recurrence. *Journal of Hydrology*, *637*, 131301. https://doi.org/10.1016/j.jhydrol.2024.131301, 2024.
- Rice, A., Whitney, M. M., Garvine, R. W., and Huq, P.: Energetics in Delaware Bay: Comparison of two box models with 710 observations. Journal of Marine Research, 66(6), 873-898. https://elischolar.library.yale.edu/journal of marine research/218, 2008.
 - Riggs, R. M., Allen, G. H., Wang, J., Pavelsky, T. M., Gleason, C. J., David, C. H., and Durand, M.: Extending global river gauge records using satellite observations, Environ. Res. Lett., 18, 64027, <u>https://doi.org/10.1088/1748-9326/acd407</u>, 2023.
- 715 <u>Saccotelli, L., Verri, G., de Lorenzis, A., Cherubini, C., Caccioppoli, R., Coppini, G., & Maglietta, R. Enhancing estuary salinity prediction: A Machine Learning and Deep Learning based approach. *Applied Computing and Geosciences*, 23, 100173. https://doi.org/10.1016/j.acags.2024.100173, 2024.</u>

Sakai, T., Omori, K., Oo, A. N., & Zaw, Y. N. Monitoring saline intrusion in the Ayeyarwady Delta, Myanmar, using data from the Sentinel-2 satellite mission. *Paddy and Water Environment*, *19*(2), 283–294. https://doi.org/10.1007/s10333-020-00837-0, 2021.

- Sampurno, J., Vallaeys, V., Ardianto, R., & Hanert, E. Integrated hydrodynamic and machine learning models for compound flooding prediction in a data-scarce estuarine delta. Nonlin. Processes Geophys., 29(3), 301–315. https://doi.org/10.5194/npg-29-301-2022, 2022.
- Schiller, R. V., and Kourafalou, V. H.: Modeling river plume dynamics with the HYbrid Coordinate Ocean Model. Ocean Modelling, 33(1), 101117. <u>https://doi.org/10.1016/j.ocemod.2009.12.005</u>, 2010.
 - Shi, X., Qin, T., Nie, H., Weng, B., and He, S.: Changes in Major Global River Discharges Directed into the Ocean. International Journal of Environmental Research and Public Health, 16(8). DOI: 10.3390/ijerph16081469, 2019.
 - Sobrinho, J., de Pablo, H., Campuzano, F., and Neves, R.: Coupling Rivers and Estuaries with an Ocean Model: An Improved Methodology. Water, 13(16), 2284. https://doi.org/10.3390/w13162284, 2021.
- 730 Sotillo, M. G., Campuzano, F., Guihou, K., Lorente, P., Olmedo, E., Matulka, A., Santos, F., Amo-Baladron. M.A., and Novellino, A.: River Freshwater Contribution in Operational Ocean Models along the European Atlantic Façade: Impact of a New River Discharge Forcing Data on the CMEMS IBI Regional Model Solution. Journal of Marine Science and Engineering, 9(4), 401. <u>https://doi.org/10.3390/jmse9040401</u>, 2021<u>a</u>.
 - Sotillo, M. G., Mourre, B., Mestres, M., Lorente, P., Aznar, R., García-León, M., Liste, M., Santana, A., Espino, M., and
- Álvarez, E.: Evaluation of the Operational CMEMS and Coastal Downstream Ocean Forecasting Services During the Storm Gloria (January 2020), https://www.frontiersin.org/articles/10.3389/fmars.2021.644525, 2021b.
 - Sprintall, J., and Tomczak, M.: Evidence of the barrier layer in the surface layer of the tropics. Journal of Geophysical Research: Oceans, 97(C5), 7305-7316. <u>https://doi.org/10.1029/92JC00407</u>, 1992.
- Staneva, J., Melet, A., Veitch, J., Matte, P. Solving Coastal Dynamics: Introduction to High Resolution Ocean Forecasting
 Services. *State Planet Discuss.*, 2024.
 - Storkey, D., Blaker, A. T., Mathiot, P., Megann, A., Aksenov, Y., Blockley, E. W., Calvert, D., Graham, T., Hewitt, H. T., Hyder, P., Kuhlbrodt, T., Rae, J. G. L., and Sinha, B.: UK Global Ocean GO6 and GO7: a traceable hierarchy of model resolutions, Geosci. Model Dev., 11, 3187–3213. <u>https://doi.org/10.5194/gmd-11-3187-2018</u>, 2018.
 - Sun, Q., Whitney, M. M., Bryan, F. O., and Tseng, Y.: A box model for representing estuarine physical processes in Earth system models. Ocean Modelling, 112, 139-153. <u>https://doi.org/10.1016/j.ocemod.2017.03.004</u>, 2017.
 - Sun, Q., Whitney, M. M., Bryan, F. O., and Tseng, Y.: Assessing the Skill of the Improved Treatment of Riverine Freshwater in the Community Earth System Model (CESM) Relative to a New Salinity Climatology. Journal of Advances in Modeling Earth Systems, 11(5), 1189-1206. <u>https://doi.org/10.1029/2018MS001349</u>, 2019.
- Suzuki, T., Yamazaki, D., Tsujino, H. et al.: A dataset of continental river discharge based on JRA-55 for use in a global ocean circulation model. Journal of Oceanography, 74, 421-429. https://doi.org/10.1007/s10872-017-0458-5, 2018.

745

720

-

- Talke S. A., and Jay, D. A.: Nineteenth Century North American and Pacific Tidal Data: Lost or Just Forgotten? Journal of Coastal Research. DOI: 10.2112/JCOASTRES-D-12-00181.1, 2013.
- Tang, S., Sun, F., Liu, W., Wang, H., Feng, Y., & Li, Z. Optimal Postprocessing Strategies With LSTM for Global StreamflowPredictioninUngaugedBasins.WaterResourcesResearch,59(7),e2022WR034352,2023.
- <u>Thao, N.T., Staneva, J., Grayek, S., Bonaduce, A., Hagemann, S., Nam, T.P., Kumar, R., & Rakovec, O. (2024): Impacts of extreme river discharge on coastal dynamics and environment: Insights from high-resolution modeling in the German Bight. Regional Studies in Marine Science, Vol 73, 103476, doi:10.1016/j.rsma.2024.103476</u>

- Thielen, J., Bartholmes, J., Ramos, M.-H., and de Roo, A.: The European Flood Alert System Part 1: Concept and development, Hydrol. Earth Syst. Sci., 13, 125-140. <u>https://doi.org/10.5194/hess-13-125-2009</u>, 2009.
 - Tonani, M., Sykes, P., King, R. R., McConnell, N., Péquignet, A.-C., O'Dea, E., Graham, J. A., Polton, J., and Siddorn, J.: The impact of a new high-resolution ocean model on the Met Office North-West European Shelf forecasting system, Ocean Sci., 15, 1133-1158. <u>https://doi.org/10.5194/os-15-1133-2019</u>, 2019.
- Tseng, Y., Bryan, F. O., and Whitney, M. M.: Impacts of the representation of riverine freshwater input in the community earth system model. Ocean Model., 105, 71-86. <u>https://doi.org/10.1029/2020MS002276</u>, 2016.
 - Tsujino, H., Urakawa, S., Nakano, H., Justin Small, R., et al. (2018). JRA-55 based surface dataset for driving ocean-sea-ice models (JRA55-do). Ocean Modelling, 130, 79-139. https://doi.org/10.1016/j.ocemod.2018.07.002
- Vallaeys, V., Lambrechts, J., Delandmeter, P., Pätsch, J., Spitzy, A., Hanert, E., & Deleersnijder, E. Understanding the circulation in the deep, micro-tidal and strongly stratified Congo River estuary. *Ocean Modelling*, *167*, 101890.
 https://doi.org/https://doi.org/10.1016/j.ocemod.2021.101890, 2021.
 - Vallaeys, V., Kärnä, T., Delandmeter, P., Lambrechts, J., Baptista, A. M., Deleersnijder, E., & Hanert, E. Discontinuous Galerkin modeling of the Columbia River's coupled estuary-plume dynamics. *Ocean Modelling*, *124*, 111–124. https://doi.org/https://doi.org/10.1016/j.ocemod.2018.02.004, 2018.
 - Vazquez-Cuervo, J., Fournier, S., Dzwonkowski, B., and Reager, J.: Intercomparison of In-Situ and Remote Sensing Salinity
- 775 Products in the Gulf of Mexico, a River-Influenced System. Remote Sensing, 10(10). https://doi.org/10.3390/rs10101590, 2018.
 - Verri, G., Barletta, I., Pinardi, N., Federico, I., Alessandri, J., & Coppini, G. Shelf slope, estuarine dynamics and river plumes in a z* vertical coordinate, unstructured grid model. *Ocean Modelling*, 184, 102235. https://doi.org/https://doi.org/10.1016/j.ocemod.2023.102235, 2023.
- 780 Verri, G., Mahmoudi Kurdistani, S., Coppini, G., and Valentini, A.: Recent Advances of a Box Model to Represent the Estuarine Dynamics: Time-Variable Estuary Length and Eddy Diffusivity. Journal of Advances in Modeling Earth Systems, 13(4), e2020MS002276, 2021.

- Verri, G., Pinardi, N., Bryan, F., Tseng, Y., Coppini, G., and Clementi, E.: A box model to represent estuarine dynamics in mesoscale resolution ocean models. Ocean Modelling, 148, 101587. <u>https://doi.org/10.1016/j.ocemod.2020.101587</u>, 2020.
- Verri, G., Pinardi, N., Oddo, P., Ciliberti, S. A., & Coppini, G. River runoff influences on the Central Mediterranean overturning circulation. Climate Dynamics, 50(5), 1675–1703. https://doi.org/10.1007/s00382-017-3715-9, 2018.
 - Vörösmarty, C. J., Fekete, B. M., and. Tucker, B. A.: Discharge compilation from The Global River Discharge (RivDIS) Project. Distributed Active Archive Center, Oak Ridge National Laboratory. PANGAEA. https://doi.org/10.1594/PANGAEA.859439, 1998.
- Vörösmarty, C. J., Green, P, Salisbury, J., and Lammers, R. B.: Global Water Resources: Vulnerability from Climate Change and Population Growth. Science, 289(5477), 284-288, DOI: 10.1126/science.289.5477.284, 2020.
 - Wahl, K. L., Thomas, W. O., and Hirsch, R. M.: The stream-gaging program of the U.S. Geological Survey. 1995.
- Yan, D., Wang, K., Qin, T. et al. (2019). A data set of global river networks and corresponding water resources zones divisions. Sci Data, 6, 219. https://doi.org/10.1038/s41597-019-0243-y, 1995.
- Yang, Y., Feng, D., Beck, H. E., Hu, W., Sengupta, A., Monache, L. D., Hartman, R., Lin, P., Shen, C., & Pan, M. Global Daily Discharge Estimation Based on Grid-Scale Long Short-Term Memory (LSTM) Model and River Routing. ESS Open Archive. https://doi.org/10.22541/essoar.169724927.73813721/v1, 2023.
- Yin, J., Stouffer, R.J., Spelman, M.J., and Griffies, S.M.: Evaluating the Uncertainty Induced by the Virtual Salt Flux
 Assumption in Climate Simulations and Future Projections. Journal of Climate, v23(1), 80-96.
 https://doi.org/10.1175/2009JCLI3084.1, 2010.
 - Young, E. F., and Holt, J. T.: Prediction and analysis of long-term variability of temperature and salinity in the Irish Sea. Journal of Geophysical Research: Oceans, 112(C1). <u>https://doi.org/10.1029/2005JC003386</u>, 2007.
 - Zhang, Y. J., Ye, F., Stanev, E. V., and Grashorn, S.: Seamless cross-scale modeling with SCHISM. Ocean Modelling, 102, 64-81. https://doi.org/10.1016/j.ocemod.2016.05.002, 2016.
 - Zhao, G., Bates, P., Neal, J., & Pang, B. Design flood estimation for global river networks based on machine learning models. *Hydrol. Earth Syst. Sci.*, 25(11), 5981–5999. https://doi.org/10.5194/hess-25-5981-2021, 2021.
 - Zweng, M. M, J. R. Reagan, J. I. Antonov, R. A. Locarnini, A. V. Mishonov, T. P. Boyer, H. E. Garcia, O.K. Baranova, D.R. Johnson, D. Seidov, M.M. Biddle. World Ocean Atlas 2013, Volume 2: Salinity. S. Levitus, Ed.; A. Mishonov, Technical
- 810 Ed.; NOAA Atlas NESDIS 74, 39 pp, 2013. Zweng, M., et al.: World Ocean Atlas 2013, Volume 2: Salinity. NOAA Atlas NESDIS 74, 2013. http://doi.org/10.7289/V55X26VD, 2013.

Competing interests

785

790

795

805

The contact author has declared that none of the authors has any competing interests.

Data and/or code availability

815 Data/code availability is not applicable to this article as no new data/code were created or analysed in this study.

Authors contribution

Pascal Matte: Conceptualization, Investigation, <u>Writing – original draft preparation</u>, Writing – review and editing. John Wilkin: Writing – review and editing. Joanna Staneva: Writing – review and editing.

Acknowledgements

820 The authors wish to thank Kristen Wilmer-Becker and members of the OceanPredict community who participated in the survey conducted in May 2023 on the status of implementation of river forcing in current OOFS, as well as three anonymous reviewers for their constructive comments on this manuscript.